arXiv:math/0610738v1 [math.DG] 24 Oct 2006

Metrics of Special Curvature with Symmetry

Brandon Dammerman
The Queen's College
University of Oxford

A thesis submitted for the degree of
Doctor of Philosophy

May 2004

Acknowledgements

Foremost, I would like to thank my supervisor, Dr. Andrew Dancer, for his patient, lucid, and insightful instruction. The results of Chapter 4 were obtained under the guidance of Prof. K. P. Tod, and I am very grateful to him for his help and instruction. Thanks also to my fellow graduate students Gil Cavalcanti, Marco Gualtieri, Steve Marshall, and Fred Witt for useful discussions.

Finally, I thank the Rhodes Trust for its generous funding and The Queen's College for granting me a lectureship in my final year.

Metrics of Special Curvature with Symmetry

Brandon Dammerman

Oxford University D. Phil. Thesis

May 2004

Abstract

Various curvature conditions are studied on metrics admitting a symmetry group. We begin by examining a method of diagonalizing cohomogeneity-one Einstein manifolds and determine when this method can and cannot be used. Examples, including the well-known Stenzel metrics, are discussed.

Next, we present a simplification of the Einstein condition on a compact four manifold with T^2-isometry to a system of second-order elliptic equations in two-variables with well-defined boundary conditions.

We then study the Einstein and extremal Kahler conditions on Kahler toric manifolds. After constructing explicitly new extremal Kahler and constant scalar curvature metrics on $\mathbb{P}(\mathcal{O}_{\mathbb{C}P^1\times\mathbb{C}P^1}\oplus\mathcal{O}_{\mathbb{C}P^1\times\mathbb{C}P^1}(1,-1))$, we demonstrate how these metrics can be obtained by continuously deforming the Fubini-Study metric on $\mathbb{C}P^3$. We also define a generalization of Kahler toric manifolds, which we call fiberwise Kahler toric manifolds, and construct new explicit extremal Kahler and constant scalar curvature metrics on both compact and non-compact manifolds in all even dimensions. We also calculate the Futaki invariant on manifolds of this type.

After describing an Hermitian non-Kahler analogue to fiberwise Kahler toric geometry, we construct constant scalar curvature Hermitian metrics with J-invariant Riemannian tensor. In dimension four, we write down explicitly new constant scalar curvature Hermitian metrics with J-invariant Ricci tensor on $\mathbb{C}P^1\times\mathbb{C}P^1$ and $\mathbb{C}P^2\sharp\overline{\mathbb{C}P^2}$.

Finally, we integrate the scalar curvature equation on a large class of cohomogeneity-one metrics.

Chapter 1

Introduction

Much of Riemannian geometry involves the search for distinguished metrics within the infinite-dimensional space of smooth metrics on a C^∞-differentiable manifold, particularly when the manifold is of dimension greater than or equal to four. Here, the term 'distinguished' refers to those metrics that exhibit additional structure or satisfy some curvature condition. For example, one could look for metrics of constant Riemannian or constant scalar curvature or whose Ricci tensor satisfies the Einstein condition. In particular, Einstein metrics are of interest to both geometers and physicists. Originally, the study of Einstein metrics was motivated by theoretical physics; space-time in the vacuum state is an Einstein manifold with zero Einstein constant. Furthermore, in the compact case Einstein metrics are critical points of the scalar curvature functional. Alternatively, one could study metrics with a compatible complex structure or metrics of special holonomy. Once a condition has been placed upon a metric, the mathematician tries to determine which manifolds admit metrics of that type, how abundant those metrics are on a given manifold, and if at all possible write down the metrics explicitly. In this thesis, we study the Einstein and constant scalar curvature conditions on manifolds of general type as well as on those admitting Hermitian and Kahler structures, in many cases deriving explicit solutions. We place particular emphasis on those cases in which the equations of the curvature conditions take a simplified form. Simplification is accomplished in two ways: we either demand that the manifold admit a large symmetry group or we use Kaluza-Klein bundle constructions. Both require an understanding of Riemannian submersions

The thesis begins with a brief review of the theory of Riemannian submersions using the formalism developed by O'Neill in [48]. Indeed, though the thesis considers various curvature conditions on metrics, some of which have compatible complex structures and some of which do not, all of the manifolds considered can be viewed as Riemannian submersions. Therefore, an understanding of the curvature conditions on a Riemannian submersion will be essential to every part of the thesis.

In Chapter 2, we also discuss a special kind of Riemannian submersion: the quotient by a group action. More precisely, we present the equations of the Einstein and constant scalar curvature conditions on a metric admitting *orthogonally transitive* isometries, i.e. those isometric group actions

in which the distribution orthogonal to the tangent space of the orbits of the action is integrable. Though not all of the metrics considered below admit orthogonally transitive isometries, many do. Isometry group actions on cohomogeneity-one manifolds are automatically orthogonally transitive, and cohomogeneity-one manifolds will be considered in Chapters 3, $5-8$, and 11. Manifolds of higher cohomogeneity under orthogonally transitive actions will be considered in Chapters $4-7$ and 9. In Chapter 2, we also present a rough topological classification of Einstein manifolds with an orthogonally transitive group action.

Most of the new material begins in Chapter 3 with our investigation of when it is possible to globally diagonalize an Einstein metric of cohomogeneity-one under the action of a compact, connected group G. The Einstein condition on a cohomogeneity-one manifold reduces to a system of ordinary differential equations, cf. [5]. While this is a great simplification of the general Einstein condition, as yet there exists no classification of cohomogeneity-one Einstein metrics even in the compact case. Indeed, such a classification appears, at the moment, to be only a distant goal. There are, however, many known examples of cohomogeneity-one Einstein metrics in all dimensions. To the author's knowledge, *all* of the known metrics can be written in diagonal form relative to a G-invariant basis. Many of these examples have been found either by looking at those group actions under which every cohomogeneity-one metric is automatically diagonalizable (called the monotypic case) or by demanding *ad hoc* that the metric be diagonal. In some special cases, it has been observed that the Einstein condition allows one to globally diagonalize certain cohomogeneity-one metrics in the non-monotypic case. For example, it is well-known to relativists that the Einstein condition on a four manifold with $SU(2)$-symmetry allows one to globally diagonalize the metric, though this result does not seem to appear in the pure mathematics literature. From that result, we devise a general procedure for determining when the Einstein condition can be used to globally diagonalize a cohomogeneity-one metric in the non-monotypic case. As an illustration, we show that this method works in the case of $SO(n+1)/SO(n-1)$, with $n \geq 3$, principal orbits (the Stenzel metrics found in [56] are of this type) but fails when the principal orbits are isomorphic to $SU(n+1)/U(n-1)$ or $SO(n_1+n_2+2)/(SO(n_1)\times SO(n_2))$ with both n_1 and n_2 greater than or equal to two.

After reviewing what is known about Einstein four manifolds admitting a large isometry group, we leave the cohomogeneity-one category to investigate compact Einstein four-manifolds with T^2-isometry in Chapter 4. Though it was known previously [39] that the torus action must be orthogonally transitive in the Einstein case, little else is known about metrics of this type in general. To date, the only known Einstein metrics of this type are the homogeneous metrics on S^4, $S^2\times S^2$, and $\mathbb{C}P^2$; the Hermitian-Einstein metric (the Page metric) on $\mathbb{C}P^2\sharp\overline{\mathbb{C}P^2}$; and the Kahler-Einstein metric (found by Siu in [55]) on $\mathbb{C}P^2\sharp 3\overline{\mathbb{C}P^2}$, though this metric has not been written down explicitly. After reviewing the topological classification of smooth torus actions on compact simply-connected four manifolds discovered by Orlik and Raymond in [49] and the *nuts* and *bolts* classification developed

by Gibbons and Hawking, we present our main result which provides a simplification of the Einstein equations. In general, the Einstein condition is an overdetermined system of partial differential equations. In the compact T^2-invariant case, we show that the system reduces to a set of four second-order elliptic equations in four independent functions in two variables together with a set of well-defined boundary conditions. This simplification is effected by making use of a natural complex structure on the two-dimensional quotient space M/T^2. While we are unable at this time to prove the existence of any new Einstein metrics, the reduction makes the system much more amenable to analytical methods for determining the existence and uniqueness of T^2-invariant Einstein metrics. It is our hope that our work will provide a useful first step in the eventual classification of such metrics. Finally, we end this section with a discussion of a simplified system in which the metric is diagonal. In this case, more can be said about the geometric structure of Einstein metrics.

Chapter 5 begins a study of curvature conditions on manifolds with large symmetry group in the presence of an integrable complex structure. Since the Einstein condition appears, at the moment, to be difficult to solve even in the relatively simple case of a four manifold with T^2-symmetry, there is little hope of arriving at a solution to the Einstein condition on a $2n$-dimensional manifold with T^n-symmetry. It seems appropriate therefore to require that the manifold admit additional structures which simplify the curvature equations. One such condition which is commonly studied is the reduction of the holonomy group to $U(n)$ from $SO(2n)$. In this case the manifold will be Kahler and the various curvature conditions can be expressed in much simplified form. When the manifold is Kahler toric, i.e. when there exists a T^n action preserving the Kahler structure, these conditions take yet simpler forms as we will see below.

Kahler toric manifolds are of interest because of, among other reasons, the presence of a Legendre transformation between complex coordinates and symplectic Darboux coordinates. A recent theorem of Abreu [1] allows one to fix a symplectic structure on a toric manifold and parameterize all of the compatible Kahler metrics via a symplectic potential. It is this property, which is absent in general Kahler geometry, that makes Kahler toric geometry so tractable. After reviewing the symplectic formalism, we discuss various curvature conditions on Kahler toric manifolds. In addition to the Kahler-Einstein condition, we also discuss the extremal Kahler condition (introduced by Calabi in [10]) and the more restrictive constant scalar curvature condition on Kahler toric manifolds, both of which have been widely studied by differential geometers. While the majority of Chapter 5 is a review of material necessary for later chapters, we do present a simplification of the scalar curvature equation on a Kahler toric manifold found by Abreu in [2].

Specific examples of Kahler toric manifolds with distinguished metrics, particularly in real dimensions four and six, are discussed in Chapter 6. A great deal is known about the existence of Kahler-Einstein toric manifolds (see [43]), though relatively few metrics are known explicitly. Before looking at explicit examples, we show how to integrate the Einstein equations to obtain the

Legendre transform of the Monge-Ampere equations of complex coordinates and write down the Futaki invariant, an obstruction to the existence of positive Kahler-Einstein metrics, in symplectic coordinates. We discuss the Kahler-Einstein metric on $\mathbb{C}P^2\sharp 3\overline{\mathbb{C}P^2}$ which has not been written down explicitly. We analyze this case in detail and propose an ansatz for the possible form of the Siu metric. Next, we proceed to write down the Kahler-Einstein metric found by Sakane on $\mathbb{P}(\mathcal{O}_{\mathbb{C}P^1\times\mathbb{C}P^1}\oplus\mathcal{O}_{\mathbb{C}P^1\times\mathbb{C}P^1}(1,-1))$ *explicitly* in symplectic coordinates. It is our hope that analyzing the form of known Kahler-Einstein toric metrics may lead to a method of writing down all of Kahler-Einstein toric metrics.

Recently, there has been a great deal of interest in finding and classifying constant scalar curvature metrics on compact Kahler manifolds. For example, a recent paper by Donaldson [18] discusses the question of finding constant scalar curvature metrics on Kahler toric manifolds and the relation to notions of stability. In the second part of Chapter 6, we look at the more general question of finding extremal Kahler metrics. We begin by reviewing the construction of the extremal Kahler metric conformal to the Hermitian-Einstein metric on $\mathbb{C}P^2\sharp\overline{\mathbb{C}P^2}$ and giving a novel interpretation of the symplectic potential of that extremal Kahler metric. We then find new *explicit* extremal Kahler metrics in dimension six. Specifically, we find a continuous two-parameter family of extremal Kahler metrics on $\mathbb{P}(\mathcal{O}_{\mathbb{C}P^1\times\mathbb{C}P^1}\oplus\mathcal{O}_{\mathbb{C}P^1\times\mathbb{C}P^1}(1,-1))$. Within this two-parameter family lie two one-parameter families of constant scalar curvature metrics. These two families intersect at one point; this point corresponds to the Kahler-Einstein metric found by Sakane. Furthermore, this two-parameter family, along with a one-parameter family of extremal Kahler metrics of nonconstant scalar curvature on a $\mathbb{C}P^2$-bundle over $\mathbb{C}P^1$, arise from a continuous deformation of the Fubini-Study metric on $\mathbb{C}P^3$.

Finally, we conclude Chapter 6 with a discussion of the connection, found by Derdzinski in [17], between Hermitian-Einstein metrics and extremal Kahler metrics in dimension four. Using a result of LeBrun in [40], we apply this connection to the Kahler toric case and exhibit the equations which must be satisfied for a non-Kahler Hermitian toric metric to be Einstein in dimension four.

In addition to the T^3-symmetry present on any six-dimensional Kahler toric manifold, all of the new extremal Kahler metrics we have constructed admit additional symmetries; they are all in fact of cohomogeneity-one under their full symmetry group. It seems natural, therefore, to look for explicit distinguished metrics on Kahler manifolds with large symmetry group. To begin such a study, we look, in Chapter 7, at Kahler manifolds of cohomogeneity-d under a compact semisimple group action. We see that the presence of a moment map and a complex structure requires that each homogeneous principal orbit, G/K, becomes a Riemannian submersion over a coadjoint orbit G/L with fibers L/K of dimension d. If we demand that each fiber be abelian (i.e. isomorphic to T^d) and restrict to the monotypic case, we recover a generalization of Kahler toric geometry which we refer to as fiberwise Kahler toric geometry. Metrics of this type can be thought of as Kahler

toric metrics with additional structure, and describing a fiberwise Kahler toric metric is equivalent to choosing a Kahler toric metric and specifying the values of a collection of constants.

Remarkably, metrics of the form described can be obtained by starting with very different geometric assumptions. In the second construction of fiberwise Kahler toric metrics we begin with a principal T^d-bundle over the product of Kahler-Einstein manifolds endowed with metrics as described in [58]. Letting $\hat{g}$ denote a family of such metrics depending on d variables, we construct a metric $g = \check{g} + \hat{g}$ where $\check{g}$ is a metric on the d-dimensional space of variables. Imposing a Kahler structure in a natural way, we retrieve metrics of the same form with the same curvature equations as those obtained by the previous construction; we also refer to these metrics as fiberwise Kahler toric. Indeed, the term fiberwise Kahler toric will be used to describe the form of the metric and not the particular construction. We end Chapter 7 by deriving the equations of the Einstein and extremal Kahler conditions.

Having defined fiberwise Kahler toric geometry, we specialize to the $d = 1$ case and begin our search for distinguished metrics. The Kahler-Einstein condition on fiberwise Kahler toric manifolds in the $d = 1$ case has been studied in great detail (see [15]). However, to the author's knowledge, the extremal Kahler and constant scalar curvature conditions have not been studied on manifolds of this type. In Chapter 8, we demonstrate how to integrate the scalar curvature equation on a fiberwise Kahler toric manifold in the $d = 1$ case. This integration will allow us to find extremal Kahler and constant scalar curvature Kahler metrics. After discussing smoothness conditions, we demonstrate that every fiberwise Kahler toric manifold admits at least one extremal Kahler metric and most admit a multi-parameter family of such metrics. This contrasts with the constant scalar curvature case in which there are examples of fiberwise Kahler toric metrics with $d = 1$ which are known to admit no constant scalar curvature metric even in real dimension four. We are able to prove the existence of a one-parameter family of constant scalar curvature metrics on most (we will make this statement more precise below) fiberwise Kahler toric metrics. This method is useful not only because it proves the existence of constant scalar curvature Kahler metrics, but also because it allows us to write the metrics down explicitly. Furthermore, it allows us to study how, on a given fiberwise Kahler toric manifold, the constant scalar curvature Kahler metrics sit within the larger family of extremal Kahler metrics. Conceivably, one could use this method to determine all of the Kahler classes that admit constant scalar curvature or extremal Kahler metrics and to study how those distinguished classes sit in the space of all Kahler classes. Unfortunately, we cannot answer all of these questions here. We do, however, go through many examples in detail. After a review of the four-dimensional case, we study the Kahler manifolds $\mathbb{P}(\mathcal{O}_{\mathbb{C}P^1\times\mathbb{C}P^1} \oplus \mathcal{O}_{\mathbb{C}P^1\times\mathbb{C}P^1}(p,q))$ and find extremal Kahler metrics for all values of p and q in $\mathbb{Z}$. Moreover, we see that constant scalar curvature metrics exists when $p > 0$ and $q < 0$ (or vice versa) or when $p = q = 0$ but not when both p and q are positive (or negative).

The case of fiberwise Kahler toric manifolds with $d > 1$ is more difficult to analyze because the curvature conditions are partial, instead of ordinary, differential equations. We cannot provide, therefore, new explicit examples of distinguished metrics. We do, however, integrate the equations of the Einstein condition to give an analogue of the Monge-Ampere equations as we did in the Kahler toric case. Also, we demonstrate how to express the Futaki invariant in the coordinates of fiberwise Kahler toric geometry.

In Chapter 10, we look for distinguished metrics which are Hermitian but non-Kahler. As in the second construction of $d = 1$ fiberwise Kahler toric metrics, we study metrics of the form $g = dt^2 + \hat{g}$ where $\hat{g}$ is a family of metrics on a principal S^1-bundle over the product of Kahler-Einstein manifolds. Again, we impose a natural integrable complex structure on the manifold. However, we no longer demand that the metric be Kahler; instead, we demand that the Riemannian curvature tensor be J-invariant, i.e. $R(\cdot,\cdot,\cdot,\cdot) = R(J\cdot, J\cdot, J\cdot, J\cdot)$. Note that the J-invariance of the Riemannian curvature tensor implies the J-invariance of the Ricci-tensor, i.e. $r(\cdot,\cdot) = r(J\cdot, J\cdot)$, and that the Riemannian and Ricci tensors of a Kahler manifold are automatically J-invariant. The Hermitian-Einstein condition on manifolds of this type was studied in [57]. In this chapter, we study the constant scalar curvature condition and demonstrate, after integrating the scalar curvature equation, a Hermitian constant scalar curvature metric always exists when each principal S^1-bundle is over a product of more than one Kahler-Einstein manifold. Therefore, most of the manifolds which do not admit constant scalar curvature Kahler metrics will admit non-Kahler constant scalar curvature Hermitian metrics with J-invariant Riemannian curvature tensor.

We place particular emphasis on the case in which the total manifold is of dimension four. In this case, the metric $\hat{g}$ is a metric on an S^1-bundle over $\mathbb{C}P^1$. In the compact case, the manifold is either $\mathbb{C}P^2$ or a Hirzebruch surface $\mathbb{P}(\mathcal{O}_{\mathbb{C}P^1} \oplus \mathcal{O}_{\mathbb{C}P^1}(q))$ where $q \in \mathbb{Z}$. It is well-known that, for $q \neq 0$, no Hirzebruch surface admits a constant scalar curvature Kahler metric. Therefore, it seems natural to widen the search for distinguished metrics. We propose that a natural generalization of both the constant scalar curvature Kahler and Hermitian-Einstein conditions, particularly in dimension four, is to look for constant scalar curvature Hermitian metrics with J-invariant Ricci tensor. In dimension four, non-Kahler Hermitian metrics with J-invariant Ricci tensor (but without necessarily being of constant scalar curvature) have been studied extensively by Apostolov and Gauduchon [4] and others. As such metrics are plentiful (some manifolds admit infinite dimensional families), one would like to impose addition restraints on the curvature tensor. An interesting additional constraint is the constant scalar curvature condition. To the author's knowledge, the only known non-Kahler Hermitian metric of constant scalar curvature with J-invariant Ricci tensor on a four manifold is the Page metric on $\mathbb{C}P^2 \sharp \overline{\mathbb{C}P^2}$. Here, we are able to construct new one-parameter families of such metrics on both $\mathbb{C}P^1 \times \mathbb{C}P^1$ and $\mathbb{C}P^2 \sharp \overline{\mathbb{C}P^2}$.

Both in the $d = 1$ fiberwise Kahler toric and the analogous Hermitian cases, we demonstrate how to integrate the scalar curvature equation. Chapter 11 investigates when it is possible to integrate the scalar curvature equation on a general cohomogeneity-one manifold which may not have a compatible complex structure. More precisely, we study manifolds of cohomogeneity-one under the action of a compact, connected group G with connected isotropy subgroup K. Assuming that the isotropy representation is composed of distinct $Ad(K)$-invariant summands (the monotypic assumption) and that there exists a summand of dimension one, we show how to integrate the scalar curvature when the Ricci tensor is invariant under the action of a natural rank-2 skew-symmetric map $\tilde{J} : TM \to TM$. We note that in the Einstein case, the Ricci tensor *must* be invariant under this map. The integration of the scalar curvature equation in the Einstein case amounts to finding a *conserved quantity* in the Hamiltonian formulation of the cohomogeneity-one Einstein equations. The discovery of such an integral may provide a first step in determining when the full integration of the Einstein equations is possible.

Chapter 2

Submersions, Isometries, and Curvature

This chapter is devoted to reviewing the theory of Riemannian submersions and deriving the curvature equations of a manifold admitting an orthogonally transitive group action. These curvature equations will be used extensively in all of the following chapters.

2.1 Riemannanian Submersions

In our exposition of the basic principles of Riemannian submersions, we follow closely the presentation found in [6]. Let (M, g) be a smooth Riemannian manifold without boundary and let $(B, \check{g})$ be a Riemannian manifold with piecewise smooth boundary. A submersion, $\pi : M \to B$, defines a tangent map $\pi_* : T_xM \to T_bB$ for each point x of M such that $\pi(x) = b$. Denote by $\mathcal{V}$ the kernel of the tangent map π_*, i.e. $\mathcal{V}$ is the distribution in TM such that, for any point x in M, $\mathcal{V}_x$ is the kernel of the map $\pi_* : T_xM \to T_bB$. In this thesis, the dimension of the distribution will be constant on all points of M which lie in the inverse image of the interior of B. However, at points in the inverse image of the boundary of B, the dimension of $\mathcal{V}$ may decrease. As M is endowed with a Riemannian metric, the orthogonal complement of $\mathcal{V}$ in TM, which will be denoted $\mathcal{H}$, is well-defined. Just as in the case of the *vertical* distribution $\mathcal{V}$, the *horizontal* distribution $\mathcal{H}$ is of constant dimension in the inverse image of the interior of B. As a final piece of notation, we denote by $\pi^{-1}(b)$ the inverse image of a point b in B. The tangent space to each point x in $\pi^{-1}(b)$ is simply the vertical space $\mathcal{V}_x$.

We say that π is a **Riemannian submersion** if it induces an isometry from $\mathcal{H}_x$ to T_bM for each point x such that $\pi(x) = b$. In this thesis, we frequently make use of a special type of Riemannian submersion known as the *quotient by an isometric action*. For $(M,\ g)$ a Riemannian manifold, let G be a closed subgroup of the isometry group of M. If we set $B = M/G$ and let $\check{g}$ be the quotient metric induced by g, then the projection from M to M/G is a Riemannian submersion. In this case,

each fiber, $\pi^{-1}(b)$, will be a homogeneous manifold isomorphic to G/K where K is the isotropy subgroup of G.

Given a Riemannian submersion, the various curvatures of M can be expressed in terms of the curvature of the base space B, the curvature of the spaces $\pi^{-1}(b)$, and two invariants [48]. The invariants, which following O'Neill we refer to as A and T, are defined in terms of the Levi-Civita connections of M and of the fibers $\pi^{-1}(b)$. Let ∇ be the Levi-Civita connection of g and let $\hat{\nabla}$ be the collection of Levi-Civita connections associated to the metrics $\hat{g}_b$ induced by g on each fiber $\pi^{-1}(b)$. Equivalently, one can define $\hat{\nabla}$ to be the vertical projection of ∇: for X and Y vector fields in TM, we define the connection of the fiber to be $\hat{\nabla}$ such that $\hat{\nabla}_X Y = \mathcal{V}\nabla_X Y$, where $\mathcal{V}X$ denotes the projection of the vector field X onto the vertical distribution (the horizontal projection is defined in an analogous way). We are now prepared to define the two invariants of a Riemannian submersion.

Definition 1. *For vector fields E and F on M, let T be the (2,1)-tensor field on M such that*

$$T_E F = \mathcal{H}\nabla_{\mathcal{V}E}\mathcal{V}F + \mathcal{V}\nabla_{\mathcal{V}E}\mathcal{H}F.$$

The various identities of the invariant T are collected in [6] and will not be reproduced here. We note, however, that T can be thought of as the collection of the second fundamental forms of the fibers $\pi^{-1}(b)$. The vertical distribution $\mathcal{V}$, since it is defined to be the kernel of a linear map of the tangent space TM, is integrable. This implies that each of the fibers $\pi^{-1}(b)$ is a submanifold of M. Therefore, one can define on each fiber the second fundamental form and it is easily checked that T restricted to a fiber is the second fundamental form of that fiber. Since the second fundamental form of a Riemannian submanifold measures the failure of that submanifold to be totally geodesic, the invariant T measures the failure of the fibers $\pi^{-1}(b)$ to be totally geodesic.

The second invariant associated to a Riemannian submersion is defined in a similar way.

Definition 2. *For vector fields E and F on M, let A be the (2,1)-tensor field on M such that*

$$A_E F = \mathcal{H}\nabla_{\mathcal{H}E}\mathcal{V}F + \mathcal{V}\nabla_{\mathcal{H}E}\mathcal{H}F.$$

The following proposition, found in [48], provides an interpretation of A.

Proposition 2.1. *For H and $\bar{H}$ horizontal vector fields, $A_H\bar{H} = \frac{1}{2}\mathcal{V}[H,\bar{H}]$.*

The tensor A can be interpreted as measuring the failure of the horizontal distribution to be integrable. Although the horizontal distribution is pointwise isometric to the tangent space of B, unless A vanishes, $\mathcal{H}$ does not define a submanifold isometric to B.

The curvature equations of a Riemannian submersion are presented in detail in [6]. For the sake of brevity, we will not reproduce the general curvatures equations of a Riemannian submersion here.

The Einstein condition on a generic Riemannian submersion is in general too difficult to solve. Therefore, additional constraints must be placed upon the submersion to make the equations more

tractable. As mentioned above, a common means of simplification is to look at the special case of a quotient by an isometric action. Since the fibers of this type of submersion are homogeneous manifolds, the terms representing their contribution to the curvature of the manifold are greatly simplified.

Either as an alternative or as a complementary means of simplification, one could set one of the two invariants A or T equal to zero. Note that setting them both equal to zero would mean that the Riemannian submersion is a simple product of the base and a fiber endowed with the product metric. Setting the invariant T equal to zero, since T is a generalization of the second fundamental form, is equivalent to demanding that the fibers be totally geodesic. For a review of submersions with totally geodesic fibers see [6] and the article by M. Wang in [41]. Setting the invariant A equal to zero is equivalent to demanding that the horizontal distribution be integrable. The condition that $A = 0$ is referred to as *orthogonal transitivity* by relativists. An orthogonally transitive action is known as a *surface-orthogonal* action elsewhere in the literature.

We will write down only the equations for the Ricci curvature of a quotient by an orthogonally transitive isometric action and refer the reader to [48] for their derivation.

Proposition 2.2. *Let g be the metric on M (we will sometimes set $g(\cdot,\cdot) = \langle\cdot,\cdot\rangle$), $\hat{g}$ the induced metric on the fibers, and $\check{g}$ the induced metric on the quotient space M/G and let r, $\hat{r}$ and $\check{r}$ to be the Ricci curvatures of those three metrics. For X and Y vertical vector fields and H and $\bar{H}$ horizontal vector fields, the Ricci curvature of M can be expressed by the following equations when the tensor A vanishes:*

$$r(X,Y) = \hat{r}(X,Y) - \langle N, T_X Y\rangle + (\tilde{\delta}T)(X,Y) \tag{2.1}$$

$$r(X,H) = \langle(\hat{\delta}T)X, H\rangle + \langle\nabla_X N, H\rangle \tag{2.2}$$

$$r(H,\bar{H}) = \check{r}(H,\bar{H}) - \langle TH, T\bar{H}\rangle + \frac{1}{2}(\langle\nabla_H N,\bar{H}\rangle + \langle\nabla_{\bar{H}} N, H\rangle) \tag{2.3}$$

where, for $\{Y_j\}$ an orthonormal basis of the vertical distribution and $\{H_j\}$ an orthonormal basis for the horizontal distribution, $N = \sum_j T_{Y_j} Y_j$, $\tilde{\delta}T(X,Y) = \sum_j \langle(\nabla_{H_j}T)_X Y, H_j\rangle$, $\hat{\delta}T = -\sum_j (\nabla_{Y_j}T)_{Y_j}$, and $\langle TH, T\bar{H}\rangle = \sum_j \langle T_{Y_j}H, T_{Y_j}\bar{H}\rangle$.

2.2 Orthogonally Transitive Isometries and the Einstein Condition

We now restrict our attention to a special kind of Riemannian submersion: the quotient by group action. We begin with a definition.

Definition 3. *Let (M^n, g) be a Riemannian manifold such that g is invariant under the action of group G. The manifold, M, is of* ***cohomogeneity-d*** *if and only if the difference, d, between the dimension of the manifold and the dimension of the principal orbits G/K is strictly less than n. (Note that while M is a smooth manifold, M/G is not smooth globally.)*

Here we have required that the dimension of the quotient space be strictly less than the dimension of the total manifold which implies that the dimension of the symmetry group G is at least one.

On manifolds of cohomogeneity-one the Riemannian submersion invariant A automatically vanishes since the one dimensional quotient manifold M/G must have zero curvature. The Einstein condition on a cohomogeneity-one manifold therefore becomes a system of differential equations in one variable. Going from the cohomogeneity-one case to manifolds with group actions of higher codimension, the complexity of the Einstein condition increases in three different ways. Firstly, the invariant A is no longer automatically equal to zero as the horizontal distribution orthogonal to the principal orbits is not necessarily integrable. Secondly, the invariant T depends on more than one variable, turning the Einstein condition into a system of partial differential equations. Finally, the Ricci tensor of the base manifold M/G is no longer zero in general. There may appear to be little hope in solving the Einstein condition in even the cohomogeneity-two case. However, if one considers a simplification of the higher cohomogeneity case, the Einstein condition becomes more tractable, particularly in low dimensions.

In light of the vanishing of the invariant A in the cohomogeneity-one case, a natural simplification of the general higher cohomogeneity case is to set the invariant A equal to zero. Below, after presenting certain properties of general higher cohomogeneity manifolds, we will restrict our attention to the case in which A vanishes.

Definition 4. *For (M, g) a cohomogeneity-d manifold, M is an* ***orthogonally transitive cohomogeneity-d manifold*** *if and only if the invariant A of the Riemannanian submersion $\pi : M \to M/G$ vanishes.*

2.2.1 Cohomogeneity-d Einstein Manifolds and Topology

Just as in the cohomogeneity one case, we have the following proposition.

Proposition 2.3. *For M an Einstein manifold of cohomogeneity-d, if the scalar curvature of M is negative then M is noncompact.*

Proof. If M is compact with negative Ricci curvature, Bochner's theorem states that all Killing vectors must vanish identically which would imply that the principal orbits are of dimension zero. □

Proposition 2.4. *For M a compact Einstein manifold of cohomogeneity-two or cohomogeneity-three, the scalar curvature is positive or the metric is flat.*

Proof. In light of the last proposition we need only consider the case in which the Einstein constant is equal to zero, i.e. $r = 0$. By Bochner's theorem, $s = 0$ implies that all Killing vectors are parallel [52] . It is easily seen that this implies the vanishing of the invariants A and T. By equations $9.36(a - c)$ of [6] and equations (2.1)-(2.3), this implies that $\check{r} = \hat{r} = 0$. As each principal orbit is homogeneous, $\hat{r} = 0$ implies that the Riemannian curvature of the principal orbit is zero. Finally, since M/G is of dimension two or three, $\check{r} = 0$ implies that $\check{R} = 0$. □

By Myer's Theorem, a compact Einstein manifold with positive scalar curvature must have finite fundamental group. Therefore, we have the following rough classification of cohomogeneity-two and -three Einstein manifolds.

For M a complete cohomogeneity-two or cohomogeneity-three Einstein manifold, one of the following holds:

- M is compact with positive scalar curvature and finite fundamental group
- M is compact and flat
- M is noncompact and has nonpositive scalar curvature.

This classification fails for manifolds of cohomogeneity greater than three (but which still have a symmetry group of dimension at least one), because on a manifold of dimension four or greater Ricci flatness is not equivalent to flatness. The simplest counter-example to extending the above classification to manifolds of higher cohomogeneity with symmetry group of at least one dimension is $S^1 \times K3$-surface. Endowed with its standard metric, this is a cohomogeneity-four Einstein manifold with zero scalar curvature which is not flat.

2.2.2 The Einstein Equations in the Orthogonally Transitive Case

Let M be a cohomogeneity-d manifold under the orthogonally transitive action of compact group G with isotropy subgroup K. The quotient by the group action yields the Riemannian submersion $\pi : M \to M/G$ where $\dim M/G = d$ as described above. A principal orbit, $\pi^{-1}(b)$, is the inverse image of a point b in the interior of the space M/G. Each principal orbit is isomorphic to G/K and of real codimension-d. Moreover, the union of the principal orbits forms an open dense set in M. On this set, the metric g, because it is invariant under an orthogonally transitive group action, can be written as

$$g = \check{g} + \hat{g}$$

where $\check{g}$ is the metric induced on the quotient space M/G and $\hat{g}$ is a d-parameter family of G-invariant metrics on the principal orbit.

To express the equations of the Einstein condition in the orthogonally transitive case, we set $\{X_i\}_{i=1,\dots,n-d}$ to be an invariant basis for each principal orbits and set

$$h_{ij} = g(X_i, X_j) = \hat{g}(X_i, X_j) = X_i^a X_{ja}.$$

For the moment, we will say no more about $\hat{g}$ other than to note that, depending on the isotropy representation, some of the metric elements h_{ij} will be zero automatically. We will make this precise in the following section.

Let $\{H_i\}_{i=1,\dots,d}$ be an *orthonormal* basis for the horizontal distribution and hence for the tangent space to M/G. We are now ready to determine the equations of the Einstein condition. The first equation we want to study is

$$r(X_i, X_j) = \hat{r}(X_i, X_j) - \langle N, T_{X_i} Y_i\rangle + (\tilde{\delta}T)(X_i, X_j) = \lambda g(X_i, X_j) = \lambda h_{ij}$$

where $g(\cdot,\cdot) = \langle\cdot,\cdot\rangle$ as above.

By the definition of T

$$T_{X_i} X_j = \mathcal{H}(X_i^a \nabla_a X_{jb}).$$

Note that we can choose the X_j and the H_i so that $[H_i, X_j] = 0$. Using this fact and the properties of tensor T, we have

$$\langle T_{X_i} X_j, H_k\rangle = -\frac{1}{2} H_a \nabla^a \langle X_i, X_j\rangle.$$

Therefore, we have demonstrated that

$$T_{X_i} X_j = -\frac{1}{2}\check{\nabla}_b h_{ij}. \tag{2.4}$$

The vector N, referred to as the *mean curvature vector* is defined to be the trace of the tensor T. That is,

$$N = h^{ij}(T_{X_i} X_j) = -\frac{1}{2} h^{ij}\check{\nabla}_b h_{ij} = -\frac{1}{2}\frac{\check{\nabla}_b(\det(h))}{\det(h)}. \tag{2.5}$$

Immediately, we can calculate one term of the Einstein condition,

$$\langle N, T_{X_i} X_j\rangle = \frac{1}{4}(\det(h))^{-1}\check{\nabla}^a \det(h)\check{\nabla}_a h_{ij} = \frac{1}{4}\frac{\langle\check{\nabla}\det(h), \check{\nabla} h_{ij}\rangle}{\det(h)}.$$

In order to find $R_{ab} X_i^a X_j^b = r(X_i, X_j)$ we must next calculate $\tilde{\delta}T$. Recalling the definition of $\tilde{\delta}T(X_i, X_j)$ and using the fact that T is a tensor field, the action of the Levi-Civita connection on tensors gives

$$\tilde{\delta}T(X_i, X_j) = \sum_k \{\langle\nabla_{H_k}(T_{X_i} X_j), H_k\rangle - \langle T_{\nabla_{H_k} X_i} X_j, H_k\rangle - \langle T_{\nabla_{H_k} X_j} X_i, H_k\rangle\}. \tag{2.6}$$

Having expressed $T_{X_i}X_j$ in terms of h_{ij}, it is straightforward to calculate (2.6). First, we note that, for any horizontal vector H, $\nabla_H(T_{X_i}X_j) = -\frac{1}{2}H^b\nabla_b\nabla_a h_{ij}$ and

$$\sum_k \langle \nabla_{H_k}(T_{X_i}X_j), H_k\rangle = -\frac{1}{2}\check{\nabla}^a\check{\nabla}_a h_{ij}.$$

The other elements of $\tilde{\delta}T$ are somewhat more difficult to compute. Recall that the X_i were chosen so that they commute with every horizontal vector field H, i.e. $\nabla_H X_i = \nabla_{X_i}H$. Furthermore, we note that $\nabla_H X_i = T_{X_i}H$ since A vanishes. Combining these results, we have

$$-\langle T_{\nabla_{H_k}X_i}X_j, H_k\rangle - \langle T_{\nabla_{H_k}X_j}X_i, H_k\rangle = \langle T_{X_j}H_k, \nabla_{H_k}X_i\rangle + \langle T_{X_i}H_k, \nabla_{H_k}X_j\rangle = 2\langle \nabla_{H_k}X_i, \nabla_{H_k}X_j\rangle.$$

From [45] we have

$$\nabla_H X_i = H^a\nabla_a X_{ib} = \frac{1}{2}h^{mk}(H^a\check{\nabla}_a h_{ki})X_{mb}$$

and therefore

$$\sum_{k=1}^{d}\langle \nabla_{H_k}X_i, \nabla_{H_k}X_j\rangle = \frac{1}{4}h^{kl}\check{\nabla}^a h_{ki}\check{\nabla}_a h_{lj} = -\frac{1}{4}h_{ik}\check{\nabla}^a h^{kl}\check{\nabla}_a h_{lj}.$$

The above results prove that

$$\tilde{\delta}T(X_i, X_j) = -\frac{1}{2}\check{\nabla}^a\check{\nabla}_a h_{ij} + \frac{1}{2}h^{kl}\check{\nabla}^a h_{ki}\check{\nabla}_a h_{lj}.$$

Finally we find that

$$r(X_i, X_j) = -\frac{1}{2}\check{\nabla}^a\check{\nabla}_a h_{ij} - \frac{1}{4}(\det(h))^{-1}\check{\nabla}^a\det(h)\check{\nabla}_a h_{ij} + \frac{1}{2}h^{kl}\check{\nabla}^a h_{ki}\check{\nabla}_a h_{lj} + \hat{r}(X_i, X_j) \tag{2.7}$$

or, equivalently,

$$r(X_i, X_j) = -\frac{1}{2}h_{ik}(\det(h))^{-\frac{1}{2}}\check{\nabla}^a((\det(h))^{\frac{1}{2}}h^{kl}\check{\nabla}_a h_{ij}) + \hat{r}(X_i, X_j). \tag{2.8}$$

We now turn to the second equation of the Einstein condition $r(X, H) = 0$ where $X = X_i$ and $H = H_j$ for some i and j. Because the O'Neill tensor A vanishes, we have that

$$r(X, H) = \langle(\hat{\delta}T)X, H\rangle + \langle\nabla_X N, H\rangle.$$

From [6], we see that

$$\hat{\delta}T = -h^{ij}(\nabla_{X_i}T)_{X_j}.$$

We consider first the term $\langle\nabla_X N, H\rangle$. Noting the N is G-invariant we see that

$$\check{\nabla}_X\langle N, H\rangle = 0 \Rightarrow \langle\nabla_X N, H\rangle + \langle N, \nabla_X H\rangle = 0.$$

Because we have chosen X to be a Killing vector field commuting with H, we have that

$$\mathcal{H}\nabla_X H = \mathcal{H}\nabla_H X = A_H X = 0.$$

Therefore,

$$\check{\nabla}_X \langle N, H \rangle = 0$$

automatically.

The term $\langle (\hat{\delta} T) X, H \rangle$ is more difficult to analyze and we follow [5] closely. In [5], $H = \frac{\partial}{\partial t}$ as they are considering the cohomogeneity-one case. However, we may take H to be any horizontal vector field without affecting the calculation. We have that

$$r(X,H) = -h^{ij} \langle (\nabla_{X_i} T)_{X_j} X, H \rangle = h^{ij} \langle (\nabla_{X_i} T)_{X_j} H, X \rangle$$

$$= h^{ij} \langle \hat{\nabla}_{X_i} (T_{X_j} H), X \rangle - h^{ij} \langle T_{\hat{\nabla}_{X_i} X_j} H, X \rangle.$$

From the symmetries of T, we see that

$$h^{ij} \langle T_{\hat{\nabla}_{X_i} X_j} H, X \rangle = -h^{ij} \langle T_{\hat{\nabla}_{X_j} X_i} H, X \rangle.$$

For all Killing vectors, Z, we know by a well-known formula that

$$\langle \hat{\nabla}_{X_i} X_j, Z \rangle = \frac{1}{2} (\langle [X_i, X_j], Z \rangle + \langle [X_i, Z], X_j \rangle + \langle [X_j, Z], X_i \rangle). \tag{2.9}$$

This implies (see [5]) that

$$-h^{ij} \langle T_{\hat{\nabla}_{X_i} X_j} H, X \rangle = -\mathrm{tr}(\mathrm{ad}_{\mathfrak{g}}(T_H X)).$$

This term is equal to zero if and only if G is unimodular. Since we have assumed that G is compact and compact groups are unimodular, we deduce that

$$r(X,H) = h^{ij} \langle \hat{\nabla}_{X_i} (T_{X_j} H), X \rangle$$

$$= h^{ij} X_j \langle T_{X_i} H, X \rangle - h^{ij} \langle \hat{\nabla}_{X_i} X, T_{X_j} H \rangle.$$

The second term is equal to zero because T is symmetric and $\hat{\nabla} X$ is anti-symmetric because X is a Killing vector field [5]. The first term becomes

$$h^{ij} X_j \langle T_{X_i} H, X \rangle = \frac{1}{2} h^{ij} H X_i \langle X_i, X_j \rangle = \frac{1}{2} (h^{ij} H \langle \hat{\nabla}_{X_i} X_j, X \rangle + h^{ij} H \langle \hat{\nabla}_{X_i} X, X_j \rangle).$$

The second term vanishes because $\hat{\nabla} X$ is anti-symmetric. The first term can be simplified using equation (2.9) to give

$$r(X,H) = \frac{1}{2} h^{ij} H \langle X_i, [X_j, X] \rangle. \tag{2.10}$$

Since this equation holds for all $X = X_k$ and H horizontal, equation $r(X,H) = 0$ of the Einstein condition implies that

$$h^{ij} \hat{\nabla} \langle X_i, [X_j, X_k] \rangle = 0 \tag{2.11}$$

for all X_k. Aside from the Ricci curvature of the base manifold, the only other terms in the Ricci tensor restricted to the horizontal distribution are $\langle TH, T\bar{H} \rangle$ and $\langle \nabla_H N, \bar{H} \rangle + \langle \nabla_{\bar{H}} N, H \rangle$ where H

and $\bar{H}$ are again arbitrary horizontal vector fields. Recalling the $N = -\frac{1}{2}h^{ij}\nabla_b h_{ij}$, a straightforward calculation shows that

$$\langle \nabla_H N, \bar{H}\rangle = -\frac{1}{2}h^{ij}H^a\bar{H}^b\nabla_a\nabla_b h_{ij} - \frac{1}{2}H^a\nabla_a h^{ij}\bar{H}^b\nabla_b h_{ij}.$$

Because both the matrix J_{ij} and the Hessian $\nabla_a\nabla_b$ are symmetric, $\langle \nabla_H N, \bar{H}\rangle = \langle \nabla_{\bar{H}} N, H\rangle$. Due to the vanishing of A and the choice of killing vectors and for $(Y_k)_{k=1,\dots,n-d}$ an orthonormal basis of the vertical distribution,

$$\langle TH, T\bar{H}\rangle = \sum_k \langle T_{Y_k}H, T_{Y_k}\bar{H}\rangle = \sum_k \langle \nabla_H Y_k, \nabla_{\bar{H}} Y_k\rangle = -\frac{1}{4}H^a\nabla_a h^{ij}\bar{H}^b\nabla_b h_{ij}.$$

Finally, we can express the Ricci curvature restricted to the horizontal distribution as

$$r_{ab}|_{\mathcal{H}} = \check{r}_{ab} - \frac{1}{2}h^{ij}\check{\nabla}_a\check{\nabla}_b h_{ij} - \frac{1}{4}\check{\nabla}_a h^{ij}\check{\nabla}_b h_{ij},$$

where $\check{r}$ is the Ricci curvature of the quotient manifold M/G endowed with its induced metric $\check{g}$.

With the above calculations, we have proven the following theorem.

Theorem 2.5. *Let (M, g) be a cohomogeneity-d manifold under the orthogonally transitive action of compact connected group G. Let $\{X_i\}_{i=1,\dots,n-d}$ be an invariant basis for the vertical distribution where $h_{ij} = g(X_i, X_j)$ and let $\check{g}$ be the metric induced on the quotient space of M/G. The Einstein condition on such a manifold is equivalent to*

- $-\frac{1}{2}\check{\triangle}h_{ij} - \frac{1}{4}\frac{\langle\check{\nabla}\det(h),\check{\nabla}h_{ij}\rangle}{\det(h)} - \frac{1}{2}h_{ik}\langle\check{\nabla}h^{kl},\check{\nabla}h_{lj}\rangle + \hat{r}(X_i, X_j) = \lambda h_{ij}$, *for all* $1 \le i, j \le n-d$,
- $h^{ij}\check{\nabla}\langle X_i, [X_j, X_k]\rangle = 0$, *for all* $1 \le k \le n-d$, *and*
- $\check{r}_{ab} - \frac{1}{2}h^{ij}\check{\nabla}_a\check{\nabla}_b h_{ij} - \frac{1}{4}\check{\nabla}_a h^{ij}\check{\nabla}_b h_{ij} = \lambda\check{g}_{ab}$.

For now, we will not make an effort to express $\hat{r}$ more explicitly. In the sequel, we will evaluate this term on a case by case basis. For a more general discussion in the cohomogeneity-one case, which readily extends to the orthogonally transitive cohomogeneity-d case, we refer the reader to [58].

2.2.3 Einstein Equations in the Monotypic Case

In this section, we give a more thorough discussion of the form of the metric $\hat{g}$. We then restrict our attention to the monotypic case in which the metric and the equations of the Einstein condition take a particularly simple form.

We again take (M, g) to be an orthogonally transitive cohomogeneity-d manifold under the action of compact connected group G. Let K be the isotropy subgroup of G, i.e.

$$K = \{g \in G : g(x) = x\}$$

for some $x \in \pi^{-1}(b)$ with b in the interior of the quotient space M/G. As mentioned in the previous section, the metric can be written as $g = \check{g} + \hat{g}$ where $\hat{g}$ is a d-parameter family of G-invariant homogeneous metrics on $G/K \cong \pi^{-1}(b)$. To describe the space of such homogeneous metrics, we identify the tangent space to G/K with an $\mathrm{Ad}(K)$-invariant subspace of the Lie algebra of G.

Let $\mathfrak{p}$ be an $Ad(K)$-invariant subspace of $\mathfrak{g}$ such that

$$\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}.$$

We choose once and for all a space $\mathfrak{p}$. though we note that the subspace $\mathfrak{p}$ is not unique. That subspace can be identified with $T_x(G/K)$ for $x \in G/K$ [6]. Under this identification, choosing a G-invariant metric on G/K is equivalent to choosing an $\mathrm{Ad}(K)$-invariant scalar product on $\mathfrak{p}$.

One can further decompose $\mathfrak{p}$ into $\mathrm{Ad}(K)$-invariant summands

$$\mathfrak{p} = \mathfrak{p}_1 \oplus ... \oplus \mathfrak{p}_m$$

where each $\mathfrak{p}_i$ is irreducible. Set $\dim\mathfrak{p}_i = d_i$ for all $1 \leq i \leq m$. We fix once and for all a bi-invariant metric Q on $\mathfrak{g}$ such that $Q(\mathfrak{k}, \mathfrak{p}) = 0$ and $Q(\mathfrak{p}_i, \mathfrak{p}_j) = 0$ for $i \neq j$. Any other G-invariant metric on G/K can be written as $Q(L\cdot, \cdot)$ where L is a positive-definite symmetric $\mathrm{Ad}(K)$-equivariant linear map [37]. That is,

$$L : \mathfrak{p} \to \mathfrak{p}.$$

The space of $\mathrm{Ad}(K)$-invariant scalar products is parameterized by the matrices L. Finally we let $\{Y_j^i\}_{j=1,...,d_i}$ be an orthonormal basis for $\mathfrak{p}_i$ relative to the metric Q.

We now specialize to the monotypic case.

Definition 5. *The isotropy representation is called* ***monotypic*** *if all of the summands, $\mathfrak{p}_i$, are pairwise inequivalent.*

By Schur's lemma, we see that any $\mathrm{Ad}(K)$-equivariant linear map, L, leaves each of the $\mathfrak{p}_i$ invariant in the monotypic case [5]. Therefore, the d-parameter family of G-invariant metrics $\hat{g}$ can be written as

$$\hat{g}(\cdot, \cdot) = \sum_{i=1}^{m} A_i Q(\cdot, \cdot)|_{\mathfrak{p}_i} \tag{2.12}$$

where the A_i are positive functions on M/G. The resulting cohomogeneity-d metric preserves the bi-invariance in the monotypic case. This gives the following proposition:

Proposition 2.6. *In the monotypic case, for Y_j^i a vertical vector field and H a horizontal vector field on an orthogonally transitive cohomogeneity-d manifold M, $r(Y_j^i, H) = 0$.*

Proof. The proof follows from the proof of the analogous statement in the cohomogeneity one case [5]. To see this we note that the second equation in (2.5) becomes

$$\sum_{i=1}^{m}\sum_{j=1}^{d_i} A_i^{-1}\check{\nabla}\langle Y_j^i,[Y_j^i,Y_l^k]\rangle = 0$$

for all Y_l^k. Because Q is bi-invariant, equation (2.12) implies that the above equation vanishes automatically.

□

A straightforward calculation using the equations of theorem (2.5) shows that for $(Y_1^1,...,Y_{d_m}^m)$ a basis for the principal orbits, the Einstein condition on an orthogonally transitive cohomogeneity-d manifold in the monotypic case becomes

$$\hat{r}(Y_i^k,Y_i^k) - \frac{1}{2}\check{\triangle}A_i + \frac{1}{2}\frac{\langle\check{\nabla}A_i,\check{\nabla}A_i\rangle}{A_i} - \frac{1}{4}\sum_{j=1}^{m} d_j\frac{\langle\check{\nabla}A_i,\check{\nabla}A_j\rangle}{A_j} = \lambda A_i \tag{2.13}$$

$$\check{r}_{ab} - \frac{1}{2}\sum_{j=1}^{m} d_j\frac{\check{\nabla}_a\check{\nabla}_b A_j}{A_j} + \frac{1}{4}\sum_{j=1}^{m} d_j\frac{\check{\nabla}_a A_j\check{\nabla}_b A_j}{A_j^2} = \lambda\check{g}_{ab} \tag{2.14}$$

for all $1\le i\le m$ and $1\le k\le d_i$ where $\hat{r}$ is the Ricci curvature of each principal orbit, $\check{r}$ is the Ricci curvature of the base space, $\check{\nabla}_a\check{\nabla}_b$ is the Hessian with respect to the base, $\check{\triangle} = \check{\nabla}^a\check{\nabla}_a$ is the Laplacian of $\check{g}$, and λ is the Einstein constant of the manifold M. If $(n-d) = \sum_{i=1}^{r} d_i$ then the trace of (2.13) can be written as

$$-\frac{\check{\triangle}\left(A_1^{\frac{d_1}{2}}\cdot...A_m^{\frac{d_m}{2}}\right)}{A_1^{\frac{d_1}{2}}\cdot...A_m^{\frac{d_m}{2}}} = (n-d)\lambda - \hat{s}.$$

This equation, or rather its non-monotypic version, will be very useful when we consider torus principal orbits since $\hat{r}=0$ in that case. Although torus principal orbits are not contained in the monotypic case, it can easily be seen from the calculations in [5] that $r(X,H)=0$ in that case as well.

Finally, the scalar curvature, S, of metric g, can be written as

$$S = \hat{S} + \check{S} + \sum_{j=1}^{m} d_j\frac{\check{\triangle}A_j}{A_j} + \frac{3}{4}\sum_{j=1}^{m} d_j\frac{\langle\check{\nabla}A_j,\check{\nabla}A_j\rangle}{A_j^2} - \frac{1}{4}\sum_{j=1}^{m}\sum_{k=1}^{m}\frac{\langle\check{\nabla}A_j,\check{\nabla}A_k\rangle}{A_jA_k}$$

where $\hat{S}$ denotes the scalar curvature of the fibers and $\check{S}$ denotes the scalar curvature of the quotient metric $\check{g}$.

Chapter 3

Diagonalizability of Cohomogeneity-One Einstein Metrics

On a manifold of cohomogeneity-one under the action of a group G, we saw above that a metric is automatically diagonal in the monotypic case. However, when two or more of the irreducible summands in the isotropy representation are equivalent, the metric need not be diagonal. The simplest example is a cohomogeneity-one metric whose principal orbits are copies of T^n. In this case, the metric is 'maximally' non-diagonalizable in the sense that all the non-diagonal entries of the fiber portion of a smooth metric may be non-zero. While a 'generic' cohomogeneity-one metric is not automatically diagonalizable in the non-monotypic case, it is known that, in certain situations, the equations of the Einstein condition allow the metric to be diagonalized globally. More precisely, in these cases it is one equation in particular of the Einstein condition which allows the metric to be globally diagonalized.

The Equation:

Let (M, g) be an n-dimensional manifold of cohomogeneity-one under the action of a compact connected group G. In the cohomogeneity-one case, the quotient space, M/G, is one-dimensional which implies that the horizontal distribution of the Riemannian submersion $\pi : M \to M/G$ is also of dimension one. As a one-dimensional distribution is automatically integrable, the O'Neill tensor A vanishes. Since $A \equiv 0$, the O'Neill formulae imply that

$$r\left(X, \frac{\partial}{\partial t}\right) = \left\langle (\delta T)X, \frac{\partial}{\partial t} \right\rangle + \left\langle \nabla_X N, \frac{\partial}{\partial t} \right\rangle$$

where X is any vertical vector field and $\frac{\partial}{\partial t}$ is the lift of the unit length vector field on the one-dimensional quotient space M/G.

Because the action of the group G is orthogonally transitive the metric can be written as

$$g = dt^2 + \hat{g} = dt^2 + g_t.$$

As above, let $\{X_i\}_{i=1,\dots,n-1}$ be an invariant basis of Killing vector fields for each of the principal orbits $\pi^{-1}(b) \cong G/K$, and let $h_{ij} = \langle X_i, X_j \rangle = \hat{g}(X_i, X_j)$. Equation (2.12) tells us that

$$r\left(X, \frac{\partial}{\partial t}\right) = \frac{1}{2} h^{ij} \frac{\partial}{\partial t} \langle X_i, [X_j, X] \rangle$$

for all vertical vector fields X. Because the vertical and horizontal distributions are orthogonal, the Einstein condition requires that $r\left(X, \frac{\partial}{\partial t}\right) = 0$ for all X. In the Einstein case, therefore, the above equation becomes

$$h^{ij} \frac{\partial}{\partial t} \langle X_i, [X_j, X] \rangle = 0 \tag{3.1}$$

Note that equation (3.1) is a necessary condition for a metric to be Einstein but it is far from sufficient.

Recall from proposition 2.6 that equation (3.1) is automatically satisfied in the monotypic case. This equation is also automatically satisfied in the abelian case in which the principal orbits are copies of T^n. However, in the non-monotypic case, there are many principal orbit types in which equation (3.1) is not automatically satisfied.

For certain principal orbit types in the non-monotypic case, it has been noted that the Einstein condition (specifically equation (3.1) of the Einstein condition) forces metrics to be diagonal. Thus far only a few individual examples of this phenomenon are known and, to the author's knowledge, no general study has been made of when the Einstein condition can, or cannot, be used to diagonalize a metric. In this chapter, we will review in detail an example of a cohomogeneity-one Einstein four manifold well known to relativists and will treat that example in the more mathematical formalism due to Berard-Bergery. Then, we will begin a more general study of the diagonalizability of Einstein metrics in the non-monotypic case.

3.1 Einstein Four Manifolds with SU(2) Principal Orbits

A Bianchi IX metric is a cohomogeneity-one metric on a four manifold in which the principal orbits are copies of $SU(2)$. That is, the group is $G = SU(2)$, the isotropy subgroup is $K = \{Id\}$, and the Lie algebra $\mathfrak{g}$ decomposes as

$$\mathfrak{g} = \mathfrak{p}_1 \oplus \mathfrak{p}_2 \oplus \mathfrak{p}_3$$

where each $\mathfrak{p}_i$ is a trivial one-dimensional subspace of $\mathfrak{g}$. Take $X_i \in \mathfrak{p}_i$ to be a basis for each principal orbit satisfying the standard Lie algebra relations for $\mathfrak{su}(2)$, namely

$$[X_1, X_2] = X_3, \quad [X_2, X_3] = X_1, \quad [X_3, X_1] = X_2.$$

Defining $h_{ij} = \langle X_i, X_j \rangle$ as above we see that

$$g_t = (h_{ij}).$$

Now, since all of the $\mathfrak{p}_i$ are equivalent one-dimensional trivial subspaces, none of the off-diagonal entries in (h_{ij}) need be zero automatically.

Equation (3.1) becomes the system

$$h^{ij} \frac{\partial}{\partial t} \langle X_i, [X_j, X_k] \rangle = 0$$

for all k.

We now make the important observation that given a point x of the quotient space one can, without loss of generality, diagonalize the metric at that point. That is,

$$h_{ij} \mid_{t=x} = h^{ij} \mid_{t=x} = 0$$

for $i \neq j$. Given this condition, equation (3.1) becomes the following system of three equations

$$(h^{33} - h^{22}) \dot{h}_{23} \mid_x = 0 \tag{3.2}$$

$$(h^{22} - h^{11}) \dot{h}_{12} \mid_x = 0 \tag{3.3}$$

$$(h^{11} - h^{33}) \dot{h}_{13} \mid_x = 0 \tag{3.4}$$

where $\dot{h}_{ij} = \frac{\partial}{\partial t} h_{ij}$. If we require that $SU(2)$ is the full isometry group (e.g. the manifold does not have $U(2)$ symmetry) we can further assume without loss of generality that the h_{ii} are distinct. Equation (3.1) then implies that

$$\dot{h}_{ij} \mid_x = 0 \tag{3.5}$$

for $i \neq j$.

Therefore, at point x the metric and the first derivative of the metric are diagonal. By differentiating equation (3.1) and evaluating at the point x, we conclude in the same way that the second derivatives of the non-diagonal entries must be zero at x as well. By continued differentiation of equation (3.1), we see that all of the derivatives of the non-diagonal entries must be zero at x. It follows immediately that the non-diagonal entries of the metric must be zero everywhere.

We have proven that equation (3.1) of the Einstein condition allows us to globally diagonalize the metric on a cohomogeneity-one four manifold with $SU(2)$ symmetry.

One may ask what happens if the metric is of cohomogeneity-one under a larger isometry group than $SU(2)$, say $U(2)$? If the four manifold is of cohomogeneity-one under the action of $G = U(2)$ then it has isotropy subgroup $K = U(1)$ and the Lie algebra decomposition becomes

$$\mathfrak{u}(2) = \mathfrak{u}(1) \oplus \mathfrak{p}_1 \oplus \mathfrak{p}_2$$

where $\dim\mathfrak{p}_1 = 1$ and $\dim\mathfrak{p}_2 = 2$. These two summands are clearly inequivalent and we are therefore in the monotypic case and the metric is automatically diagonal.

Having worked through a particular example in detail, we now wish to look at the question of diagonalization of Einstein metrics in the non-monotypic case in greater generality.

3.2 The Program

From the example of an Einstein four manifold of cohomogeneity-one under the action of $SU(2)$, we can derive a program for determining the degree to which a cohomogeneity-one Einstein metric can be diagonalized. We here describe, in general terms, the steps of this program. In the following sections, we will put this program into more detailed practice.

- **Step 1:** After specifying the principal orbit type G/K choose a decomposition $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ as above. Determine what non-diagonal elements may exist in the metric. Recall that in the monotypic case there are no non-diagonal elements to the metric.

- **Step 2:** Assume the number of non-diagonal entries in the metric g_t is greater than zero. We can diagonalize the metric g_t at a point x in M/G.

- **Step 3:** Having diagonalized the metric at a point, we evaluate equation (3.1) of the Einstein condition. At the point x, because the metric is diagonal, we will obtain equations involving only the derivatives of the non-diagonal entries and not the derivatives of the diagonal elements.

- **Step 4:** The equations obtained in Step 3 will allow us to deduce that certain linear combinations of the derivatives of the non-diagonal elements are equal to zero at x.

- **Step 5:** Finally, we deduce that those linear combinations of non-diagonal elements whose derivatives vanish, must be zero on the whole of M. If sufficiently many non-diagonal elements vanish, then we will be able to globally diagonalize the metric.

These five steps provide a broad outline of our method, the details of each step will be made clear below.

3.3 Preliminary Calculations

Let (M^n, g) be a cohomogeneity-one Riemannian manifold under the action of a compact connected Lie group G. Each principal orbit is of the form G/K where K is the isotropy subgroup. We choose an $Ad(K)$-invariant complement $\mathfrak{p}$ to $\mathfrak{k}$ such that decomposition of $\mathfrak{g}$ such that

$$\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p} = \mathfrak{k} \oplus \mathfrak{p}_1 \oplus ... \oplus \mathfrak{p}_m \tag{3.6}$$

where each $\mathfrak{p}_\mathfrak{i}$ is an irreducible summand of dimension d_i. As above, we fix once and for all a bi-invariant metric Q on $\mathfrak{g}$ such that $Q(\mathfrak{k}, \mathfrak{p}) = 0$ and $Q(\mathfrak{p}_i, \mathfrak{p}_j) = 0$ for all $i \neq j$.

Let $\{Y_j^i\}_{j=1,...,d_i}$ be an orthonormal basis relative to the metric Q for each $\mathfrak{p}_i$. Because of the bi-invariance of Q we have that

$$Q(Y_j^i, [Y_k^l, Y_r^s]) = -Q(Y_r^s, [Y_k^l, Y_j^i]) \tag{3.7}$$

for all i, j, k, l, r, and s.

Again, if all of the summands $\mathfrak{p}_i$ are distinct then the metric, g, is automatically diagonal and equation $r(X, \frac{\partial}{\partial t}) = 0$ is automatically satisfied. Therefore, we will be here interested exclusively in the non-monotypic case.

Non-diagonal elements of the metric on a cohomogeneity-one manifold will arise if and only if $\mathfrak{p}_i \simeq \mathfrak{p}_j$ for two or more of the $\mathfrak{p}_\mathfrak{i}$. By reordering if necessary, we can write the decomposition of $\mathfrak{g}$ as

$$\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}_1 \oplus ... \oplus \mathfrak{p}_r \oplus \mathfrak{p}_{r+1} \oplus ... \oplus \mathfrak{p}_m$$

where, for all $1 \leq i \leq r$, $\mathfrak{p}_\mathfrak{i} \simeq \mathfrak{p}_j$ for some $1 \leq j \leq r$ and, for $r+1 \leq k \leq m$, $\mathfrak{p}_k$ is distinct from all other summands. In this case, we can write the metric on the fibers as

$$g_t(\cdot, \cdot) = g_t(\cdot, \cdot) \mid_{\mathfrak{p}_1 \oplus ... \oplus \mathfrak{p}_r} \overset{\perp}{\bigoplus_{i=r+1,...,m}} A_i(t) Q(\cdot, \cdot) \mid_{\mathfrak{p}_i} .$$

From the bi-invariance of Q we have that

$$g_t(Y_j^i, [Y_j^i, X]) = 0 \tag{3.8}$$

for all $r+1 \leq i \leq m$, $1 \leq j \leq d_i$, and $X \in \mathfrak{g}$.

Therefore, using equation (3.8) and letting $\{Z_j\}$ be an invariant basis for $\mathfrak{p}_1 \oplus ... \oplus \mathfrak{p}_r$ with $h_{ij} = \langle Z_i, Z_j \rangle = g_t(Z_i, Z_j)$, we see that equation (3.1) simplifies to

$$h^{ij} \frac{\partial}{\partial t} \langle Z_i, [Z_j, X] \rangle = 0 \tag{3.9}$$

for $X \in \mathfrak{p}$. (Note that in equation (3.1) the h_{ij} are the metric functions relative to basis $\{X_i\}$ of $\mathfrak{p} = \mathfrak{p}_1 \oplus ... \oplus \mathfrak{p}_m$ while in equation (3.9) the h_{ij} are the metric functions relative to basis $\{Z_i\}$ of $\mathfrak{p}_1 \oplus ... \oplus \mathfrak{p}_r$. In the sequel, we work with an explicit basis which will eliminate any ambiguity.)

Below, we make use of equation (3.9) extensively. To get an idea for how it is applied, let us first consider the simplest type of cohomogeneity-one metric which is not automatically diagonalizable.

3.4 The Case of Two Equivalent Summands

Given the decomposition of equation (3.6), consider the case in which

- $\mathfrak{p}_1 \simeq \mathfrak{p}_2$ and
- all other $\mathfrak{p}_i$, for $i \geq 3$, are distinct.

As in the previous section, we take Q to be a fixed bi-invariant metric on $\mathfrak{g}$ and take $\{Y_j^i\}_{j=1,\ldots,d_i}$ to be an orthonormal basis for each $Q\mid_{\mathfrak{p}_i}$. In this case, $d_1 = d_2 = d$

Now, the fiber metric g_t, following the discussion above, can be written as

$$g_t(\cdot,\cdot) = g_t(\cdot,\cdot)\mid_{\mathfrak{p}_1\oplus\mathfrak{p}_2} \overset{\perp}{\bigoplus_{i=3,\ldots,m}} A_i(t)Q(\cdot,\cdot)\mid_{\mathfrak{p}_i}$$

Before proceeding, we will need a more precise expression for $g_t(\cdot,\cdot)\mid_{\mathfrak{p}_1\oplus\mathfrak{p}_2}$ in terms of the bi-invariant form $Q\mid_{\mathfrak{p}_1\oplus\mathfrak{p}_2}$. To do so, we follow closely the work of Kerr in [37] who studied Einstein metrics on homogeneous manifolds whose isotropy representations contain equivalent summands.

We want to parameterize $g_t\mid_{\mathfrak{p}_1\oplus\mathfrak{p}_2}$in terms of $Q\mid_{\mathfrak{p}_1\oplus\mathfrak{p}_2}$ and functions of t. Put another way, we want to find the number of distinct non-diagonal entries a general fiber metric $g_t\mid_{\mathfrak{p}_1\oplus\mathfrak{p}_2}$ has. To answer this question, we must determine the dimension of the space of intertwining maps $I : \mathfrak{p}_1 \to \mathfrak{p}_2$.

Any fiber metric restricted to $\mathfrak{p}_1 \oplus \mathfrak{p}_2$ can be written as

$$g_t(\cdot,\cdot)\mid_{\mathfrak{p}_1\oplus\mathfrak{p}_2} = Q(h\cdot,\cdot)\mid_{\mathfrak{p}_1\oplus\mathfrak{p}_2}$$

where h is a positive definite $Ad(K)$-invariant linear map $h : \mathfrak{p}_1 \oplus \mathfrak{p}_2 \to \mathfrak{p}_1 \oplus \mathfrak{p}_2$. To parameterize, the space of $Ad(K)$-equivariant maps between the pair of equivalent representations, we first note that $\mathfrak{p}_1$ and $\mathfrak{p}_2$ are irreducible as real representations. However, the complexification of, say, $\mathfrak{p}_1$ is not necessarily irreducible. After complexifying $\mathfrak{p}_1$, there are three possibilities

- $\mathfrak{p}_1$ is **orthogonal** iff $\mathfrak{p}_1 \otimes \mathbb{C}$ is irreducible
- $\mathfrak{p}_1$ is **unitary** iff $\mathfrak{p}_1 \otimes \mathbb{C} = \varphi \oplus \overline{\varphi}$ and φ is not equivalent to $\overline{\varphi}$
- $\mathfrak{p}_1$ is **symplectic** iff $\mathfrak{p}_1 \otimes \mathbb{C} = \varphi \oplus \overline{\varphi}$ and $\varphi \simeq \overline{\varphi}$.

The space of intertwining maps is 1-dimensional in the orthogonal case, 2-dimensional in the unitary case, and 4-dimensional in the symplectic case [37]. Below, we consider each of these cases separately.

3.4.1 The Orthogonal Case

When $\mathfrak{p}_1$ and $\mathfrak{p}_2$ are orthogonal representations, we have that

$$g_t|_{\mathfrak{p}_1 \oplus \mathfrak{p}_2} = \begin{pmatrix} h_{11}\mathrm{Id}_d & h_{12}\mathrm{Id}_d \\ h_{12}\mathrm{Id}_d & h_{22}\mathrm{Id}_d \end{pmatrix}$$

relative to basis $(Y_1^1, ..., Y_d^1, Y_1^2, ..., Y_d^2)$ where $h_{ij} = h_{ij}(t)$ [37]. Therefore, in the orthogonal case, there is only one non-diagonal function h_{12}. When can we conclude that $h_{12} = 0$ on an Einstein metric?

First, we calculate that

$$r\left(X, \frac{\partial}{\partial t}\right) = \frac{1}{2} h^{ij} \frac{\partial}{\partial t} \left(\sum_{k=1}^{d} \langle Y_k^i, [Y_k^j, X] \rangle \right)$$

for all $X \in \mathfrak{p}$. Here we have used the Einstein summation convention and have noted that $g_t(Y_k^i, Y_l^j) = 0$ for $k \neq l$.

Having accomplished Step One of the program by determining the number of independent non-diagonal elements of the metric, we progress to Step Two and diagonalize the metric at point $x \in M$. That is, $h_{12} \mid_x = 0$. This implies that

$$r\left(X, \frac{\partial}{\partial t}\right) \mid_x = \frac{1}{2} h^{ii} \frac{\partial}{\partial t} \left(\sum_{k=1}^{d} \langle Y_k^i, [Y_k^i, X] \rangle \right) \mid_x \tag{3.10}$$

$$= \frac{1}{2} h^{11} \frac{\partial}{\partial t} \left(\sum_{k=1}^{d} \langle Y_k^1, [Y_k^1, X] \rangle \right) \mid_x + \frac{1}{2} h^{22} \frac{\partial}{\partial t} \left(\sum_{k=1}^{d} \langle Y_k^2, [Y_k^2, X] \rangle \right) \mid_x .$$

Recalling the form of the fiber metric determined above and the bi-invariance of Q, we have that

$$\langle Y_k^1, [Y_k^1, X] \rangle = g_t(Y_k^1, [Y_k^1, X]) = h_{11} Q(Y_k^1, [Y_k^1, X]) + h_{12} Q(Y_k^2, [Y_k^1, X])$$

$$= h_{12} Q(Y_k^2, [Y_k^1, X]) = -h_{12} Q([Y_k^1, Y_k^2], X).$$

Performing a similar calculation for $\langle Y_k^2, [Y_k^2, X] \rangle$, we have demonstrated that

$$r\left(X, \frac{\partial}{\partial t}\right) \mid_x = \frac{1}{2} (h^{22} - h^{11}) \dot{h}_{12} \sum_{k=1}^{d} Q([Y_k^1, Y_k^2], X) \mid_x .$$

Step Three tells us to evaluate equation $r\left(X, \frac{\partial}{\partial t}\right) = 0$ of the Einstein condition at point x. In the Einstein case, we have that

$$(h^{22} - h^{11}) \dot{h}_{12} \sum_{k=1}^{d} Q([Y_k^1, Y_k^2], X) \mid_x = 0$$

We can assume that there exists a point x at which $h^{11} \neq h^{22}$. This could only fail if $h = h_{11} Id_{2d}$ which implies that $g_t = h_{11} Q$. In that case, diagonalizability is automatic. We have

$$\left[\sum_{k=1}^{d} Q([Y_k^1, Y_k^2], X) \right] \dot{h}_{12} \mid_x = 0. \tag{3.11}$$

If $\sum_{k=1}^{d} Q([Y_k^1, Y_k^2], X) \neq 0$ for some $X \in \mathfrak{p}$ we deduce that $\dot{h}_{12} \mid_x = 0$. However, since $h_{12} \mid_x = \dot{h}_{12} \mid_x = 0$ we can then conclude that $h_{12} \equiv 0$ and the metric is diagonal.

Because Q is nondegenerate, $\sum_{k=1}^{d} Q([Y_k^1, Y_k^2], X)$ is nonzero for some X if and only if the vector $\sum_{k=1}^{d} [Y_k^1, Y_k^2]$ is nonzero. We have proven the following theorem.

Theorem 3.1. *Let (M, g) be an Einstein manifold of cohomogeneity-one under the action of a compact, connected group G such that the isotropy representation has two equivalent* ***orthogonal*** *summands of dimension d, with all others distinct. The Einstein metric can be diagonalized if*

$$\sum_{k=1}^{d} [Y_k^1, Y_k^2]_{\mathfrak{p}} \neq 0 \tag{3.12}$$

where the Y_j^i are as defined above.

We now review two examples in which we can apply this theorem. In the first example, equation (3.1) of the Einstein condition can be used to diagonalize a cohomogeneity-one metric but in the second equation (3.1) cannot be so used.

Example: $G/K \cong SO(n+1)/SO(n-1)$

The Stiefel manifold is the homogeneous space $V_2(\mathbb{R}^2) = SO(n+1)/SO(n-1)$ for $n \geq 3$. Let (M, g) be a cohomogeneity-one manifold with principal orbits isomorphic to Stiefel manifolds. In this case, the Lie algebra $\mathfrak{g} = \mathfrak{so}(n+1)$ and $\mathfrak{k} = \mathfrak{so}(n-1)$. Letting E_{ij} the matrix with 1 in the ijth entry and -1 in the jith entry, we first embed K in G so that

$$\mathfrak{k} \cong \begin{pmatrix} 0 & 0 \\ 0 & \mathfrak{so}(n-1) \end{pmatrix} \subset \mathfrak{so}(n+1).$$

The isotropy representation is

$$\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}_1 \oplus \mathfrak{p}_2 \oplus \mathfrak{p}_3$$

and we can take $\mathfrak{p}_j = \text{span}\{E_{j,2+i} \mid 1 \leq i \leq n-1\}$ for $j = 1, 2$, and $\mathfrak{p}_3 = \text{span}\{E_{12}\}$. This decomposition is not unique since $\mathfrak{p}_1 \simeq \mathfrak{p}_2$. Therefore, the cohomogeneity-one metric g is not automatically diagonal.

To determine whether g can be diagonalized in the Einstein case, we must calculate the vector $\sum_{k=1}^{n-1} [E_{1,2+k}, E_{2,2+k}]_{\mathfrak{p}}$. It is straightforward to determine that

$$\sum_{k=1}^{n-1} [E_{1,2+k}, E_{2,2+k}]_{\mathfrak{p}} = -(n-1)E_{12} \neq 0.$$

Therefore, by theorem (3.1), any cohomogeneity-one Einstein metric with principal orbit isomorphic to $SO(n+1)/SO(n-1)$ can be diagonalized globally.

Stenzel in [56] constructed Ricci-flat Kahler metrics on the tangent bundle to the n-sphere, TS^n. These metrics are of cohomogeneity-one with principal orbits isomorphic to Stiefel manifolds (the $n = 2$ case is equivalent to the Eguchi-Hanson metric and the $n = 3$ case was found by Candelas and de la Ossa in [11]). Dancer and Strachan in [12] constructed Einstein metrics of negative Einstein constant on manifolds of this type. In each of these cases, the metrics were found by restricting to the diagonal case. Up to now it was unclear whether or not one could construct non-diagonal cohomogeneity-one Einstein manifolds of this type which were inequivalent to the diagonal ones. The above proves that the diagonal case is equivalent to the general case and requiring the metric to be diagonal need not be considered an additional constraint.

Example: $G/K \cong SO(n_1 + n_2 + 2)/(SO(n_1) \times SO(n_2))$

We now consider a generalization of a Stiefel manifold. Consider (M, g) a cohomogeneity-one manifold with principal orbits isomorphic to $SO(n_1 + n_2 + 2)/(SO(n_1) \times SO(n_2))$ where n_1 and n_2 are positive integers.

Letting E_{ij} be the matrix with 1 in the ijth entry and -1 in the jith entry. We can embed $SO(n_1) \times SO(n_2)$ in $SO(n_1 + n_2 + 2)$ so that on the Lie algebra level we have

$$\mathfrak{k} \cong \begin{pmatrix} 0 & 0 & 0 \\ 0 & \mathfrak{so}(n_1) & 0 \\ 0 & 0 & \mathfrak{so}(n_2) \end{pmatrix} \subset \mathfrak{so}(n_1 + n_2 + 2).$$

The isotropy representation becomes

$$\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}_1 \oplus \mathfrak{p}_2 \oplus \mathfrak{p}_3 \oplus \mathfrak{p}_4 \oplus \mathfrak{p}_5 \oplus \mathfrak{p}_6$$

where $\mathfrak{g} = \mathfrak{so}(n_1 + n_2 + 1)$, $\mathfrak{k} = \mathfrak{so}(n_1) \oplus \mathfrak{so}(n_2)$, $\mathfrak{p}_1 \cong (\mathbb{R}^{n_1} \otimes \mathbb{R}^{n_2})$, $\mathfrak{p}_2 \simeq \mathfrak{p}_3 \cong \mathbb{R}^{n_1}$, $\mathfrak{p}_4 \simeq \mathfrak{p}_5 \cong \mathbb{R}^{n_2}$, and $\mathfrak{p}_6 \cong \mathbb{R}$.

We see that $\mathfrak{p}_2 = \text{span}\{E_{1,2+j} \mid 1 \leq j \leq n_1\}$, $\mathfrak{p}_3 = \text{span}\{E_{2,2+j} \mid 1 \leq j \leq n_1\}$, $\mathfrak{p}_4 = \text{span}\{E_{1,2+n_1+j} \mid 1 \leq j \leq n_2\}$, $\mathfrak{p}_5 = \text{span}\{E_{2,2+n_1+j} \mid 1 \leq j \leq n_2\}$, and $\mathfrak{p}_6 = \text{span}\{E_{12}\}$. Clearly, we have two sets of equivalent orthogonal summands. This implies that a general cohomogeneity-one metric of this type will have two non-diagonal functions. To determine whether g can be diagonalized globally in the Einstein case, we first calculate that

$$\sum_{i=1}^{n_1} [E_{1,2+j}, E_{2,2+j}]_{\mathfrak{p}} = -n_1 E_{12}$$

and

$$\sum_{i=1}^{n_2} [E_{1,2+n_1+j}, E_{2,2+n_1+j}]_{\mathfrak{p}} = -n_2 E_{12}.$$

Given this calculation, it is straightforward that the Einstein condition cannot be diagonalized globally by the method described above. This follows because we have only one condition which is not sufficient to force the two non-diagonal functions to be zero.

3.4.2 The Unitary Case

When $\mathfrak{p}_1$ and $\mathfrak{p}_2$ are equivalent unitary representations, we have that $g_t(\cdot,\cdot) \mid_{\mathfrak{p}_1\oplus\mathfrak{p}_2} = Q((h+f)\cdot,\cdot) \mid_{\mathfrak{p}_1\oplus\mathfrak{p}_2}$ where h and f are of the form

$$h = \begin{pmatrix} h_{11}\mathrm{Id}_d & h_{12}\mathrm{Id}_d \\ h_{12}\mathrm{Id}_d & h_{22}\mathrm{Id}_d \end{pmatrix}$$

and

$$f = \begin{pmatrix} & & & f\mathrm{Id}_{\frac{d}{2}} \\ & & -f\mathrm{Id}_{\frac{d}{2}} & \\ & -f\mathrm{Id}_{\frac{d}{2}} & & \\ f\mathrm{Id}_{\frac{d}{2}} & & & \end{pmatrix}$$

relative to a basis $(Y_1^1,...,Y_d^1,Y_1^2,...,Y_d^2)$ where $h_{ij}=h_{ij}(t)$ and $f=f(t)$. Therefore, in the unitary case there are two non-diagonal functions. When can we conclude that $h_{12}\equiv 0$ and $f\equiv 0$ in the Einstein case?

As above, we diagonalize the metric at a point $x\in M$. That is, $h_{12}\mid_x = 0$ and $f\mid_x = 0$. At x, we calculate that

$$r\left(X,\frac{\partial}{\partial t}\right)\mid_x = \frac{1}{2}h^{ii}\frac{\partial}{\partial t}\left(\sum_{k=1}^{d}\langle Y_k^i,[Y_k^i,X]\rangle\right)\mid_x$$

$$= \frac{1}{2}h^{11}\frac{\partial}{\partial t}\left(\sum_{k=1}^{d}\langle Y_k^1,[Y_k^1,X]\rangle\right)\mid_x + \frac{1}{2}h^{22}\frac{\partial}{\partial t}\left(\sum_{k=1}^{d}\langle Y_k^2,[Y_k^2,X]\rangle\right)\mid_x$$

for all $X\in\mathfrak{p}$. Noting the bi-invariance of Q we calculate that

$$\langle Y_k^1,[Y_k^1,X]\rangle = g_t(Y_k^1,[Y_k^1,X]) = -h_{12}Q([Y_k^1,Y_k^2],X) - fQ([Y_k^1,Y_{k+\frac{d}{2}}^2],X)$$

for $1\le k\le\frac{d}{2}$ and

$$\langle Y_k^1,[Y_k^1,X]\rangle = g_t(Y_k^1,[Y_k^1,X]) = -h_{12}Q([Y_k^1,Y_k^2],X) + fQ([Y_k^1,Y_{k-\frac{d}{2}}^2],X)$$

for $\frac{d}{2}\le k\le d$ and for all $X\in\mathfrak{p}$.

After calculating $\langle Y_k^2,[Y_k^2,X]\rangle$, we have that

$$r(X,\frac{\partial}{\partial t})\mid_x = \frac{1}{2}(h^{22}-h^{11})\left[\dot{h}_{12}\mid_x\sum_{k=1}^{d}Q([Y_k^1,Y_k^2],X) + \dot{f}\mid_x\sum_{k=1}^{\frac{d}{2}}Q([Y_k^1,Y_{k+\frac{d}{2}}^2]-[Y_{k+\frac{d}{2}}^1,Y_k^2],X)\right].$$

As in the orthogonal case, we can assume that $h_{11}\mid_x \neq h_{22}\mid_x$. Equation $r(X,\frac{\partial}{\partial t})\mid_x = 0$ of the Einstein condition becomes

$$\dot{h}_{12}\mid_x\sum_{k=1}^{d}Q([Y_k^1,Y_k^2],X) + \dot{f}\mid_x\sum_{k=1}^{\frac{d}{2}}Q([Y_k^1,Y_{k+\frac{d}{2}}^2]-[Y_{k+\frac{d}{2}}^1,Y_k^2],X) = 0 \tag{3.13}$$

for all $X\in\mathfrak{p}$.

We can now prove the unitary analogue of theorem (3.1).

Theorem 3.2. *Let (M, g) be an Einstein manifold of cohomogeneity-one under the action of a compact connected group G such that the isotropy representation has two equivalent* ***unitary*** *summands of dimension d, with all other summands distinct. The Einstein metric can be diagonalized globally if*

$$\dim\left(\text{span}\left\{\sum_{k=1}^{d}[Y_k^1, Y_k^2]_{\mathfrak{p}}, \sum_{k=1}^{\frac{d}{2}}([Y_k^1, Y_{k+\frac{d}{2}}^2]_{\mathfrak{p}} - [Y_{k+\frac{d}{2}}^1, Y_k^2]_{\mathfrak{p}})\right\}\right) = 2$$

where the Y_j^i are as defined above.

Proof. If the span of the above vectors is of dimension two then equation (3.13) implies that $\dot{h}_{12}\mid_x = \dot{f}\mid_x = 0$.

Therefore, we have demonstrated that, in the Einstein case, $h_{12}\mid_x = f\mid_x = \dot{h}_{12}\mid_x = \dot{f}\mid_x$. This implies that $h_{12} \equiv 0$ and $f \equiv 0$. □

Example: $G/K \cong SU(n+1)/U(n-1)$

This is a well-known homogeneous manifold which was studied by Calabi and others. It can be viewed as a bundle over $\mathbb{C}P^n \cong SU(n+1)/U(n)$ with fiber $S^{2n-1} \cong U(n)/U(n-1)$. We represent $\mathfrak{su}(n+1)$ as the set of $(n+1)\times(n+1)$ complex matrices A such that $A^* + A = 0$ and $\text{tr}(A) = 0$. We embed $U(n-1)$ in $SU(n+1)$ so that $\mathfrak{u}(n-1)$ sits in $\mathfrak{su}(n+1)$ as

$$\mathfrak{k} \cong \mathfrak{u}(n-1) \cong \begin{pmatrix} \mathfrak{u}(n-1) & 0 \\ 0 & 0 \end{pmatrix} \subset \mathfrak{su}(n+1).$$

Because the notation is cumbersome, we will only calculate the $n = 2$ result and leave the rest to the reader. Schematically, we can write $\mathfrak{su}(3)$ as

$$\begin{pmatrix} ia & \alpha & \beta \\ -\bar{\alpha} & ib & \gamma \\ -\bar{\beta} & -\bar{\gamma} & i(-b-a) \end{pmatrix}$$

where $a, b \in \mathbb{R}$ and α, β, and $\gamma \in \mathbb{C}$. Let

$$\mathfrak{u}(1) = \text{span}\left\{\begin{pmatrix} ia & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}\right\}.$$

The isotropy representation is

$$\mathfrak{so}(n+1) = \mathfrak{u}(n-1) \oplus \mathfrak{p}_1 \oplus \mathfrak{p}_2 \oplus \mathfrak{p}_3 \oplus \mathfrak{p}_4 \oplus \mathfrak{p}_5$$

where $\mathfrak{p}_1 \simeq \mathfrak{p}_2$ are equivalent unitary representations of real dimension $2n-2$ and $\mathfrak{p}_3 \simeq \mathfrak{p}_4 \simeq \mathfrak{p}_5$ are trivial one-dimensional representations. Moreover,

$$\mathfrak{p}_3 \oplus \mathfrak{p}_4 \oplus \mathfrak{p}_5 \cong \mathfrak{su}(2).$$

In this case, there are two sets of equivalent summands. That is a general metric would be of the form

$$g_t(\cdot,\cdot)|_{\mathfrak{p}_1\oplus\mathfrak{p}_2} = Q((h+f),\cdot,\cdot)|_{\mathfrak{p}_1\oplus\mathfrak{p}_2}$$

with h and f as above and

$$g_t(\cdot,\cdot))|_{\mathfrak{p}_3\oplus\mathfrak{p}_4\oplus\mathfrak{p}_5} = Q(\bar{h}\cdot,\cdot)|_{\mathfrak{p}_3\oplus\mathfrak{p}_4\oplus\mathfrak{p}_5}$$

where

$$\bar{h} = \begin{pmatrix} \bar{h}_{11} & \bar{h}_{12} & \bar{h}_{13} \\ \bar{h}_{12} & \bar{h}_{22} & \bar{h}_{23} \\ \bar{h}_{13} & \bar{h}_{23} & \bar{h}_{33} \end{pmatrix}$$

For our purposes we wish to study the pair of equivalent unitary representations. Therefore, we partially diagonalize the cohomogeneity-one metric.

Assumption A:

$$\bar{h}_{12} = \bar{h}_{13} = \bar{h}_{23} = 0.$$

Under this assumption, there are only two non-diagonal functions: h_{12} and f. Given the above assumption, can we diagonalize a cohomogeneity-one Einstein metric with this principal orbit type?

In this case,

$$\mathfrak{p}_1 = \text{span}\left\{ Y_1^1 = \begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, Y_2^1 = \begin{pmatrix} 0 & i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \right\}$$

and

$$\mathfrak{p}_2 = \text{span}\left\{ Y_1^2 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix}, Y_2^2 = \begin{pmatrix} 0 & 0 & i \\ 0 & 0 & 0 \\ i & 0 & 0 \end{pmatrix} \right\}.$$

A simple calculation demonstrates that $\text{span}\{[Y_1^1, Y_1^2]_{\mathfrak{p}} + [Y_2^1, Y_2^2]_{\mathfrak{p}}, [Y_1^1, Y_2^2]_{\mathfrak{p}} - [Y_2^1, Y_1^2]_{\mathfrak{p}}\}$ is of dimension two. Therefore, by theorem (3.2), any cohomogeneity-one Einstein metric with principal orbits isomorphic to $G/K \cong SU(n+1)/U(n-1)$ such that Assumption A holds can be diagonalized globally.

Alternatively, we could make a different assumption.

Assumption B:

$$h_{12} = f = 0.$$

As before, a cohomogeneity-one metric satisfying Assumption B is only partially diagonalized as $\bar{h}$ may contain non-diagonal entries. In this case, however, the situation is slightly more complicated as there are now three equivalent summands $\mathfrak{p}_3, \mathfrak{p}_4$, and $\mathfrak{p}_5$ which are no longer automatically orthogonal. We will go over the general case of more than two equivalent summands below. However, we do not need those results to analyze this particular case. Note that

$$\mathfrak{p}_3 \oplus \mathfrak{p}_4 \oplus \mathfrak{p}_5 \cong \mathfrak{su}(2).$$

Therefore, given Assumption B, diagonalizing a cohomogeneity-one Einstein metric with principal orbits isomorphic to $G/K \cong SU(n+1)/U(n-1)$ is equivalent to diagonalizing a cohomogeneity-one Einstein metric with $SU(2)$ principal orbits. That, of course, is possible.

To summarize, a cohomogeneity-one Einstein metric with $G/K \cong SU(n+1)/U(n-1)$ can be diagonalized globally if Assumption A or Assumption B hold. If neither hold, we cannot diagonalize the metric in the Einstein case using this method.

3.4.3 The Symplectic Case

When $\mathfrak{p}_1$ and $\mathfrak{p}_2$ are symplectic representations, we have that $g_t(\cdot,\cdot) \mid_{\mathfrak{p}_1 \oplus \mathfrak{p}_2} = Q((h+f)\cdot,\cdot) \mid_{\mathfrak{p}_1 \oplus \mathfrak{p}_2}$ where

$$h = \begin{pmatrix} h_{11}\mathrm{Id}_d & \\ & h_{22}\mathrm{Id}_d \end{pmatrix}$$

and

$$f = \begin{pmatrix} & & f^1\mathrm{Id}_{\frac{d}{2}} & f^2\mathrm{Id}_{\frac{d}{2}} \\ & & f^3\mathrm{Id}_{\frac{d}{2}} & f^4\mathrm{Id}_{\frac{d}{2}} \\ f^1\mathrm{Id}_{\frac{d}{2}} & f^2\mathrm{Id}_{\frac{d}{2}} & & \\ f^3\mathrm{Id}_{\frac{d}{2}} & f^4\mathrm{Id}_{\frac{d}{2}} & & \end{pmatrix}$$

relative to basis $(Y_1^1, ..., Y_d^1, Y_1^2, ..., Y_d^2)$. Note that we have defined h differently than we had before. In the symplectic case, there are four non-diagonal functions. When can we conclude that all the $f^j = 0$ globally in the Einstein case? We can diagonalize the metric at a point $x \in M$. That is $f^j \mid_x = 0$ for $1 \leq j \leq 4$. At x we calculate that

$$r\left(X, \frac{\partial}{\partial t}\right) \mid_x = \frac{1}{2}h^{11}\frac{\partial}{\partial t}\left(\sum_{k=1}^{d}\langle Y_k^1, [Y_k^1, X]\rangle\right) \mid_x + \frac{1}{2}h^{22}\frac{\partial}{\partial t}\left(\sum_{k=1}^{d}\langle Y_k^2, [Y_k^2, X]\rangle\right) \mid_x$$

for all $X \in \mathfrak{p}$. Making use of the bi-invariance of Q we have

$$\langle Y_k^1, [Y_k^1, X]\rangle = g_t(Y_k^1, [Y_k^1, X]) = -f^1 Q([Y_k^1, Y_k^2], X) - f^3 Q([Y_k^1, Y_{k+\frac{d}{2}}^2], X)$$

for $1 \leq k \leq \frac{d}{2}$, and

$$\langle Y_k^1, [Y_k^1, X]\rangle = g_t(Y_k^1, [Y_k^1, X]) = -f^2 Q([Y_k^1, Y_{k-\frac{d}{2}}^2], X) - f^4 Q([Y_k^1, Y_k^2], X)$$

for $\frac{d}{2} + 1 \leq k \leq d$ for all $X \in \mathfrak{p}$.

Similar calculations hold for $\langle Y_k^2, [Y_k^2, X]\rangle$ and we have that

$$r\left(X, \frac{\partial}{\partial t}\right) \mid_x = \frac{1}{2}(h^{22} - h^{11})\{\dot{f}^1 \mid_x \sum_{k=1}^{\frac{d}{2}} Q([Y_k^1, Y_k^2], X) + \dot{f}^3 \mid_x \sum_{k=1}^{\frac{d}{2}} Q([Y_k^1, Y_{k+\frac{d}{2}}^2], X)$$

$$+\dot{f}^2 \mid_x \sum_{k=\frac{d}{2}+1}^{d} Q([Y_k^1, Y_{k-\frac{d}{2}}^2], X) + \dot{f}^4 \mid_x \sum_{k=\frac{d}{2}+1}^{d} Q([Y_k^1, Y_k^2], X)\}.$$

We can assume, as above, that $h^{11} \mid_x \neq h^{22} \mid_x$. With this assumption, equation $r(X, \frac{\partial}{\partial t}) \mid_x = 0$ of the Einstein condition becomes

$$\dot{f}^1 \mid_x Q\left(\sum_{k=1}^{\frac{d}{2}} [Y_k^1, Y_k^2], X\right) + \dot{f}^3 \mid_x Q\left(\sum_{k=1}^{\frac{d}{2}} [Y_k^1, Y_{k+\frac{d}{2}}^2], X\right) \tag{3.14}$$

$$+\dot{f}^2 \mid_x Q\left(\sum_{k=\frac{d}{2}+1}^{d} [Y_k^1, Y_{k-\frac{d}{2}}^2], X\right) + \dot{f}^4 \mid_x Q\left(\sum_{k=\frac{d}{2}+1}^{d} [Y_k^1, Y_k^2], X\right) = 0$$

for all $X \in \mathfrak{p}$.

We now have the symplectic analogue to theorem (3.1).

Theorem 3.3. *Let (M, g) be an Einstein manifold of cohomogeneity-one under the action of a compact, connected group G such that the isotropy representation has two equivalent symplectic summands of dimension d with all others distinct. The Einstein metric can be diagonalized globally if*

$$\dim\left(\text{span}\left\{\sum_{k=1}^{\frac{d}{2}} [Y_k^1, Y_k^2]_{\mathfrak{p}}, \sum_{k=1}^{\frac{d}{2}} [Y_k^1, Y_{k+\frac{d}{2}}^2]_{\mathfrak{p}}, \sum_{k=\frac{d}{2}+1}^{d} [Y_k^1, Y_{k-\frac{d}{2}}^2]_{\mathfrak{p}}, \sum_{k=\frac{d}{2}+1}^{d} [Y_k^1, Y_k^2]_{\mathfrak{p}}\right\}\right) = 4$$

where the Y_i^j are as defined above.

Proof. If the span of the above vectors is of dimension four then equation (3.14) implies that $\dot{f}^i \mid_x = 0$ for all $1 \leq i \leq 4$. Since all the $f^i \mid_x = 0$, we deduce that $f^i \equiv 0$ for all $1 \leq i \leq 4$. □

3.5 The Case of r Equivalent Summands

Having analyzed the orthogonal, unitary, and symplectic case for isotropy representations with only pairs of equivalent summands, we now consider the case in which

$$\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}_1 \oplus ... \oplus \mathfrak{p}_r \oplus \mathfrak{p}_{r+1} \oplus ... \oplus \mathfrak{p}_m$$

where $\mathfrak{p}_i \simeq \mathfrak{p}_j$ for $1 \leq i, j \leq r$ and all other $\mathfrak{p}_l$ for $l > r$ are distinct.

As above, we fix a bi-invariant metric Q on $\mathfrak{g}$ where $\{Y_j^i\}_{j=1,...,d_i}$ forms a Q-orthonormal basis for each $\mathfrak{p}_i$. The fiber metrics g_t can be written

$$g_t(\cdot,\cdot) = g_t(\cdot,\cdot) \mid_{\mathfrak{p}_1\oplus\ldots\oplus\mathfrak{p}_r} \overset{\perp}{\bigoplus_{i=r+1,\ldots,m}} A_i(t)Q(\cdot,\cdot) \mid_{\mathfrak{p}_i}$$

Since the restriction of g_t to $\mathfrak{p}_1 \oplus ... \oplus \mathfrak{p}_r$ is $Ad(K)$-invariant, it can be written as $g_t(\cdot,\cdot) \mid_{\mathfrak{p}_1\oplus\ldots\oplus\mathfrak{p}_r} = Q(h\cdot,\cdot) \mid_{\mathfrak{p}_1\oplus\ldots\oplus\mathfrak{p}_r}$ where h is a positive definite $Ad(K)$-invariant map $h : \mathfrak{p}_1 \oplus ... \oplus \mathfrak{p}_r \to \mathfrak{p}_1 \oplus ... \oplus \mathfrak{p}_r$. The form of h depends upon whether the representation $\mathfrak{p}_1$ is orthogonal, unitary, or symplectic. We again consider each of these case individually.

3.5.1 The Orthogonal Case

When $\mathfrak{p}_1, ..., \mathfrak{p}_r$ are orthogonal representations, we have that h is given by the matrix

$$h = (h_{ij} Id_d)$$

where $1 \leq i, j \leq r$, $h_{ij} = h_{ij}(t)$ and $d = d_1 = ... = d_r$. This implies that a generic cohomogeneity-one manifold of this type has $\frac{r(r-1)}{2}$ independent non-diagonal functions. We want to determine when all of these functions can be set equal to zero at every point of an Einstein manifold without loss of generality.

We calculate that

$$r\left(X, \frac{\partial}{\partial t}\right) = \frac{1}{2} h^{ij} \frac{\partial}{\partial t}\left(\sum_{k=1}^{d} \langle Y_k^i, [Y_k^j, X]\rangle\right)$$

for all $X \in \mathfrak{p}$. We can, without loss of generality, diagonalize the metric at a point $x \in M$ which implies that $h_{ij} \mid_x = 0$ for $i \neq j$. Given this assumption, we have

$$r\left(X, \frac{\partial}{\partial t}\right) \mid_x = \frac{1}{2} \sum_{i=1}^{r} h^{ii} \frac{\partial}{\partial t}\left(\sum_{k=1}^{d} \langle Y_k^i, [Y_k^i, X]\rangle\right) \mid_x .$$

From the bi-invariance of Q, we calculate that

$$\langle Y_k^i, [Y_k^i, X]\rangle = g_t(Y_k^i, [Y_k^i, X]) = \sum_{j=1}^{r} h_{ij} Q(Y_j^j, [Y_k^i, X])$$

$$= -\sum_{j=1}^{r} h_{ij} Q([Y_k^i, Y_k^j], X).$$

Therefore, at x, the Einstein condition implies that

$$\sum_{1\leq i<j\leq r} \left\{ (h^{jj} - h^{ii}) Q\left(\sum_{k=1}^{d} [Y_k^i, Y_k^j], X\right)\right\} \dot{h}_{ij} \mid_x = 0 \tag{3.15}$$

for all $X \in \mathfrak{p}$.

To simplify our presentation we define vectors Z_{ij} by the equation

$$Z_{ij} = \sum_{k=1}^{d} [Y_k^i, Y_k^j]_{\mathfrak{p}}$$

for $1 \leq i < j \leq r$.

We are now ready to prove the diagonalization theorem for this type of group action.

Theorem 3.4. *Let (M, g) be an Einstein manifold of cohomogeneity-one under the action of a compact, connected group G such that the isotropy representation has at most r equivalent **orthogonal** summands of dimension d with all others distinct. The Einstein metric can be diagonalized globally if*

$$\dim(\text{span}\{Z_{ij} \mid 1 \leq i < j \leq r\}) = \frac{r(r-1)}{2}. \tag{3.16}$$

Proof. As above, we can assume that $h_{ii} \mid_x \neq h_{jj} \mid_x$ for $i \neq j$. Therefore, if the span of the Z_{ij}'s has dimension $\frac{r(r-1)}{2}$, then equations (3.15) become $\frac{r(r-1)}{2}$ independent equations in the $\dot{h}_{ij} \mid_x$ terms. Since there are $\frac{r(r-1)}{2}$ such terms, we deduce that $\dot{h}_{ij} \mid_x = 0$ for $i \neq j$. Since we already have that $h_{ij} \mid_x = 0$ for $i \neq j$, we have that $h_{ij} \equiv 0$ for all $i \neq j$. The metric is then globally diagonal. □

3.5.2 The Unitary Case

When $\mathfrak{p}_1, ..., \mathfrak{p}_r$ are equivalent unitary representations of dimension d, we have that $g_t(\cdot, \cdot) \mid_{\mathfrak{p}_1 \oplus ... \oplus \mathfrak{p}_r} = Q((h+f)\cdot, \cdot) \mid_{\mathfrak{p}_1 \oplus ... \oplus \mathfrak{p}_r}$ relative to basis $(Y_1^1, ..., Y_d^r)$ where h is of the form

$$h = (h_{ij} Id_d)$$

for $1 \leq i, j \leq r$ and f is defined by

$$f_{ij} = g_t(Y_k^i, Y_{k+\frac{d}{2}}^j)$$

for $i < j$ and $1 \leq k \leq \frac{d}{2}$ and

$$f_{ij} = -g_t(Y_k^i, Y_{k-\frac{d}{2}}^j)$$

for $i < j$ and $\frac{d}{2} + 1 \leq k \leq d$.

After diagonalizing the metric at point $x \in M$, that is $h_{ij} \mid_x = f_{ij} \mid_x = 0$ for $i \neq j$, equation $r(X, \frac{\partial}{\partial x}) \mid_x = 0$ of the Einstein condition becomes

$$\sum_{1 \leq i < j \leq r} (h^{jj} - h^{ii}) \left\{ \dot{h}_{ij} \mid_x \sum_{k=1}^{d} Q([Y_k^i, Y_k^j], X) + \dot{f}_{ij} \mid_x \sum_{k=1}^{\frac{d}{2}} Q([Y_k^i, Y_{k+\frac{d}{2}}^j] - [Y_{k+\frac{d}{2}}^i, Y_k^2], X) \right\} = 0 \tag{3.17}$$

for all $X \in \mathfrak{p}$.

To simplify our presentation, we define the vectors

$$Z_{ij} = \sum_{k=1}^{d} [Y_k^i, Y_k^j]_{\mathfrak{p}}$$

and

$$W_{ij} = \sum_{k=1}^{\frac{d}{2}} ([Y_k^i, Y_{k+\frac{d}{2}}^j]_{\mathfrak{p}} - [Y_{k+\frac{d}{2}}^i, Y_k^j]_{\mathfrak{p}})$$

for $1 \leq i < j \leq r$.

Given the above formalism, we can now prove the following theorem.

Theorem 3.5. *Let (M, g) be an Einstein manifold of cohomogeneity-one under the action of a compact, connected group G such that the isotropy representation has at most r equivalent **unitary** summands of dimension d with all others distinct. The Einstein metric can be diagonalized globally if*

$$\dim(\text{span}\{Z_{ij}, W_{ij} \mid 1 \leq i < j \leq r\}) = r(r-1).$$

Proof. The proof of this theorem is similar to the proof of the analogous theorem in the orthogonal case. The only difference is that, in the unitary case, there are $r(r-1)$ independent non-diagonal functions. The rest follows in an identical fashion. □

3.5.3 The Symplectic Case

When $\mathfrak{p}_1, ..., \mathfrak{p}_r$ are equivalent symplectic representations of dimension d, we have that $g_t(\cdot,\cdot) \mid_{\mathfrak{p}_1 \oplus ... \oplus \mathfrak{p}_r} = Q((h+f)\cdot,\cdot) \mid_{\mathfrak{p}_1 \oplus ... \oplus \mathfrak{p}_r}$ relative to basis $(Y_1^1, ..., Y_d^r)$ where h is of the form

$$h = \begin{pmatrix} h_{11} & & & \\ & h_{22} & & \\ & & ... & \\ & & & h_{rr} \end{pmatrix}$$

and f is defined, for $1 \leq i, j \leq r$ by

$$f_{ij}^1 = g_t(Y_k^i, Y_k^j)$$

for $1 \leq k \leq \frac{d}{2}$,

$$f_{ij}^3 = g_t(Y_k^i, Y_{k+\frac{d}{2}}^j)$$

for $1 \leq k \leq \frac{d}{2}$,

$$f_{ij}^2 = g_t(Y_k^i, Y_{k-\frac{d}{2}}^j)$$

for $\frac{d}{2} + 1 \leq k \leq d$, and

$$f_{ij}^4 = g_t(Y_k^i, Y_k^j)$$

for $\frac{d}{2} + 1 \leq k \leq d$.

After diagonalizing the metric at point $x \in M$, that is $f_{ij}^l \mid_x = 0$ for all l, i, and j, equation $r(X, \frac{\partial}{\partial t}) \mid_x = 0$ of the Einstein condition becomes

$$\sum_{1 \leq i < j \leq r} (h^{jj} - h^{ii}) \{\dot{f}_{ij}^1 \mid_x Q\left(\sum_{k=1}^{\frac{d}{2}} [Y_k^i, Y_k^j], X\right) + \dot{f}_{ij}^3 \mid_x Q\left(\sum_{k=1}^{\frac{d}{2}} [Y_k^i, Y_{k+\frac{d}{2}}^j], X\right) \tag{3.18}$$

$$+\dot{f}^2_{ij} \mid_x Q\left(\sum_{k=\frac{d}{2}+1}^{d}[Y^i_k, Y^j_{k-\frac{d}{2}}], X\right) + \dot{f}^4_{ij} \mid_x Q\left(\sum_{k=\frac{d}{2}+1}^{d}[Y^i_k, Y^j_k], X\right)\} = 0$$

for all $X \in \mathfrak{p}$. To simplify our presentation, we define vectors

$$Z^1_{ij} = \sum_{k=1}^{\frac{d}{2}}[Y^i_k, Y^j_k]_{\mathfrak{p}}, \quad Z^3_{ij} = \sum_{k=1}^{\frac{d}{2}}[Y^i_k, Y^j_{k+\frac{d}{2}}]_{\mathfrak{p}}, \quad Z^2_{ij} = \sum_{k=\frac{d}{2}+1}^{d}[Y^i_k, Y^j_{k-\frac{d}{2}}]_{\mathfrak{p}},$$

and

$$Z^4_{ij} = \sum_{k=\frac{d}{2}+1}^{d}[Y^i_k, Y^j_k]_{\mathfrak{p}}$$

for $1 \leq i < j \leq r$. Given the above formalism, we can now prove the following theorem.

Theorem 3.6. *Let (M, g) be an Einstein manifold of cohomogeneity-one under the action of compact, connected group G such that the isotropy representation has at most r equivalent **symplectic** summands of dimension d with all others distinct. The Einstein metric can be diagonalized globally if*

$$\dim(\text{span}\{Z^l_{ij} \mid 1 \leq l \leq 4, 1 \leq i < j \leq r\}) = 2r(r-1).$$

Proof. The proof of this theorem is identical to the proof of the analogous theorems in the orthogonal and unitary case, except that in the symplectic case there are $2r(r-1)$ non-diagonal equations which need to be eliminated. □

3.6 The Case of a Trivial Isotropy Subgroup

Let M be an n-dimensional manifold of cohomogeneity-one under the action of group G such that each principal orbit is a copy of G, i.e. the isotropy subgroup K is trivial. In this case, the Lie algebra $\mathfrak{g}$ decomposes into $n-1$ irreducible summands of dimension one

$$\mathfrak{g} = \mathfrak{p}_1 \oplus ... \oplus \mathfrak{p}_{n-1}. \tag{3.19}$$

We choose a basis $\{Y^i\}_{i=1,...n-1}$ as described above. (Note that since each summand is of dimension one, we have omitted the subscript k.)

We have the following corollary to theorem (3.4) .

Corollary 3.7. *Let (M^n, g) be an Einstein manifold of cohomogeneity-one under the action of a compact group G such that the isotropy subgroup is trivial, i.e. $K = \{1\}$. Equation $r(X, \frac{\partial}{\partial t}) = 0$ of the Einstein condition can be used to globally diagonalize the metric g if and only if $\dim M = 4$ and $G \cong SU(2)$.*

Proof. As in the previous section, define $Z_{ij} = [Y^i, Y^j] \mid_{\mathfrak{p}} = [Y^i, Y^j]$ for $i < j$. Because each of the equivalent summands is one dimensional, each is an orthogonal representation and we can apply

theorem (3.4). In this case, $r = n-1$. Therefore, equation $r(X, \frac{\partial}{\partial t}) = 0$ of the Einstein can be used to diagonalize the metric globally if and only if $\dim(\text{span}\{Z_{ij}\}) = \frac{(n-1)(n-2)}{2}$. But the span of the set of Z_{ij}'s must be less than or equal to the dimension of each principal orbit, $n-1$. The inequality $\frac{(n-1)(n-2)}{2} \leq (n-1)$ implies that $n \leq 4$. Clearly, the dimension of the cohomogeneity-one manifold is greater than one, otherwise the principal orbits would be trivial.

We have shown that the dimension of the manifold is two, three, or four. If the manifold has dimension two, then the manifold is diagonal trivially and $r(X, \frac{\partial}{\partial t})$ automatically. If the manifold has dimension three, then the group G has dimension two. The only compact Lie Group of dimension two is abelian. Therefore, equation $r(X, \frac{\partial}{\partial t})$ of the Einstein condition is satisfied automatically. We cannot then use that equation to diagonalize the metric. If the manifold has dimension four, then G is of dimension three. The only compact Lie groups of dimension three (up to finite quotient) are T^3 and $SU(2)$. Since T^3 is abelian, $r(X, \frac{\partial}{\partial t}) = 0$ automatically and we cannot use this equation to diagonalize the metric. Finally, we have already seen how we can diagonalize an Einstein metric with $SU(2)$ principal orbits. □

Chapter 4

Compact Einstein Four Manifolds with Torus Isometry

Having discussed a problem concerning Einstein metrics of cohomogeneity-one in the previous chapter, we now consider the Einstein condition on a manifold of cohomogeneity-two. In particular, we specialize to the case of a compact Einstein four manifold admitting a T^2 isometry. After discussing what is known about Einstein four manifolds with large symmetry group, we review the topological classification of smooth torus actions on compact, simply-connected four manifolds due to Orlik and Raymond [49]. Also, we will review the nuts and bolts formalism developed by Gibbons and Hawking for retrieving topological data from the vanishing of Killing vectors to the case of a compact four manifold with torus symmetry. Although the action of a T^2 isometry on a compact four manifold need not be automatically orthogonally transitive, a result in [39] tells us that the O'Neill tensor A must vanish if the metric is Einstein. Using this result we are able to substantially simplify the equations of the Einstein condition by making use of a natural complex structure on the quotient space M/T^2. After this simplification, we see that the Einstein condition becomes four second-order *elliptic* equations of four functions in two variables. After reviewing in detail all of the known examples of compact Einstein four manifolds admitting a T^2-isometry we look at the special case in which the Einstein metric is globally diagonal. In this special case, we derive some further consequences of the Einstein condition.

4.1 Compact Einstein Four Manifolds with Symmetry

The Einstein condition on manifolds of dimension two and three is equivalent to the constant curvature condition; therefore, the Einstein condition on manifolds of dimension two and three reduces to a topological problem. Any manifold of dimension two admits an Einstein metric since every two dimensional manifold admits a metric of constant curvature [6]. A three dimensional Einstein manifold has a universal covering diffeomorphic to $\mathbb{R}^3$ or S^3. At present there exists no classification of Einstein four manifolds even in the compact case, but some success has been achieved in classifying

Einstein four manifolds with large symmetry groups. Homogeneous four manifolds are effectively classified by the following theorem.

Theorem 4.1. *[32] Every homogeneous Einstein four manifold is symmetric.*

As all symmetric spaces were classified by Cartan, it is known that the only compact homogeneous Einstein four manifolds are T^4, S^4, $S^2 \times S^2$, and $\mathbb{C}P^2$ as well as certain finite quotients of these manifolds. The work of Cartan together with the above theorem gives the following result.

Theorem 4.2. *If (M, g) is a compact homogeneous Einstein four manifold, then, up to homothety, M must be T^4, S^4, $\mathbb{R}P^4$, $\mathbb{C}P^2$, $S^2 \times S^2$, $S^2 \times \mathbb{R}P^2$, $(S^2 \times S^2)/\{\pm(1,1)\}$, or $\mathbb{R}P^2 \times \mathbb{R}P^2$ endowed with their standard metrics.*

Since the Einstein condition on homogeneous four manifolds is understood, it is logical to next consider the Einstein condition on cohomogeneity-one four manifolds. Cohomogeneity-one manifolds were studied by Berard-Bergery who was able to classify all compact Einstein cohomogeneity-one four manifolds, barring one exception.

Theorem 4.3. *[44] If (M, g) is a compact cohomogeneity-one Einstein four manifold which is not homogeneous and does not have principal orbits isometric to $SU(2)$ then, up to homothety, M is $(S^2 \times S^2)/\{\pm(\delta_1, -1)\}$, $(S^2 \times S^2)/\{\pm(\delta_2, -1)\}$, $(S^2 \times \mathbb{R}P^2)/\{\pm(\delta_2, -1)\}$, $\mathbb{C}P^2 \sharp \overline{\mathbb{C}P^2}$, or $\mathbb{C}P^2 \sharp \mathbb{R}P^2$, where $\delta_1 = \mathrm{diag}(-1, 1, 1)$ and $\delta_2 = \mathrm{diag}(-1, -1, 1)$. The first three manifolds are endowed with their standard metrics and the last two are endowed with the D. Page metric or its $\mathbb{Z}_2$ quotient*

The Einstein condition on manifolds of cohomogeneity-one with $SU(2)$ principal orbits gives rise to what are known as the *Bianchi IX* equations. These have been studied extensively [13] but there exists no complete classification of their solutions.

Aside from the Bianchi IX case, therefore, compact cohomogeneity-one Einstein four manifolds have been completely classified. In order to complete the classification of compact Einstein manifolds with isometry group of dimension greater than or equal to one, it remains to consider the cohomogeneity-two and cohomogeneity-three cases. In the cohomogeneity-two case, since each principal orbit must be a homogeneous two manifold, the only possible principal orbit types are S^2, $\mathbb{R}P^2$, and T^2. The following theorem by Derdzinski eliminates the first two orbit possibilities.

Theorem 4.4. *[17] If (M, g) is a compact cohomogeneity-two Einstein four manifolds with principal orbits isometric to S^2 or $\mathbb{R}P^2$ then M must be homogeneous or cohomogeneity-one.*

Therefore, we need only consider the case of T^2 principal orbits.

Finally, a compact cohomogeneity-three manifold must have S^1 principal orbits. Below, we will study the Einstein condition on compact cohomogeneity-two four manifolds with principal orbits isometric to T^2. We will also look briefly at cohomogeneity-three four manifolds with S^1 principal orbits by imposing the additional condition of orthogonal transitivity.

4.2 Compact Simply-Connected Four Manifolds Admitting a Smooth T^2 Action

As in Chapter 2, any compact Einstein four manifold with two commuting Killing vectors must be T^4 or a manifold with positive Einstein constant and finite fundamental group. For the sake of simplicity, we will, for the moment, restrict our attention to the simply-connected case. This restriction does not greatly reduce the complexity of the problem but will allow for a cleaner presentation. Due to the work of Orlik and Raymond, there exists a topological classification of smooth four manifolds admitting a smooth torus action [49]. In this section, we will give a brief review of their results following the summary found in [35].

Let M be a smooth compact simply-connected four manifold admitting a smooth, effective torus action and set $G = T^2 = SO(2) \times SO(2)$.

Proposition 4.5. *[49] The action of G on M must have fixed points.*

At fixed points, the stabilizer is the whole of G. In addition to fixed points, M admits one-dimensional subspaces upon which the stability group is a subgroup isomorphic to $SO(2) = S^1$. There are many such subgroups, each of which can be described by a pair of integers.

The group G can be given coordinates (φ, ψ) where $0 \leq \varphi < 2\pi$ and $0 \leq \psi < 2\pi$. Every subgroup of G isomorphic to $SO(2)$ can be written as $G(m, n)$ where $m\varphi + n\psi = 0$ and m and n are coprime integers. Any two such subgroups $G(m, n)$ and $G(m', n')$ will generate the homology of G if and only if $mn' - m'n = \pm 1$ [49]. We will study not only the quotient space M/G but also the quotient of M by the subgroups $G(m, n)$. Therefore, we state two propositions which will be used below.

Proposition 4.6. *[49] If M is oriented, then $M/G(m, n)$ is a three space with boundary for all coprime integer pairs m and n.*

Proposition 4.7. *[49] If M is compact and oriented, then M/G is a compact two manifold with piecewise smooth boundary.*

Note that in the simply-connected case, M/G will be a compact two manifold with a piecewise smooth boundary polygon. In [49], it is demonstrated that $\partial M/G$ is a one-dimensional manifold containing a finite number of isolated fixed points. Therefore, M/G can be viewed as a polygon whose interior points have G orbits in M, whose edges away from the vertices have S^1 orbits, and whose vertices have point orbits in M. Furthermore, the stabilizer of the orbits along the interior of a given edge must be constant, while there is a 'jump' in the stabilizer group at the endpoints [49]. Therefore each edge of the polygon can be described by its stabilizer group $G(m, n)$. At the intersection of two edges, the stabilizers $G(m, n)$ and $G(m', n')$ of the S^1 orbits of the two edges must generate the homology of G. Therefore, at each vertex $mn' - m'n = 1$.

The above results demonstrate that any smooth, effective T^2 action on a compact, simply-connected four manifold can be described by a finite number of coprime integer pairs, $(m_i, n_i)_{1\leq i\leq k}$ where $(m_i, n_i) \sim (-m_i, -n_i)$ and $m_i n_{i+1} - m_{i+1} n_i = \pm 1$ for $1 \leq i < k$ and $m_k n_1 - m_1 n_1 = \pm 1$. The following theorem states the conditions under which two sets of integer pairs define the same action on the same manifold.

Theorem 4.8. *[49] There exists an equivariant diffeomorphism between $M = \{(m_1, n_1), ..., (m_k, n_k)\}$ and $M' = \{(m'_1, n'_1), ..., (m'_{k'}, n'_{k'})\}$ if and only if $k = k'$ and, for some fixed integer t, $(m_i, n_i) = (m'_{t+i}, n'_{t+i})$ or $(m_i, n_i) = (-m'_{t+i}, -n'_{t+i})$ for $i = 1, ..., k$.*

Given such a set of integer pairs, one can uniquely identify the manifold corresponding to the action described. Indeed, any such manifold admitting a smooth T^2 action must be S^4 or connected sums of $\mathbb{C}P^2$, $\overline{\mathbb{C}P^2}$, and $S^2 \times S^2$ [49]. This identification of integer pair sets with the manifolds listed above is not a bijection since, in general, any such manifold will admit many inequivalent torus actions. It is well known that $(S^2 \times S^2)\sharp\mathbb{C}P^2$ is equivalent to $\mathbb{C}P^2\sharp\overline{\mathbb{C}P^2}\sharp\mathbb{C}P^2$. Therefore, aside from S^4 and $\sharp^n(S^2 \times S^2)$, all compact four manifolds admitting a torus action are of the form $k\mathbb{C}P^2\sharp l\overline{\mathbb{C}P^2}$.

Although we will not here reproduce the entirety of the classification, we note along with Joyce in [34] that certain topological information can be readily obtained from a collection of integer pairs defining a torus action. Given an action defined by $\{(m_1, n_1), ..., (m_k, n_k)\}$, because the integer pairs are defined only up to sign, we can choose the integer pairs so that $m_j \geq 0$ and $n_j > 0$ whenever $m_j = 0$. The Euler characteristic of the manifold, M, admitting that action is given by

$$\chi(M) = k,$$

where k is the number of sides of the polygon forming the boundary of the quotient manifold M/G. The signature of the manifold defined by the same torus action is

$$\tau(M) = b_+ - b_- = m_1 n_k - m_k n_1 + \sum_{j=1}^{k-1} m_{j+1} n_j - m_j n_{j+1}.$$

Recall that $\chi(S^4) = 2$ and $\tau(S^4) = 0$ and that $\chi(\sharp^n(S^2 \times S^2)) = 2n + 2$ and $\tau(\sharp^n(S^2 \times S^2)) = 0$. Finally, we recall that, for $M = k\mathbb{C}P^2\sharp l\overline{\mathbb{C}P^2}$, $\chi(M) = 2 + k + l$ and $\tau(M) = k - l$ [41]. The Euler characteristic and the signature do not uniquely determine the torus action or even the manifold itself. For example, $S^2 \times S^2$ and $\mathbb{C}P^2\sharp\overline{\mathbb{C}P^2}$ both have Euler characteristic four and signature zero. The difference is that $S^2 \times S^2$ is spin while $\mathbb{C}P^2\sharp\overline{\mathbb{C}P^2}$ is not.

Theorem 4.9. *[19], [21] A smooth simply connected oriented four manifold is determined up to homeomorphism by its Euler characteristic, χ, its signature, τ, and whether it is spin or non-spin.*

Since a given torus action can be associated to at most one manifold, one must be able to distinguish between torus actions on spin manifolds and those on non-spin manifolds. By analyzing the classification of Orlik and Raymond, it can be deduced that a manifold is spin if and only if $m_{j+2}n_j - m_j n_{j+2}$ as well as $m_2 n_k - m_k n_2$ and $m_1 n_{k-1} - m_{k-1} n_1$ are even.

These topological invariants can be used to tell us which manifolds admit which types of torus action. The Euler characteristic of a manifold M is equal to the number of sides of the polygon forming the boundary of M/G. Therefore, it immediately follows that two polygons with a different number of sides cannot possibly bound the orbit space of the same manifold. Below, when we endeavor to endow these manifolds with a T^2-invariant metric, the signature of the manifold will provide us with detailed information relating to the vanishing of Killing vectors.

4.3 The Einstein Equations

The classification of Orlik and Raymond was used to show that a simply-connected compact four manifold admitting a metric with a torus isometry must be S^4, $\sharp^l(S^2 \times S^2)$, or $k\mathbb{C}P^2 \sharp l\overline{\mathbb{C}P^2}$. However, the Hitchin-Thorpe inequality informs us that many of these manifolds cannot admit Einstein metrics of any kind.

Theorem 4.10. *[30] If M is a smooth compact oriented manifold admitting an Einstein metric, the*

$$2\chi(M) \geq 3|\tau(M)| \tag{4.1}$$

with equality if and only if the manifold is locally hyper-Kahler.

Both S^4 and $\sharp^l(S^2 \times S^2)$ have zero signature, so they satisfy the Hitchin-Thorpe inequality. However, many of the other manifolds are excluded. The Euler characteristic of a manifold of type $k\mathbb{C}P^2 \sharp l\overline{\mathbb{C}P^2}$ is given by $\chi = 2 + k + l$ and the signature is given by $\tau = k - l$. Since no such manifold can be locally hyper-Kahler, we have the following

Corollary 4.11. *A manifold $k\mathbb{C}P^2 \sharp l\overline{\mathbb{C}P^2}$ cannot admit an Einstein metric unless $4 + 5k > l > \frac{k-4}{5}$.*

When determining which of the remaining manifolds admit an Einstein metric with a T^2 isometry, there is no reason, *a priori*, that the action of the torus must be orthogonally transitive. However, because of the commuting Killing vectors must vanish somewhere on a compact manifold M, we have the following proposition.

Proposition 4.12. *[39] For M a compact Einstein four manifold with two commuting Killing vectors, the invariant A of the Riemannian submersion $\pi : M \to M/G$ vanishes and, equivalently, the action of the G is orthogonally transitive.*

Proof. Take X^a and Y^a to be the two commuting Killing vectors. Since X and Y commute, $\mathcal{L}_Y X = 0$ which implies that $\mathcal{L}_Y(X \wedge dX) = 0$. If we define $\phi = X \wedge dX$ and $\psi = Y \wedge dY$ ($\phi_{abc} = X_{[a}\nabla_b X_{c]}$ and $\psi_{abc} = Y_{[a}\nabla_b Y_{c]}$) then $\mathcal{L}_Y \phi = 0$ and $\mathcal{L}_X \psi = 0$. The well-known *twist scalars* α and β are defined to be $\alpha = *(Y \wedge \phi) = Y_{[a}\phi_{bcd]}\epsilon^{abcd}$ and $\beta = *(X \wedge \psi) = X_{[a}\psi_{bcd]}\epsilon^{abcd}$.

In order to determine α, we first calculate $\nabla^a(Y_{[a}\phi_{bcd]})$.

$$\nabla^a(Y_{[a}\phi_{bcd]}) = \frac{1}{4}\nabla^a(Y_a\phi_{bcd} - Y_b\phi_{cda} + Y_c\phi_{dab} - Y_d\phi_{abc}).$$

Knowing that $\nabla^a Y_a$ and $\mathcal{L}_Y\phi = 0$, a straightforward calculation shows that

$$\nabla^a(Y_{[a}\phi_{bcd]}) = \frac{1}{4}(-Y_b\nabla^a\phi_{cda} + Y_c\nabla^a\phi_{dab} - Y_d\nabla^a\phi_{abc}).$$

Now,

$$\nabla^a\phi_{abc} = \frac{1}{3}\nabla^a(X_a\nabla_b X_c + X_b\nabla_c X_a + X_c\nabla_a X_b).$$

Using the antisymmetry of ∇X and the fact that $\nabla_b\nabla_c X_d = R_{abcd}X^a$, we have

$$\nabla^a\phi_{abc} = \frac{1}{3}(X^a R_{dabc}X^d + X_b R_d{}^a{}_{ca}X^d + X_c R_d{}^a{}_{ab}X^d).$$

Using the Bianchi identities, we see that the first term vanishes and the above becomes

$$\nabla^a\phi_{abc} = \frac{1}{3}(-X_b R_{cd}X^d + X_c R_{bd}X^d).$$

Since M is an Einstein manifold $R_{ab} = \lambda g_{ab}$ and we see that $\nabla^a\phi_{abc} = 0$ and therefore $\nabla^a(Y_{[a}\phi_{bcd]}) = 0$. This implies that $\nabla_a\alpha = 0$ and, equivalently, that α is constant.

By the same argument, β is seen to be a constant. Indeed, since X and Y must vanish, we find that $\alpha = *(Y \wedge \phi) = \beta = *(X \wedge \psi) = 0$. The vanishing of the twist scalars is a necessary and sufficient condition for the action of the torus to be orthogonally transitive by [45].

□

Let X_1 and X_2 be two independent commuting Killing vectors. As above, we set $h = (h_{ij}) = (g_{ab}X_i^a X_b^j) = (X_i^a X_{ja})$ where the entries h_{ij} are functions of the coordinates of M/G. The previous theorem implies that on the open union of principal T^2 orbits, an Einstein metric can be written as $g = g_{M/G} + h = \breve{g} + h$, where $\breve{g}$ is a smooth metric on the two dimensional interior of M/G. Below, we let ∇ denote the Levi-Civita connection of g and let $\breve{\nabla}$ denote the Levi-Civita connection of $\breve{g}$. Now, because the metric h is positive definite on M/G, $\det h \geq 0$. More precisely, we know that $\det(h) > 0$ on the interior of M/G and $\det(h) = 0$ on $\partial M/G$. More precisely, $\det(h) = 0$ on the fixed point set of the actions of all the S^1 subgroups of T^2.

Proposition 4.13. *For (M, g) a compact four manifold with two commuting Killing vectors, the Einstein condition is equivalent to the following system of partial differential equations:*

$$-\frac{1}{2}\check{\triangle}h_{ij} - \frac{1}{4}\frac{\langle\check{\nabla}\det(h), \check{\nabla}h_{ij}\rangle}{\det(h)} - \frac{1}{2}h_{ik}\langle\check{\nabla}h^{kl}, \check{\nabla}h_{lj}\rangle = \lambda h_{ij} \tag{4.2}$$

$$-\frac{1}{2}h^{ij}\check{\nabla}_a\check{\nabla}_b h_{ij} - \frac{1}{4}\check{\nabla}_a h^{ij}\check{\nabla}_b h_{ij} = -\check{r}_{ab} + \lambda\check{g}_{ab} = (\lambda - \frac{\check{s}}{2})\check{g}_{ab}. \tag{4.3}$$

Proof. This is immediate from the calculations above giving the Einstein equations for an orthogonally transitive action. Because each principal orbit is a torus, $\hat{r} = 0$. For the last line we have used the fact that for any two-dimensional metric $\check{g}$, the Ricci tensor can be written as $\check{r} = (\frac{\check{s}}{2})\check{g}$. □

The Einstein condition on this class of manifold is a system of partial differential equations in two variables. This system must satisfy a host of boundary conditions on the points of $\partial(M/G)$. Indeed a great deal of information about the structure of M/G can be obtained by applying these boundary conditions to the equations of the Einstein condition. As it stands, the Einstein condition on a compact four manifold with T^2-symmetry is a system of six partial differential equation, three in (4.2) and three in (4.3), in two variables. We can show that, so long as *all* of the equations are satisfied on the boundary, then only *four* of these equations must be satisfied on the interior of M/G for the Einstein condition to hold.

First, we write equations (4.2) and (4.3) in a slightly different form by decomposing them into their traces and their trace-free parts. Let

$$\rho = \sqrt{\det(h)}.$$

We see that (4.2) can be written more compactly as

$$-\frac{1}{2}h_{ik}\rho^{-1}\check{\nabla}_a(\rho h^{kl}\check{\nabla}^a h_{lj}) = \lambda h_{ij}.$$

Multiplying both sides of this equation by h^{ki} we have

$$\check{\nabla}_a(\rho h^{kl}\check{\nabla}^a h_{jl}) = -2\lambda\rho\delta^k_j.$$

This can be written more schematically as

$$\check{\nabla}_a(\rho h^{-1}\check{\nabla}^a h) = -2\lambda\rho\mathrm{Id}_2 \tag{4.4}$$

where Id_2 is the 2×2 identity matrix. To separate the trace and the trace-free parts of this equation define matrix K by

$$h = \rho K.$$

This implies that $h^{-1} = \rho^{-1}K^{-1}$ and $\det K = 1$. Substituting this into equation (4.4) implies that

$$\check{\nabla}_a(K^{-1}\check{\nabla}^a(\rho K)) = -2\lambda\rho\mathrm{Id}_2$$

which expands to

$$(\check{\triangle}\rho)\mathrm{Id}_2 + \check{\nabla}_a(\rho K^{-1}\check{\nabla}^a K) = -2\lambda\rho\mathrm{Id}_2$$

Taking the trace of this equation gives

$$2\check{\triangle}\rho + \mathrm{tr}[\check{\nabla}_a(\rho K^{-1}\check{\nabla}^a K)] = -4\lambda\rho$$

which is equivalent to

$$2\check{\triangle}\rho + \check{\nabla}_a(\rho\ \mathrm{tr}[K^{-1}\check{\nabla}^a K]) = -4\lambda\rho.$$

Recall that $\mathrm{tr}[K^{-1}\check{\nabla}^a K] = (\mathrm{det}K)^{-1}\check{\nabla}^a \mathrm{det}K = 0$ since $\mathrm{det}K = 1$. Therefore, the trace of equation (4.2) is equivalent to

$$\check{\triangle}\rho = -2\lambda\rho. \tag{4.5}$$

It is clear from the above calculation that the trace-free part of equation (4.2) becomes

$$\check{\nabla}_a(\rho K^{-1}\check{\nabla}^a K) = 0. \tag{4.6}$$

We next perform similar calculations for equation (4.3). Writing this equation schematically gives

$$\frac{1}{2}\mathrm{tr}[h^{-1}\check{\nabla}_a\check{\nabla}_b h + \frac{1}{2}\check{\nabla}_a h^{-1}\check{\nabla}_b h] = (\frac{\check{s}}{2} - \lambda)\check{g}_{ab}$$

which is equivalent to

$$\frac{1}{2}\check{\nabla}_a\mathrm{tr}[h^{-1}\check{\nabla}_b h] - \frac{1}{4}\mathrm{tr}[\check{\nabla}_a h^{-1}\check{\nabla}_b h] = (\frac{\check{s}}{2} - \lambda)\check{g}_{ab}.$$

Recalling that $\mathrm{tr}[h^{-1}\check{\nabla}_b h] = \rho^{-2}\check{\nabla}_b(\rho^2)$, this becomes

$$\rho^{-1}\check{\nabla}_a\check{\nabla}_b\rho - \rho^{-2}\check{\nabla}_a\rho\check{\nabla}_b\rho + \frac{1}{4}\mathrm{tr}[h^{-1}\check{\nabla}_a h h^{-1}\check{\nabla}_b h] = (\frac{\check{s}}{2} - \lambda)\check{g}_{ab}.$$

Once again we set $h = \rho K$ which implies that

$$\frac{1}{4}\mathrm{tr}[h^{-1}\check{\nabla}_a h h^{-1}\check{\nabla}_b h] = \frac{1}{4}\mathrm{tr}[\rho^{-2}\check{\nabla}_a\rho\check{\nabla}_b\rho\mathrm{Id}_2] + \frac{1}{4}\mathrm{tr}[K^{-1}\check{\nabla}_b K]\rho^{-1}\check{\nabla}_a\rho + \frac{1}{4}\mathrm{tr}[K^{-1}\check{\nabla}_a K K^{-1}\check{\nabla}_b K].$$

The second term vanishes, and equation (4.3) becomes

$$\rho^{-1}\check{\nabla}_a\check{\nabla}_b\rho - \frac{1}{2}\rho^{-2}\check{\nabla}_a\rho\check{\nabla}_b\rho + \frac{1}{4}\mathrm{tr}[K^{-1}\check{\nabla}_a K K^{-1}\check{\nabla}_b K] = (\frac{\check{s}}{2} - \lambda)\check{g}_{ab}. \tag{4.7}$$

Taking the trace of this equation implies that

$$\rho^{-1}\check{\triangle}\rho - \frac{1}{2}\rho^{-2}\langle\check{\nabla}\rho, \check{\nabla}\rho\rangle + \frac{1}{4}\mathrm{tr}[K^{-1}\check{\nabla}_a K K^{-1}\check{\nabla}^a K] = \check{s} - 2\lambda. \tag{4.8}$$

Combining this with equation (4.5) gives

$$\check{s} = -\frac{1}{2}\rho^{-2}\langle\check{\nabla}\rho, \check{\nabla}\rho\rangle + \frac{1}{4}\mathrm{tr}[K^{-1}\check{\nabla}_a K K^{-1}\check{\nabla}^a K]. \tag{4.9}$$

For the moment, we will not write out the trace-free part of equation (4.3).

So far, we have considered $\check{g}$ in a coordinate invariant way. For our purposes, we wish to express $\check{g}$ in terms of *isothermal coordinates*. As the quotient space is two-dimensional, the induced metric $\check{g}$, like all other metrics on a two-dimensional manifold [6], is conformally flat. Therefore, we can set

$$\check{g} = \Omega^2(dx^2 + dy^2)$$

where $\Omega = \Omega(x, y)$. These are called *isothermal coordinates*. A complex structure can be placed, locally, on M/G by setting

$$z = x + iy.$$

We define the complex differential in the standard way,

$$\partial = \frac{1}{2}(\frac{\partial}{\partial x} - i\frac{\partial}{\partial y}).$$

We wish to calculate the Einstein equations (4.2) and (4.3), or more specifically their traces and their trace-free parts, in terms of the function Ω and the operators ∂ and $\bar{\partial}$. These equations, expressed in complex coordinates, are collected in the following corollary to the above proposition.

Corollary 4.14. *On a compact four-dimensional manifold (M, g) admitting a T^2-isometry, the Einstein condition is equivalent to the following system of equations*

$$\partial\bar{\partial}\rho = -\frac{1}{2}\lambda\rho\Omega^2 \tag{4.10}$$

$$\partial(\rho K^{-1}\bar{\partial}K) + \bar{\partial}(\rho K^{-1}\partial K) = 0 \tag{4.11}$$

$$\partial\bar{\partial}\log\Omega = \frac{1}{4}\frac{\partial\rho\bar{\partial}\rho}{\rho^2} - \frac{1}{8}\mathrm{tr}[K^{-1}\partial K K^{-1}\bar{\partial}K] \tag{4.12}$$

$$8\frac{\partial\log\Omega\partial\rho}{\rho} - 4\frac{\partial\partial\rho}{\rho} + 2\frac{\partial\rho\partial\rho}{\rho^2} - \mathrm{tr}[K^{-1}\partial K K^{-1}\partial K] = 0 \tag{4.13}$$

where ρ, Ω, and the matrix K are as defined above.

Proof. The proof is a straightforward calculation; simply evaluate equations (4.2) and (4.3) for the coordinates given. We point out that (4.10) is the equivalent to the trace of (4.2) and (4.11) is equivalent to the trace-free part of (4.2). Furthermore, equation (4.12) is obtained by subtracting (4.10) from the trace of equation (4.3). Finally, (4.13) is equivalent to the trace-free part of (4.3).

We note that in terms of the isothermal coordinates and the given complex structure, $\check{s} = -8\partial\bar{\partial}\log\Omega$. □

For convenience we define Q by

$$Q = 8\frac{\partial\log\Omega\partial\rho}{\rho} - 4\frac{\partial\partial\rho}{\rho} + 2\frac{\partial\rho\partial\rho}{\rho^2} - \mathrm{tr}[K^{-1}\partial K K^{-1}\partial K].$$

Equation (4.13) is then equivalent to $Q = 0$. We have the following proposition.

Proposition 4.15. *If (4.10), (4.11), and (4.12) hold (equivalently if (4.2) and the trace of (4.3) hold) then the complex quantity ρQ is holomorphic. That is,*

$$\bar{\partial}(\rho Q) = 0.$$

Proof. From above,

$$\rho Q = 8\partial\log\Omega\partial\rho - 4\partial\partial\rho + 2\frac{\partial\rho\partial\rho}{\rho} - \rho\mathrm{tr}[K^{-1}\partial K K^{-1}\partial K].$$

Taking the $\bar{\partial}$ derivative, and using the fact that $\partial\bar{\partial} = \bar{\partial}\partial$, gives

$$\bar{\partial}(\rho Q) = 8\partial\bar{\partial}\log\Omega\partial\rho + 8\partial\log\Omega\partial\bar{\partial}\rho - 4\partial\partial\bar{\partial}\rho + 4\frac{\partial\bar{\partial}\rho\partial\rho}{\rho} - 2\frac{\bar{\partial}\rho\partial\rho\partial\rho}{\rho^2}$$
$$-\bar{\partial}\rho\mathrm{tr}[K^{-1}\partial K K^{-1}\partial K] - \rho\bar{\partial}\{\mathrm{tr}[K^{-1}\partial K K^{-1}\partial K]\}.$$

By equation (4.10), we find that

$$-4\partial\partial\bar{\partial}\rho = 2\lambda\partial(\rho\Omega^2) = 2\lambda\Omega^2\partial\rho + 4\lambda\rho\Omega^2\partial\log\Omega,$$

$$8\partial\log\Omega\partial\bar{\partial}\rho = -4\lambda\rho\Omega^2\partial\log\Omega,$$

and

$$4\frac{\partial\bar{\partial}\rho\partial\rho}{\rho} = -2\lambda\Omega^2\partial\rho.$$

Equation (4.12) implies that

$$8\partial\bar{\partial}\log\Omega\partial\rho = 2\frac{\partial\rho\partial\rho\bar{\partial}\rho}{\rho^2} - \partial\rho\mathrm{tr}[K^{-1}\partial K K^{-1}\bar{\partial}K].$$

Together these results imply that

$$\bar{\partial}(\rho Q) = -\rho\bar{\partial}\{\mathrm{tr}[K^{-1}\partial K K^{-1}\partial K]\} - (\partial\rho)\mathrm{tr}[K^{-1}\partial K K^{-1}\bar{\partial}K] - (\bar{\partial}\rho)\mathrm{tr}[K^{-1}\partial K K^{-1}\partial K].$$

We calculate that

$$\bar{\partial}\{\mathrm{tr}[K^{-1}\partial K K^{-1}\partial K]\} = 2\mathrm{tr}[\bar{\partial}(K^{-1}\partial K)K^{-1}\partial K]$$
$$= \mathrm{tr}\{[\bar{\partial}(K^{-1}\partial K) + \partial(K^{-1}\bar{\partial}K)]K^{-1}\partial K\}$$
$$= \frac{1}{\rho}\mathrm{tr}\{[\bar{\partial}(\rho K^{-1}\partial K) + \partial(\rho K^{-1}\bar{\partial}K)]K^{-1}\partial K\} - \frac{\partial\rho}{\rho}\mathrm{tr}[K^{-1}\partial K K^{-1}\bar{\partial}K] - \frac{\bar{\partial}\rho}{\rho}\mathrm{tr}[K^{-1}\partial K K^{-1}\partial K].$$

To justify the second equality we note that

$$\mathrm{tr}[\bar{\partial}(K^{-1}\partial K)K^{-1}\partial K] = -\mathrm{tr}[K^{-1}\bar{\partial}K K^{-1}\partial K K^{-1}\partial K] + \mathrm{tr}[K^{-1}\bar{\partial}\partial K K^{-1}\partial K].$$

Since the operators ∂ and $\bar{\partial}$ commute and $\mathrm{tr}(ABC) = \mathrm{tr}(BCA) = \mathrm{tr}(CAB)$ we see that

$$\mathrm{tr}[\bar{\partial}(K^{-1}\partial K)K^{-1}\partial K] = \mathrm{tr}[\partial(K^{-1}\bar{\partial}K)K^{-1}\partial K].$$

This implies that

$$\bar{\partial}(\rho Q) = -\mathrm{tr}\{[\bar{\partial}(\rho K^{-1}\partial K) + \partial(\rho K^{-1}\bar{\partial}K)]K^{-1}\partial K\}.$$

Equation (4.11) implies that the right-hand side vanishes. This complete the proof. □

Since the manifold M is assumed to be compact, the quotient space M/G is automatically compact. By the results of Orlik and Raymond discussed above, the quotient space M/G is simply-connected. Because ρQ is holomorphic, by Cauchy's Theorem,

$$\rho Q|_{\partial(M/G)} = 0 \Rightarrow \rho Q|_{M/G} \equiv 0.$$

Since $\rho > 0$ on the interior of M/G (this must be true for the metric to be nondegenerate there), we see that

$$\rho Q|_{M/G} \equiv 0 \Rightarrow Q|_{\mathrm{int}(M/G)} = 0$$

where $\mathrm{int}(M/G)$ denotes the interior of the quotient space. It is clear that if Q is identically zero on the interior of a compact set, then it must be zero at the boundary as well; that is,

$$Q|_{\mathrm{int}(M/G)} \Rightarrow Q|_{M/G} \equiv 0.$$

This is precisely equation (4.13) of the Einstein condition.

Theorem 4.16. *If equations (4.10)-(4.12) of the Einstein condition hold on a compact four manifold with T^2-symmetry then equation (4.13) need only hold at the boundary of M/T^2 for the metric to be Einstein everywhere.*

In essence, we have seen that the Einstein condition on a compact four dimensional manifold with T^2-symmetry reduces from a system of six partial differential equations to a system of *four* partial differential equations in four variables along with additional boundary conditions.

4.4 Nuts and Bolts

In [26], Gibbons and Hawking study Einstein four manifolds admitting at least one Killing vector and reveal how topological information of the manifold can be recovered from examining the fixed point set a Killing vector. On a four manifold, a given Killing vector can vanish either on a two-surface, called a bolt, or at a point, called a nut. In the case of a four manifold admitting a torus symmetry, there is a two dimensional family of Killing vectors and any given Killing vector will vanish on a collection of nuts and bolts. After reviewing the theory developed in [26], we will apply that theory to the special case of a compact Einstein four manifold with two commuting Killing vectors.

Given an oriented four manifold M admitting a one-parameter isometry group, we denote by X the Killing vector generating that isometry group, $\mu : M \to M$. At a fixed point p of the isometric action, the Killing vector X vanishes and the map $\mu_* : T_pM \to T_pM$ becomes an isometry. As noted above, the matrix $\nabla X = X_{a;b}$ is an antisymmetric 4×4 matrix and at a fixed point, any such matrix must have rank 0, 2, or 4. Were the rank of the matrix ∇X to equal zero, then the rank would be zero everywhere and the action of the isometry would be trivial. Therefore, we assume that the rank is either 2 or 4 at a fixed point.

If ∇X has rank two, then the Killing vector X vanishes on a totally geodesic two dimensional submanifold whose tangent space in TM is invariant under the action of the map μ_* and the submanifold of fixed points is referred to as a *bolt* . The two-dimensional distribution orthogonal to the tangent space of this submanifold will be rotated by this action and the period of that rotation will equal $2\pi\kappa^{-1}$ where κ is a rational number called the *surface gravity of the bolt*. We note along with Gibbons and Hawking that κ is the nonzero skew eigenvalue of ∇X relative to an orthonormal frame.

On the other hand, if ∇X has rank 4 at a fixed point p of μ, then p must be isolated as no subspace of T_pM is acted on trivially by the map μ_*. Any such fixed point is called a *nut* (the name refers to the fixed point found in the Taub-NUT metric). At a nut, the matrix ∇X will have two skew-eigenvalues κ_1 and κ_2. The map μ_* will rotate with periods $2\pi\kappa_1^{-1}$ and $2\pi\kappa_2^{-1}$ two orthogonal subspaces of T_p. Nuts can be divided into sets of *nuts* and *antinuts*. The point p is referred to as a nut when $\kappa_1\kappa_2$ is positive and as an antinut when $\kappa_1\kappa_2$ is negative.

Using fixed point theorems, Gibbons and Hawking were able to express the Euler characteristic and the signature of the manifold M in terms of the number of nuts and antinuts and the Euler characteristics of the bolts of a given Killing vector. We will here present the formulae for the Euler characteristic and the signature without proof and refer the reader to [26]. Given a manifold M with Killing vector X vanishing on N_+ nuts, N_- antinuts, and n bolts $(B_i)_{i=1,\ldots,n}$ with Euler characteristics χ_i, the Euler characteristic of M is given by

$$\chi(M) = N_+ + N_- + \sum_{i=1}^{n} \chi_i$$

and the signature of M is given by

$$\tau = N_+ - N_-.$$

We now want to apply this nuts and bolts formalism to the case of a simply connected compact Einstein four manifold with two commuting Killing vectors. In the following discussion, a 'side' of the polygon boundary of M/G will be defined to include its endpoints. Given a side A of the polygon, there is a Killing vector, say X_1, which vanishes on the two sphere $\pi^{-1}(A)$. For simplicity, we will say that X_1 has a bolt at A when, strictly speaking, it actually has a bolt at $\pi^{-1}(A)$. Any other Killing vector must have a nut (or an antinut) at each of the two vertices contained in that side. Any two nuts located at the boundary points of a side of the polygon can be seen to be contained in the bolt of the Killing vector vanishing on that side. For a fixed Killing vector X, a vertex will be viewed as either an isolated nut or as part of a bolt, i.e. a given vertex will be viewed as a nut of vector field X if and only if it is not part of a bolt of X. Consequently, any Killing vector of the torus will vanish on a certain collection of nuts, antinuts, and bolts. From each such collection, one can

retrieve the topological information of the manifold and, of course, any two collections associated to two different Killing vectors on the same manifold must encode the same topological data.

Recalling the classification of Orlik and Raymond, any smooth torus action on a compact simply connected smooth four manifold can be described as a set of coprime integer pairs $(m_i, n_i)_{i=1,\ldots,k}$. In the case of an isometric torus action, each integer pair defines which Killing vector of the torus is vanishing on the side of the polygon associated to that pair. For example, if side A is described by $(1, 0)$ then it is Killing vector X_1 that vanishes on side A. Because the set of coprime integer pairs describing a given torus action contains all of the topological data of the manifold, the nuts and bolts data must be compatible with the set of integer pairs. In order to understand this compatibility, we must be able to express the surface gravities of the bolts in terms of the functions F_i and their derivatives.

Without loss of generality, we take X_1 to be the Killing vector vanishing on side A of $\partial M/G$. Because X_1 has a bolt at A, the matrix ∇X_1 has rank two. If we take ∂_1 and ∂_2 to be an orthonormal basis for M/G and take ∂_3 and ∂_4 to be an orthonormal basis for the vertical distribution, then ∇X_1 will, on all of M looks schematically like

$$\nabla X_1 = \begin{pmatrix} 0 & 0 & -a & -b \\ 0 & 0 & -c & -d \\ a & c & 0 & 0 \\ b & d & 0 & 0 \end{pmatrix}.$$

The upper left and lower right hand two by two submatrices are equal to zero because of orthogonal transitivity and the commuting of the Killing vectors respectively.

Gibbons and Hawking state that the 'surface gravity' of the bolt at A is given by the skew eigenvalue of ∇X_1 relative to an orthonormal basis. Put another way, the surface gravity of the bolt at A, which we will label as κ_A, satisfies the equation

$$\nabla_a X_{1b} \nabla^a X_1^b|_A = \kappa_A^2. \tag{4.14}$$

We see that $\nabla_a X_{1b} \nabla^a X_1^b|_A$ must be a constant because for Killing vector X_1

$$\nabla_b \nabla_c X_{1d} = R_{abcd} X_1^a$$

which vanishes when X_1 does. In order to express the surface gravity of the bolt at A in terms of the h_{ij} and their derivatives we note that

$$\nabla_b X_{1a} \nabla^b X_1^a|_A = \nabla^b (\frac{1}{2} \nabla_b h_{11})|_A.$$

This implies that

$$\frac{1}{2} \triangle h_{11}|_A = \kappa_A^2.$$

On a general side of type (m, n), we have the following:

Proposition 4.17. *The bolt corresponding to a side with stabilizer group defined by the coprime integer pair (m, n) has* ***surface gravity*** *κ satisfying the equation*

$$m^2 \triangle h_{11} + 2mn \triangle h_{12} + n^2 \triangle h_{22} = 2\kappa^2. \tag{4.15}$$

4.5 Examples

At first blush, demanding a torus symmetry appears extremely restrictive. However, all known Einstein metrics on compact four manifolds admitting a symmetry group of dimension greater than or equal to one admit a torus isometry. Aside from the possible exception of the Bianchi IX case, all Einstein metrics on compact four manifolds with symmetry group of dimension at least two *must* admit a torus isometry. Explicit Einstein metrics of manifolds of this type are few; in the compact simply connected case only four such metrics are known (on S^4, $\mathbb{C}P^2$, $S^2 \times S^2$, and $\mathbb{C}P^2 \sharp \overline{\mathbb{C}P^2}$) and all of them are homogeneous or cohomogeneity one. Complex geometry has been used to prove the existence of a T^2 invariant Kahler-Einstein metric on $\mathbb{C}P^2 \sharp 3\overline{\mathbb{C}P^2}$ but the metric is not known explicitly. Below, we review in detail those metrics that are known and examine the torus symmetry of each.

4.5.1 (S^4, can)

The four sphere admits only one smooth torus action up to equivariant diffeomorphism [49]. That action can be described by the set of coprime integer pairs $\{((1,0),(0,1)\}$. Indeed, this is the only action whose quotient space is bounded by a polygon with two sides. The canonical metric on the four sphere can be written as $d\sigma_4^2 = dR^2 + \sin^2 R d\sigma_3^2$ where $d\sigma_3^2$ is the standard metric on S^3. If we use the coordinates of the Hopf fibration then the metric becomes

$$d\sigma_4^2 = dR^2 + \sin^2 R d\theta^2 + \sin^2 R \sin^2\theta d\varphi^2 + \sin^2 R \cos^2\theta d\psi^2.$$

∂_φ and ∂_ψ are the two commuting Killing vectors of the T^2 action. Relative to these vectors the matrices h become

$$h = \begin{pmatrix} \sin^2 R \sin^2\theta & 0 \\ 0 & \sin^2 R \cos^2\theta \end{pmatrix}.$$

It is easy to check that these equations are satisfied by the above metric on S^4 and that $\lambda = 3$ and $\check{s} = 2$.

What is the structure of M/G? The metric induced on the quotient space is $dR^2 + \sin^2 R d\theta^2$ which is the standard metric on the two sphere. The vertices are located at $R = 0$ and $R = \pi$ as the rank of J becomes zero at those points and the sides are located at $\theta = 0$ and $\theta = \frac{\pi}{2}$. Note that ∂_φ has a bolt at $\theta = 0$ and ∂_ψ has a bolt at $\theta = \frac{\pi}{2}$. Any other Killing vector will have a nut and an antinut at $R = 0$ and $R = \pi$ or vice versa.

4.5.2 ($\mathbb{C}P^2$, Fubini-Study Metric)

Like the four sphere, two dimensional complex projective space admits only one torus action up to equivariant diffeomorphism [49]. That torus action can be described by the set $\{(0,1),(1,0),(1,1)\}$. Note that transposing two of the coprime integer pairs would give the torus action on $\overline{\mathbb{C}P^2}$. The Fubini-Study metric can be written as

$$ds^2 = dR^2 + \sin^2 R d\theta^2 + \sin^2 R \sin^2\theta(1 - \sin^2 R \sin^2\theta)d\varphi^2 + \sin^2 R \cos^2\theta(1 - \sin^2 R \cos^2\theta)d\psi^2$$

$$-2\sin^4 R \sin^2\theta cos^2\theta d\varphi d\psi$$

where ∂_φ and ∂_ψ are the two commuting Killing vectors. The matrices h and F then become

$$h = \begin{pmatrix} \sin^2 R \sin^2\theta(1 - \sin^2 \sin^2\theta) & -\sin^4 R \sin^2\theta \cos^2\theta \\ -\sin^4 R \sin^2\theta \cos^2\theta & \sin^2 R \cos^2\theta(1 - \sin^2 R \cos^2\theta) \end{pmatrix}.$$

For this metric, the scalar curvature of the base manifold is $\check{s} = 2$ and the Einstein constant is $\lambda = 6$. The quotient space of the this manifold is a triangle on the standard two-sphere whose induced metric is the canonical sphere metric. The sides of the triangle are located at $\theta = 0$, $\theta = \frac{\pi}{2}$, and $R = \frac{\pi}{2}$.

Killing vector ∂_ϕ has a bolt at $\theta = 0$, Killing vector ∂_ψ has a bolt at $\theta = \frac{\pi}{2}$, and Killing vector $\partial_\phi + \partial_\psi$ has a bolt at $R = \frac{\pi}{2}$.

4.5.3 ($S^2 \times S^2$, can)

The standard Einstein metric on $S^2 \times S^2$ can be written as

$$ds^2 = dR^2 + d\theta^2 + \sin^2 R d\varphi^2 + \sin^2\theta d\psi^2$$

where ∂_φ and ∂_ψ are the two commuting Killing vectors. This metric is invariant under the torus action defined by the set $\{(0,1),(1,0),(0,1),(1,0)\}$. Relative to this action, the matrices h become

$$h = \begin{pmatrix} \sin^2 R & 0 \\ 0 & \sin^2\theta \end{pmatrix}.$$

As in the case of the standard metric on S^4, $h_{12} \equiv 0$. For this metric, $\check{s} = 0$ and $\lambda = 1$.

The quotient space for this torus action is a rectangle whose induced metric is the flat Euclidean metric on $\mathbb{R}^2$. The sides of the rectangle are located at $R = 0$, $R = \pi$, $\theta = 0$, $\theta = \pi$.

Killing vector ∂_φ has bolts at $R = 0$ and $R = \pi$ and vector ∂_ψ has bolts at $\theta = 0$ and $\theta = \pi$.

4.5.4 ($\mathbb{C}P^2\sharp\overline{\mathbb{C}P^2}$, D. Page Metric)

The Page metric on $\mathbb{C}P^2\sharp\overline{\mathbb{C}P^2}$ can be written as

$$ds^2 = \frac{3}{\lambda}(1+\nu^2)\{\frac{1-\nu^2\cos^2 R}{3-\nu^2-\nu^2(1+\nu^2)\cos^2 R}dR^2 + \frac{(1-\nu^2\cos^2 R)}{3+6\nu^2-\nu^4}d\theta^2$$

$$+\frac{1-\nu^2\cos^2 R}{3+6\nu^2-\nu^4}\sin^2\theta d\phi^2 + \frac{3-\nu^2-\nu^2(1+\nu^2)\cos^2 R}{(3+\nu^2)^2(1-\nu^2\cos^2 R)}\sin^2 R(d\psi - \sin^2\frac{1}{2}\theta d\phi)^2\}$$

where $\nu \approx 0.2817$ is a solution to the equation $\frac{4\nu(3+\nu^2)}{3+6\nu^2-\nu^4} = 1$ and ∂_ϕ and ∂_ψ are the two commuting Killing vectors. This metric is invariant under the torus action defined by the set $\{(0,1),(1,1),(0,1),(1,0)\}$. The scalar curvature, $\check{s}$, of the quotient space, M/G, is a function of R only and is not constant. It is straightforward to calculate that

$$\check{s} = \frac{2}{3}\lambda\nu^2\frac{((\nu^4+\nu^6)\cos^6 R + (-3\nu^2-3\nu^4)\cos^4 R + (6-3\nu^2+3\nu^4)\cos^2 R - 3 + \nu^2)}{((\nu^6+\nu^8)\cos^6 R + (-3\nu^4-3\nu^6)\cos^4 R + (3\nu^2+3\nu^4)\cos^2 R - 1 - \nu^2)}.$$

The quotient space of this torus action is a four-sided polygon with metric of nonconstant scalar curvature. The sides of the polygon are located at $R = 0$, $R = \pi$, $\theta = 0$, and $\theta = \pi$.

Killing vector ∂_ψ has bolts at $R = 0$ and $R = \pi$. These geodesics are also lines of constant negative curvature of the base. We note that the scalar curvature on these two geodesics is in fact equal. Killing vector ∂_ϕ has a bolt at $\theta = 0$ and Killing vector $\partial_\phi + \partial_\psi$ has a bolt at $\theta = \pi$. The curvature on these two geodesics is the same, though it is not constant. Rather, it symmetric about the $R = \frac{\pi}{2}$ axis and is periodic. It is negative at the endpoints $R = 0$ and $R = \pi$ and positive around $R = \frac{\pi}{2}$.

4.6 The Diagonal Case

Above, we discussed the Einstein condition on compact four manifolds with T^2 symmetry, i.e. with two commuting Killing vectors. Using a result in [39] we demonstrated that the Einstein condition requires that the action be orthogonally transitive. This allowed us to write the metric as $g = \check{g} + h$ as discussed above. Using isothermal coordinates, the two-dimensional metric $\check{g}$ is always diagonalizable. However, it is not always possible to diagonalize the matrix h. One could seek to classify all Einstein metrics on compact four manifolds with two commuting Killing vectors such that $h_{12} \equiv 0$ on all of M. As a first step towards this goal, we present the Einstein equations in the diagonal case and give a couple of preliminary results. Below, we again specialize to the simply-connected case.

From the topological classification of Orlik and Raymond discussed above, any torus action can be described by a set of coprime integer pairs $\{(m_i, n_i)\}_{i=1,\ldots,k}$. For h_{12} to vanish on all of M, each (m_i, n_i) must equal $(1,0)$ or $(0,1)$. Furthermore, since the same Killing vector can not vanish on adjacent sides, the number of coprime integer pairs k must be even and the pairs must alternate

between $(1,0)$ and $(0,1)$. Such a manifold would have Euler characteristic $\chi = k$ and signature $\tau = 0$. Therefore, when $k = 2$, $M = S^4$ and for $k > 2$ and $l = k-2$, $M = \sharp^l(S^2 \times S^2)$. The standard metrics on S^4 and $S^2 \times S^2$ described in detail above are each diagonalizable. No Einstein metrics are known to exist on connected sums of two or more copies of $S^2 \times S^2$.

The Einstein condition on such a manifold takes a particularly simple form because of the vanishing of h_{12}. To better distinguish this case from the case of a general torus action we set $h_{11} = f_1^2$ and $h_{22} = f_2^2$. This substitution is possible because the functions h_{11} and h_{22} must be positive on the interior of M/G for the metric g to be positive definite.

Proposition 4.18. *The metric on an Einstein four manifold admitting a T^2-isometry such that the metric is globally diagonalizable can be written as $g = g_{M/G} + f_1^2 d\phi^2 + f_2^2 d\psi^2$. The Einstein condition becomes the following system of differential equations*

$$\frac{\check{\triangle} f_1}{f_1} + \frac{\langle \check{\nabla} f_1, \check{\nabla} f_2 \rangle}{f_1 f_2} = -\lambda \tag{4.16}$$

$$\frac{\check{\triangle} f_2}{f_2} + \frac{\langle \check{\nabla} f_1, \check{\nabla} f_2 \rangle}{f_1 f_2} = -\lambda \tag{4.17}$$

$$\frac{\check{\triangle} f_1}{f_1} + \frac{\check{\triangle} f_2}{f_2} = \check{s} - 2\lambda \tag{4.18}$$

along with the additional boundary conditions.

4.7 $\sharp^l(S^2 \times S^2)$

The quotient space of the torus action on S^4 reviewed above is a two-sided polygon on the two-sphere endowed with the canonical metric. This quotient metric has constant positive curvature. Similarly, the quotient space of the torus action on $S^2 \times S^2$ is a four-sided polygon on Euclidean space endowed with its canonical metric. This quotient metric has zero curvature.

Any T^2-invariant Einstein metric on $\sharp^l(S^2 \times S^2)$ which is globally diagonalizable must have a quotient space bounded by a polygon with $2l + 2$ sides. It is reasonable to ask whether the quotient space of such an Einstein metric might, for $l \geq 2$, have constant negative curvature. Unfortunately this supposition is too optimistic.

Proposition 4.19. *The quotient space of a T^2-invariant Einstein metric on $\sharp^l(S^2 \times S^2)$ for $l \geq 2$ which is globally diagonalizable cannot have constant curvature.*

Proof. The quotient space cannot have constant nonnegative curvature because the boundary polygon has more than four sides. Setting $V = f_1 f_2$ we note that V is nonzero on the interior of the

quotient space M/G and zero on the boundary $\partial M/G$. Therefore, V must have a critical point on the interior of M/G. At that point

$$\nabla V = f_1 \nabla f_2 + f_2 \nabla f_1 = 0$$

which implies that

$$f_1^2 |\nabla f_2|^2 + 2 f_1 f_2 \langle \nabla f_1, \nabla f_2 \rangle + f_2^2 |\nabla f_1|^2 = 0.$$

Equations (4.16) and (4.17) imply that $\langle \nabla f_1, \nabla f_2 \rangle = -\frac{\check{s}}{2} f_1 f_2$. Together with the above we have

$$f_1^2 |\nabla f_2|^2 - \check{s} f_1^2 f_2^2 + f_2^2 |\nabla f_1|^2 = 0.$$

On the interior of M/G, the functions f_1 and f_2 must be nonzero in order to ensure smoothness at interior points. Therefore, $\check{s}$ must be nonnegative. This implies that $\check{s}$ is nonconstant. □

Though our prediction concerning the quotient space structure of an Einstein metric failed, we are able to say something about the bolts of such an Einstein metric should it exist.

Let A_i denote the $(l+1)$-sides of type $(1,0)$ and let B_i denote the $(l+1)$-sides of type $(0,1)$. Let $\mathrm{Area}(A_i)$ and $\mathrm{Area}(B_i)$ denote the areas of the bolts corresponding to side A_i and B_i. If we set the period of both the S^1-actions to be 2π, then the surface gravity becomes

$$\kappa_{A_i}^2 = \kappa_{B_i}^2 = 1$$

for all $1 \leq i \leq l+1$ [26].

If we let η_{A_i} and η_{B_i} be the *outward* pointing unit normal vector fields to A_i and B_i respectively. The surface gravity equation implies that

$$\eta_{A_i} f_1 = \eta_{B_i} f_2 = -1.$$

Furthermore, along a side B_i, the Einstein condition requires that $\check{\nabla} f_1$ be tangent to all B_i and the $\check{\nabla} f_2$ be tangent to all A_i. We deduce immediately that

$$\eta_{A_i} f_2 = \eta_{B_i} f_1 = 0.$$

Consider equation (4.16) of the Einstein condition. This equation can be written as

$$f_2 \check{\triangle} f_1 + \langle \check{\nabla} f_1, \check{\nabla} f_2 \rangle = -\lambda f_1 f_2.$$

Taking the integral of both sides gives

$$\int_{M/G} f_2 \check{\triangle} f_1 + \langle \check{\nabla} f_1, \check{\nabla} f_2 \rangle d\mathrm{vol}_{\check{g}} = -\lambda \int_{M/G} f_1 f_2 d\mathrm{vol}_{\check{g}}. \tag{4.19}$$

The right-hand side of this equation becomes

$$-\lambda \int_{M/G} f_1 f_2 d\mathrm{vol}_{\check{g}} = -\frac{\lambda}{4\pi^2} \mathrm{Vol}(M)$$

where $\mathrm{Vol}(M)$ is the total volume of the manifold.

To analyze the left-hand side of (4.19), we make use of Green's first identity. Let u and v be smooth functions on Ω a compact region with piecewise smooth boundary. Let $(\partial\Omega)_i$ be the ith codimension-one element of the boundary and let η_i be the outward pointing unit normal vector field to the face $(\partial\Omega)_i$. Green' first identity states that

$$\int_\Omega v\triangle u + \langle\nabla u, \nabla v\rangle d\mathrm{vol}_\Omega = \sum_i \int_{(\partial\Omega)_i} v(\eta_i u) d\omega_i$$

where $d\omega_i$ is the area form of the face $(\partial\Omega_i)$.

Using this identity, the left-hand side of equation (4.19) becomes

$$\int_{M/G} f_2 \check{\triangle} f_1 + \langle\check{\nabla} f_1, \check{\nabla} f_2\rangle d\mathrm{vol}_{\check{g}} = \sum_{i=1}^{l+1} \int_{A_i} f_1(\eta_{A_i} f_2) dA_i + \sum_{i=1}^{l+1} \int_{B_i} f_1(\eta_{B_i} f_2) dB_i$$

where dA_i and dB_i are the elements of lengths of sides A_i and B_i. Recalling that $f_1|_{A_i} = 0$ and $\eta_{B_i} f_2 = -1$. We see

$$\int_{M/G} f_2 \check{\triangle} f_1 + \langle\check{\nabla} f_1, \check{\nabla} f_2\rangle d\mathrm{vol}_{\check{g}} = -\sum_{i=1}^{l+1} \int_{B_i} f_1 dB_i = -\frac{1}{2\pi} \sum_{i=1}^{l+1} \mathrm{Area}(B_i).$$

Performing a similar calculation for equation (4.17), we see that

$$2\pi \sum_{i=1}^{l+1} \mathrm{Area}(A_i) = 2\pi \sum_{i=1}^{l+1} \mathrm{Area}(B_i) = \lambda \mathrm{Vol}(M)$$

in the Einstein case. What we have shown is that the Einstein condition requires the sum of the areas of the bolts of ∂_ϕ to equal the sum of areas of the bolts of ∂_ψ. Moreover, each of these sums is a multiple of the total volume of the manifold. It would be interesting to see if we could devise an analogue of this result in the non-diagonal case.

4.8 Einstein Four Manifolds with an Orthogonally Transitive S^1 Action

Note that in the following discussion, we could consider the more general case of a manifold with one Killing vector whose associated action is orthogonally transitive. Only when we restrict ourselves to the compact case will it be necessary for the action to be that of an S^1 isometry. A manifold with an orthogonally transitive $S^1 = G$ action is a cohomogeneity $(n-1)$ manifold with Riemannian submersion invariant $A \equiv 0$. On such a manifold the metric on an open dense submanifold can be written as $g = \check{g} + f^2 d\varphi^2$ where $\check{g}$ is the induced metric on the quotient space, ϕ is the coordinate on S^1, and f is a function of the base manifold. The quotient space M/G is an $(n-1)$ dimensional manifold and the Einstein condition on M becomes a system of partial differential equations in the

variables of quotient space. As this case is monotypic and $\hat{r} \equiv 0$, the Einstein condition is given by the following two equations

$$\frac{\check{\triangle} f}{f} = -\lambda \tag{4.20}$$

$$\check{r} - \frac{\check{\mathrm{D}}\mathrm{d}f}{f} = \lambda \check{g}. \tag{4.21}$$

where $\check{\mathrm{D}}\mathrm{d}f$ is the Hessian of the metric $\check{g}$. Taking the trace of (4.21), we have

$$\frac{\check{\triangle} f}{f} = \check{s} + (1-n)\lambda.$$

This implies that $\check{s} = (n-2)\lambda$; therefore, the scalar curvature of the base manifold is a constant of the same sign as the Einstein constant. Given the above observations, one can rephrase the Einstein condition in the following way

Proposition 4.20. *The Einstein condition on a manifold M^n with an orthogonally transitive S^1 isometry is equivalent to the equation*

$$\check{\mathrm{D}}\mathrm{d}f - \check{\triangle} f \check{g} - f\check{r} = 0 \tag{4.22}$$

subject to the constraint that $\check{s} = (n-2)\lambda$ is the Einstein constant.

Equation (4.22) also arises in a very different context. For M a compact manifold, the scalar curvature map is a quasi-linear map of second order

$$s : \mathcal{M} \to C^\infty M$$

$$g \mapsto s_g$$

where $\mathcal{M}$ is the space of all Riemannian metrics which is open in $\Gamma^2 M$ the space of all symmetric bilinear differential forms [6].

The scalar curvature map has at every point g of $\mathcal{M}$ a differential

$$s'_g : \Gamma^2 M \to C^\infty M$$

such that

$$s'_g h = -\triangle_g(\mathrm{tr}_g h) + \delta_g(\delta_g h) - g(r_g, h)$$

where δ is the formal adjoint of d.

The formal adjoint of the map s'_g, which is denoted by $(s'_g)^*$, is an overdetermined elliptic operator

$$(s'_g)^* : C^\infty M \to \Gamma^2 M$$

defined by

$$(s'_g)^* f = \mathrm{Dd}_g f - (\triangle_g f)g - f r_g.$$

Returning to equation (4.22), we see that, if M is an Einstein manifold, then f must lie in the kernel of $(\check{s}'_{\check{g}})^*$. We have the following proposition of J. P. Bourguignon

Proposition 4.21. *[8] If* $\ker(\check{s}'_{\check{g}})^* \neq 0$ *then either* $(M/G, \check{g})$*, which is a manifold of dimension* $(n-1)$ *is Ricci-flat and* $\ker(\check{s}'_{\check{g}})^* = \mathbb{R} \cdot \mathbb{I}$ *or the scalar curvature is a strictly positive constant* $\frac{\check{s}_{\check{g}}}{n-2}$

Chapter 5

Kahler Toric Manifolds

Above, in our study of Einstein four manifolds with torus symmetry we noted that the known T^2-invariant Einstein metrics on $S^2 \times S^2$, $\mathbb{C}P^2$, and $\mathbb{C}P^2 \sharp 3\overline{\mathbb{C}P^2}$ are Kahler-Einstein. Compact Kahler manifolds of dimension $2n$ admitting a T^n isometry, of which these Kahler-Einstein metrics are examples, have been studied in great detail by algebraic geometers, differential geometers, and physicists because they have many novel geometric properties. Their geometry can be described using complex coordinates, by cones and fans as in [23], or using symplectic Darboux coordinates, by polytopes in $\mathbb{R}^n$ as in [28], [16], and [1]. Much work has been done to classify all the Kahler-Einstein [43] and constant scalar curvature Kahler metrics [18] on these manifolds but many open questions remain. In this chapter, we will review the basic definitions of Kahler toric geometry following the work of Delzant, Guillemin, and Abreu who discuss Kahler toric geometry in symplectic coordinates and will study the Einstein and extremal Kahler conditions on these manifolds. While this section contains mostly background material for later work, we also present a simplification of the scalar curvature equation written by Abreu in [2].

5.1 Kahler Toric Geometry

Definition 6. *A **Kahler Toric Manifold** is a Kahler manifold (M^{2n}, J, ω) which admits a smooth Hamiltonian T^n action preserving the Kahler structure.*

On the level of Riemannian geometry, such a manifold gives rise to the following submersion (over an open dense set)

$$T^n \to M \to M/T^n \tag{5.1}$$

with torus principal orbits. Note that M/T^n is not a smooth manifold but is a stratified space. In the general Riemannian case, the O'Neill invariants A and T would be nontrivial. However, in the Kahler toric case, we see that the number of invariants of the submersion is greatly reduced.

Theorem 5.1. *On a Kahler toric manifold, the O'Neill invariant A of the associated Riemannian submersion is equal to zero.*

Proof. Take $\{\partial_{\phi_i}\}_{i=1,..,n}$ to be a basis for T^n. Because the torus action is Hamiltonian, there exists a moment map

$$\mu : M \to (\mathfrak{t}^n)^* \cong (\mathbb{R}^n)^*$$

such that

$$d\mu(v)(X) = \omega(X, v) = g(JX, v)$$

for all $v \in TM$ and $X \in \mathfrak{t}^n$. Because J is one-to-one , a vector field v is vertical if and only if $d\mu(v) = 0$. Let $v = \partial_{\phi_i}$ be such a vertical vector field. Therefore, $g(JX, \partial_{\phi_i}) = -g(X, J\partial_{\phi_i}) = 0$ for all $X \in \mathfrak{t}^n$. This implies that $J\partial_{\phi_i}$ is a horizontal vector field. Therefore we may take $\{-J\partial_{\phi_i}\}_{i=1..n}$ to be a basis for the horizontal distribution. To prove that $A = 0$ we will demonstrate that this is in fact a commuting basis implying that the horizontal distribution is integrable.

The vector fields ∂_{ϕ_i} are by assumption automorphisms of the complex structure. Such a vector field satisfies the following

$$\mathcal{L}_{\partial_{\phi_i}} J = 0 \Leftrightarrow [\partial_{\phi_i}, JX] = J[\partial_{\phi_i}, X]$$

for all $X \in TM$. From this equation we deduce immediately that

$$[\partial_{\phi_i}, J\partial_{\phi_j}] = [\partial_{\phi_i}, \partial_{\phi_j}] = 0$$

for all i and j. As J is an integrable complex structure, its Nijenhuis tensor, N_J, vanishes. This implies that

$$N_J(\partial_{\phi_i}, \partial_{\phi_j}) = [J\partial_{\phi_i}, J\partial_{\phi_j}] - J[J\partial_{\phi_i}, \partial_{\phi_j}] - J[\partial_{\phi_i}, J\partial_{\phi_j}] - [\partial_{\phi_i}, \partial_{\phi_j}] = 0.$$

It follows immediately that

$$[J\partial_{\phi_i}, J\partial_{\phi_j}] = 0$$

for all i and j.

Therefore, the basis for the horizontal distribution is integrable and the O'Neill tensor A, which is a measure of the failure of the horizontal distribution to be integrable, is equal to zero. □

The above proof demonstrated that the basis $\{-J\partial_{\phi_i}\}_{i=1,..n}$ is a coordinate basis for the horizontal distribution; we can set $\partial_{u_i} = -J\partial_{\phi_i}$. Because the vector fields ∂_{ϕ_i} and the complex structure J are invariant, we find that $\{\partial_{u_i}\}_{i=1,..n}$ is a basis for the quotient space M/T^n.

We are now in a position to describe the metric on a Kahler toric manifold. First, we set $h_{ij} = g(\partial_{\phi_i}, \partial_{\phi_j})$. Because the metric is invariant under the action of J, we see that $h_{ij} = g(\partial_{u_i}, \partial_{u_j})$. We have shown that a Kahler toric metric, g, is of the form

$$g = \begin{pmatrix} h & 0 \\ 0 & h \end{pmatrix}$$

where $h = (h_{ij})$ and the manifold M has been given the real coordinates $(u_1, ..., u_n, \phi_1, ..., \phi_n)$.

The complex structure J relative to this basis can be written as

$$J = \begin{pmatrix} 0 & -\mathrm{Id}_n \\ \mathrm{Id}_n & 0 \end{pmatrix}$$

Complex coordinates on this manifold can then be defined as $z_i = u_i + i\phi_i$ where the torus acts by $t \cdot (u + i\phi) = u + i(\phi + t)$.

The Kahler form ω is given by $\omega(\cdot, \cdot) = g(J\cdot, \cdot)$ which implies that

$$\omega = \begin{pmatrix} 0 & -h \\ h & 0 \end{pmatrix}.$$

Because the metric is Kahler, there exists, locally, a potential η for the Kahler form. The function η can be chosen to be a function of the u_i's only since the Kahler form is invariant under the action of the torus. It is straightforward to demonstrate that

$$h_{kl} = \frac{\partial^2 \eta}{\partial u_k \partial u_l}.$$

Above, we have described the Kahler toric metric in terms of standard complex coordinates. To do this, we first fixed the complex structure J to be the standard skew-symmetric matrix and arrived at different Kahler metrics by altering the Kahler potential η. Next, we will look at Kahler toric geometry relative to a different set of coordinates by fixing the symplectic structure and varying the complex structure J. We will see that the different complex structures can be parameterized by a 'symplectic' potential. This feature is unique to Kahler toric geometry and the existence of these 'symplectic' coordinates will greatly simplify curvature calculations.

To effect this change in coordinates, we note that $\breve{g} = (h_{ij})$ in the complex coordinate system. Now, the moment map μ of the Hamiltonian torus action is given by $\mu(u, \phi) = \frac{\partial \eta}{\partial u}$ [1]. We construct a new coordinate system $(x_1, ..., x_n, \phi_1, ..., \phi_n)$ on M by defining $x_i = \frac{\partial \eta}{\partial u_i}$ and fixing the coordinate ϕ (effectively we are mapping $\mathbb{R}^n \times T^n$ to $\mathbb{R}^n \times T^n \cong P^o \times T^n$ where P^o is the interior of the image of the moment map). Under the action of this map the Kahler form ω becomes $\omega = dx \wedge d\phi$. The Kahler toric metric has been transformed into Darboux coordinates. In these coordinates, the complex structure becomes

$$J = \begin{pmatrix} 0 & -h \\ h^{-1} & 0 \end{pmatrix}$$

and the metric becomes

$$g = \begin{pmatrix} h^{-1} & 0 \\ 0 & h \end{pmatrix}.$$

In effect, all we have done is to change the coordinate on the quotient space M/T^n so that $\breve{g} = (h_{ij}(x_1, ..., x_n))$.

Guillemin, in [28], demonstrated that h^{-1} itself derives from a potential in symplectic coordinates just as h does in complex coordinates. That is

$$h^{kl} = \frac{\partial^2 \Phi}{\partial x_k \partial x_l}$$

where $\Phi = \Phi(x_1, ..., x_n)$ does not depend on the ϕ_i because the metric is invariant under the torus action. Furthermore, the potentials η and Φ are Legendre dual in the sense that $\eta(u) + \Phi(x) = \sum_j \frac{\partial \eta}{\partial u_j} \frac{\partial \Phi}{\partial x_j}$. We will refer to the coordinate system (x, ϕ) as the symplectic coordinates and to the potential Φ as the symplectic potential to distinguish it from the Kahler potential η. This notation is somewhat misleading in that Φ determines the complex structure while the symplectic structure is fixed in standard Darboux form.

Thus far, all of our calculations have been local. We now wish to study the global structure of a Kahler toric manifold. To do so, we ignore for the moment the complex structure and concentrate only on the symplectic structure.

Let (M^{2n}, ω) be a compact symplectic manifold with an effective Hamiltonian action of T^n. Because the action is Hamiltonian there exists a T^n equivariant moment map $\mu : M \to (\mathfrak{t}^n)^* \cong (\mathbb{R}^n)^*$. It is well-known that the image of such a moment map is a convex polytope $P = \mu(M) \subset \mathbb{R}^n$. Delzant gave the following characterization of the polytopes that are the moment polytopes of symplectic toric manifolds.

Definition 7. *A convex polytope P in $\mathbb{R}^n$ is **Delzant** if*

1. *n edges meet at every vertex p*
2. *the edges meeting at p are rational, i.e. each edge is of the form $p + tv_i$ where $v_i \in \mathbb{Z}^n$*
3. *the $v_1, ..., v_n$ form a basis for $\mathbb{Z}^n$.*

Delzant associated to every polytope satisfying the above condition a symplectic toric manifold (M_P, ω_P). That manifold is the symplectic quotient of $\mathbb{R}^{2d}$ (where d is the number of sides of the polytope) endowed with its standard symplectic structure. This procedure is described in [16] and we will not review it here. In this procedure, every Delzant polytope is in one-to-one correspondence with a symplectic toric manifold.

Viewed in this way, a polytope does not designate a compatible complex structure and therefore does not itself determine a Kahler toric metric. That said, one can associate to each polytope a 'canonical' compatible complex structure to the toric symplectic manifold. This is done by taking $\mathbb{R}^{2d} \cong \mathbb{C}^d$ with its standard linear complex structure. This complex structure descends via the symplectic quotient to a compatible complex structure on the symplectic toric manifold (M_P, ω_P). One can therefore assign a 'canonical' Kahler toric metric to each polytope. It is important to emphasize that this 'canonical' metric is not the only Kahler toric metric whose image under the

moment map μ is the given polytope P. A different compatible T^n-invariant complex structure would give a different Kahler toric metric.

Delzant was able to determine the 'symplectic' potential Φ_P of the 'canonical' Kahler toric metric associated to a polytope P. After reviewing the construction of this potential, we discuss the work of Abreu which characterizes all other Kahler toric metrics that can be associated to polytope P.

One can describe a Delzant polytope by a set of inequalities $\langle x, \mu_m \rangle \geq \lambda_m$ for $m = 1, ..., d$ where d is the number of faces of the polytope, λ_m is a constant, and μ_m is a primitive element of the lattice $\mathbb{Z}^n \subset \mathbb{R}^n$. Define the function $l_m(x)$ by

$$l_m(x) = \langle x, \mu_m \rangle - \lambda_m.$$

Delzant showed that the 'symplectic' potential

$$\Phi_P = \frac{1}{2} \sum_{m=1}^{d} l_m \log(l_m)$$

determines a smooth Kahler toric metric on the manifold M with symplectic structure defined by the polytope P. Note that there are many other Kahler metrics associated to a given polytope just as there are many complex structures compatible with the symplectic structure defined by the polytope. It is possible to choose different potentials which will give rise to different complex structures.

It is worthwhile to pause to determine the significance of the constants μ_m and λ_m. Given a polytope defined by a set of inequalities, what happens if we change the values of a particular μ_m or a particular λ_m? If one chooses a different primitive element, μ_m, of the integer lattice, there are two possibilities. First, the resulting polytope may no longer be Delzant and may not give rise to a smooth manifold. Second, if the resulting polytope is Delzant then the corresponding symplectic structure is altered. The associated differentiable manifold may be the same or different but the symplectic structure will have been modified. On the other hand, if we perturb slightly the value of λ_m the corresponding differentiable manifold is unchanged while the cohomology class of the associated symplectic form ω is perturbed slightly. Therefore, fixing a polytope $\{\mu_m, \lambda_m\}$ is equivalent to fixing a manifold, a symplectic structure on the manifold, and the Kahler class of any metric on the manifold. Varying the potential Φ while leaving fixed the polytope P changes the complex structure of the Kahler metric [28].

As mentioned above, the 'canonical' Kahler metric constructed by Delzant is not the only Kahler metric which can be associated to a given polytope. Abreu found a means of describing all the compatible complex structures.

Theorem 5.2. *[1] Let (M_P, ω_P) be a toric symplectic manifold associated to a Delzant polytope $P \subset \mathbb{R}^n$ and J any compatible complex structure. Then J is determined by a potential Φ of the form*

$$\Phi = \Phi_P + \Psi$$

where Ψ *is smooth on* P*, the matrix* $h^{-1} = \left(\frac{\partial^2 \Phi}{\partial x^k \partial x^l}\right)$ *is positive definite on* P^0 *and has determinant*

$$\det(h^{-1}) = \left[\delta(x) \prod_{m=1}^{d} l_m(x)\right]^{-1}$$

where δ*, which depends on* Ψ*, is smooth and strictly positive on the whole of* P*.*

To summarize, any Kahler toric metric can be described by a symplectic potential which is the sum of the 'canonical' potential associated to the polytope P (which is determined by the symplectic structure) and a function satisfying the conditions of theorem (5.2).

5.2 Curvature Equations and Conditions

In this section, we will review the expression of the scalar curvature of a Kahler toric manifold in symplectic coordinates discovered by Abreu [1]. We will also present a slightly simplified version of the scalar curvature equation. Next, we will study the Extremal Kahler condition and will derive the full Einstein equation on a Kahler toric manifold

5.2.1 The Scalar Curvature Equation

Before we calculate the scalar curvature in symplectic coordinates, recall that the matrix h_{kl}, when expressed in complex coordinates (u, ϕ), is derived from a potential η such that $h_{kl} = \frac{\partial^2 \eta}{\partial u^k \partial u^l}$. Conversely, when expressed in symplectic coordinates, (x, ϕ), the inverse matrix h^{kl} is derived from a potential Φ such that $h^{kl} = \frac{\partial^2 \Phi}{\partial x^k \partial x^l}$. Given the definitions of x and u of the previous section we readily derive the following relations

$$h^{kl} \frac{\partial}{\partial u^l} = \frac{\partial}{\partial x^k} \tag{5.2}$$

and

$$h_{kl} \frac{\partial}{\partial x^k} = \frac{\partial}{\partial u^l}. \tag{5.3}$$

Note: Here we are using a modified Einstein summation convention. *Whenever* an index is repeated twice on the same side of an equality, we sum over that index. We will use this modified convention consistently throughout so there should be no chance for confusion with the typical convention in which one only sums over repeated indices when one of indices is raised and one is lowered.

It is well-known that in complex coordinates (u, ϕ), the scalar curvature of a Kahler toric manifold can be written as

$$S = -h^{ij} \frac{\partial^2 \log(\det(h))}{\partial u^i \partial u^j}. \tag{5.4}$$

The following theorem gives an expression for the scalar curvature in terms of the symplectic coordinates.

Theorem 5.3. *[2] The scalar curvature of a Kahler toric manifold can be expressed, relative to the coordinates (x, ϕ) as*

$$S = -\frac{\partial^2 h_{ij}}{\partial x^i \partial x^j} \tag{5.5}$$

where $h_{ij} = h_{ij}(x_1, ..., x_n)$ as above.

Proof. Using equation (5.3) we see that

$$S = -h^{ij}\frac{\partial^2 \log(\det(h))}{\partial u^i \partial u^j} = -\frac{\partial}{\partial x^i}\left(\frac{\partial \log(\det(h))}{\partial u^i}\right)$$

$$= \frac{\partial}{\partial x^i}\left(\frac{\partial \log(\det(h^{-1}))}{\partial u^i}\right) = \frac{\partial}{\partial x^j}\left(h_{ij}\frac{\partial \log(\det(h^{-1}))}{\partial x^i}\right).$$

Following Abreu, we make use of the following identity: For $V = (v^{ab})$ and $Y = (y^{ab})$ where V and Y are symmetric and Y is positive definite then

$$v^{ab}\frac{\partial}{\partial y^{ab}}(\log(\det Y)) = \text{trace}(VY^{-1}).$$

Using this identity for $Y = h^{ab}$ and $V = h_{ij}$ we have

$$S = \frac{\partial}{\partial x^j}\left(h_{ij}\frac{\partial \log(\det(h^{-1}))}{\partial x^i}\right) = \frac{\partial}{\partial x^j}\left(h_{ij}\frac{\partial \log(\det(h^{-1}))}{\partial h^{ab}}\frac{\partial h^{ab}}{\partial x^i}\right)$$

$$= \frac{\partial}{\partial x^j}\left(h_{ij}\frac{\partial h^{ab}}{\partial x^i}h_{ab}\right) = \frac{\partial}{\partial x^j}\left(h_{ij}\frac{\partial h^{ib}}{\partial x^a}h_{ab}\right) = -\frac{\partial}{\partial x^j}\left(\frac{\partial h_{ij}}{\partial x^a}h^{ib}h_{ab}\right)$$

this implies immediately that

$$S = -\frac{\partial^2 h_{ij}}{\partial x^i \partial x^j}.$$

□

We are able to give a modified version of the Abreu formula which is somewhat simpler and will prove useful later. First, we will require the following lemma.

Lemma 5.4. *On a Kahler toric manifold, setting $h_{ij} = \det(\text{h})\mathcal{M}_{\text{ij}}$,*

$$\frac{\partial}{\partial x^i}\left(\frac{1}{\det(h)}h_{ij}\right) = \frac{\partial \mathcal{M}_{ij}}{\partial x^i} = 0 \tag{5.6}$$

for all j.

Proof. Note that $\mathcal{M}$ is the adjugate matrix of h^{-1}. Recall that in symplectic coordinates h^{kl} derives from a potential ϕ. After expressing, h^{-1} in terms of this potential, and calculating the adjugate matrix $\mathcal{M}$ it is easy to check that equation (5.6) holds for $n = 1$ or 2. In the $n = 1$ case, h^{11} is the only term and $\mathcal{M}_{11} = 1$. Equation (5.6) holds easily. In the $n = 2$ case,

$$h^{-1} = \begin{pmatrix} \Phi_{x_1x_1} & \Phi_{x_1x_2} \\ \Phi_{x_1x_2} & \Phi_{x_2x_2} \end{pmatrix}$$

and

$$\mathcal{M} = \begin{pmatrix} \Phi_{x_2x_2} & -\Phi_{x_1x_2} \\ -\Phi_{x_1x_2} & \Phi_{x_1x_1} \end{pmatrix}.$$

Equation (5.6) again follows immediately. A straightforward but cumbersome induction argument shows that equation (5.6). We omit the proof here.

□

Theorem 5.5. *The scalar curvature of a Kahler toric manifold can be written in coordinate* (x, ϕ) *as*

$$S = -\frac{1}{\det(h)} h_{ij} \frac{\partial^2 \det(h)}{\partial x^i \partial x^j}. \tag{5.7}$$

Proof. The scalar curvature can be written as

$$S = -\frac{\partial^2 h_{ij}}{\partial x^i \partial x^j} = -\frac{\partial^2((\det(h))\mathcal{M}_{ij})}{\partial x^i \partial x^j} = -\frac{\partial}{\partial x^i}\left(\mathcal{M}_{ij}\frac{\partial \det(h)}{\partial x^j}\right) - \frac{\partial}{\partial x^i}\left(\det(h)\frac{\partial \mathcal{M}_{ij}}{\partial x^j}\right).$$

From the proof of Abreu's formula we see that

$$S = -\frac{\partial}{\partial x^j}\left(h_{ij}\frac{\partial \log(\det(h))}{\partial x^i}\right) = -\frac{\partial}{\partial x^i}\left(\mathcal{M}_{ij}\frac{\partial \det(h)}{\partial x^j}\right).$$

Using equation(5.6), the scalar curvature becomes

$$S = -\frac{\partial}{\partial x^i}\left(\mathcal{M}_{ij}\frac{\partial \det(h)}{\partial x^j}\right) = -\frac{\partial \mathcal{M}_{ij}}{\partial x^i}\frac{\partial \det(h)}{\partial x^j} - \mathcal{M}_{ij}\frac{\partial^2 \det(h)}{\partial x^i \partial x^j} = -\mathcal{M}_{ij}\frac{\partial^2 \det(h)}{\partial x^i \partial x^j}.$$

□

Equation (5.7) and equation(5.4) are very similar in form; this similarity is obscured by the form of Abreu because the equation he derived contains elements which cancel upon inspection. Therefore, it appears that the form of the scalar curvature function given by (5.7) will be more appropriate for performing calculations.

Below, we will make frequent use of equation (5.6) while analyzing both Kahler toric and fiberwise Kahler toric manifolds.

Before beginning our discussion of the extremal Kahler condition, we first calculate, for use later, the Laplacians of the Kahler toric metric, g, and of the quotient metric, $\check{g}$. The general formula for the Laplacian of a metric g is

$$\Delta f = \frac{1}{\sqrt{\det(g)}}\frac{\partial}{\partial y^j}\left(\sqrt{\det(g)}g^{ij}\frac{\partial f}{\partial y^i}\right)$$

where $(y_1, ..., y_m)$ is a basis for an m-dimensional manifold. In symplectic coordinates, a Kahler toric metric g can be written as $g = \begin{pmatrix} h^{-1} & 0 \\ 0 & h \end{pmatrix}$. This implies that $\det(g) = 1$. Restricted to invariant functions, the Laplacian becomes

$$\Delta f = \frac{\partial}{\partial x^j}\left(h_{ij}\frac{\partial f}{\partial x^i}\right) \tag{5.8}$$

for $f = f(x_1, ..., x_n)$. Defining the matric $\mathcal{M}$ as above and recalling equation (5.6), we have

$$\Delta f = \frac{\partial}{\partial x^j}\left(\det(h)\mathcal{M}_{ij}\frac{\partial f}{\partial x^i}\right) = \det(h)\mathcal{M}_{ij}\frac{\partial^2 f}{\partial x^i \partial x^j} + \mathcal{M}_{ij}\frac{\partial \det(h)}{\partial x^i}\frac{\partial f}{\partial x^j} + \det(h)\frac{\partial \mathcal{M}_{ij}}{\partial x^i}\frac{\partial f}{\partial x^j}.$$

From this we immediately have that

$$\Delta f = h_{ij}\frac{\partial^2 f}{\partial x^i \partial x^j} + \frac{1}{\det(h)}h_{ij}\frac{\partial \det(h)}{\partial x^i}\frac{\partial f}{\partial x^j}. \tag{5.9}$$

Next, we calculate $\check{\Delta}$, the Laplacian of the quotient metric $\check{g} = (h^{-1})$. Using the general formula for the Laplace operator, we see that

$$\check{\Delta} f = \frac{1}{\sqrt{\det(h^{-1})}}\frac{\partial}{\partial x^j}\left(\sqrt{\det(h^{-1})}h_{ij}\frac{\partial f}{\partial x^i}\right) = \sqrt{\det(h)}\frac{\partial}{\partial x^j}\left(\frac{1}{\sqrt{\det(h)}}\det(h)\mathcal{M}_{ij}\frac{\partial f}{\partial x^i}\right)$$

Using equation (5.6)

$$\check{\Delta} f = \det(h)\mathcal{M}_{ij}\frac{\partial^2 f}{\partial x^i \partial x^j} + \frac{1}{2}\mathcal{M}_{ij}\frac{\partial \det(h)}{\partial x^j}\frac{\partial f}{\partial x^i}.$$

Therefore,

$$\check{\Delta} f = h_{ij}\frac{\partial^2 f}{\partial x^i \partial x^j} + \frac{1}{2}\frac{1}{\det(h)}h_{ij}\frac{\partial \det(h)}{\partial x^j}\frac{\partial f}{\partial x^i}. \tag{5.10}$$

This formula allows us to prove the following proposition

Proposition 5.6. *On a Kahler toric manifold with the metric defined in symplectic coordinates as above, we have the following identity*

$$\check{\Delta}\left(\sqrt{\det(h)}\right) = -\left(\frac{S}{2}\right)\sqrt{\det(h)}. \tag{5.11}$$

Therefore, on a manifold of constant scalar curvature S the function $\sqrt{\det(h)}$ is an eigenfunction of the quotient Laplacian with eigenvalue $\frac{S}{2}$.

Proof. Equation (5.10) acting on function $\sqrt{\det(h)}$ implies that

$$\check{\Delta}\sqrt{\det(h)} = h_{ij}\frac{\partial^2 \sqrt{\det(h)}}{\partial x^i \partial x^j} + \frac{1}{2}\frac{1}{\det(h)}h_{ij}\frac{\partial \det(h)}{\partial x^j}\frac{\partial \sqrt{\det(h)}}{\partial x^i}.$$

Expanding the derivatives and noting the equation (5.7) for the scalar curvature finishes the proof. □

5.2.2 The Extremal Kahler Condition

As mentioned in the introduction, Einstein metrics are critical points of the scalar curvature functional $\mathbf{S} : g \to \int_M S_g \mathrm{vol}_g$. Calabi introduced another functional on metrics restricted to a fixed Kahler class. The functional is defined to be

$$\mathbf{Ss} : g \to \int_M s_g^2 \mathrm{vol}_g. \tag{5.12}$$

Definition 8. *Critical points of the functional **Ss** are called **Extremal Kahler Metrics**.*

By deriving the Euler-Lagrange equation for this functional, Calabi proved the following theorem.

Theorem 5.7. *[10] A Kahler metric is extremal if and only if the gradient ∇S is a holomorphic vector field.*

It follows immediately from this theorem that any constant scalar curvature metric is in fact extremal Kahler. Also, Calabi found that, if one exists, an extremal Kahler metric is unique within a given Kahler class [10].

What is the extremal Kahler condition on a Kahler toric manifold? In complex coordinates, the metric can be written as $g = \begin{pmatrix} h & 0 \\ 0 & h \end{pmatrix}$ relative to coordinates (u, ϕ). The complex coordinates themselves can be written as $z_i = u_i + i\phi_i$. Since the metric is invariant under the action of T^n, the scalar curvature, S, is a function of the u_i only. The gradient of S can be written

$$\nabla S = \left(h^{kl} \frac{\partial S}{\partial u^l} \right) \frac{\partial}{\partial z_k}.$$

Since the coefficients, $h^{kl} \frac{\partial S}{\partial u^l}$, of the above equation are real, the vector field ∇S is holomorphic if and only if these coefficients are constant. The extremal Kahler condition is equivalent to the following system of equations:

$$h^{kl} \frac{\partial S}{\partial u^l} = \frac{\partial S}{\partial x^k} = \alpha_k$$

where the α_i are constants.

The above proves the following theorem due to Abreu:

Theorem 5.8. *[2] A Kahler Toric metric is Extremal Kahler if and only if the scalar curvature satisfies*

$$S = \sum_{k=1}^{n} \alpha_k x_k + \beta \tag{5.13}$$

where β and the α_k are constants and the x_i are as defined above.

Later, we will return to the question of finding extremal Kahler toric metrics and we discuss the intimate connection between extremal Kahler and Kahler-Einstein four manifolds discovered by Derdzinski.

5.2.3 The Einstein Condition

To calculate the Einstein condition on a Kahler toric manifold in symplectic coordinates, we begin by looking at the Einstein condition in complex coordinates $(z_1, ..., z_n)$ where $z_i = u_i + i\phi_i$. On a Kahler manifold, the Ricci form, $\rho(\cdot, \cdot) = r(J\cdot, \cdot)$, is given by the expression

$$\rho = -i\partial\overline{\partial}\log(\det(g_{a\overline{\beta}})).$$

The Kahler-Einstein condition becomes equivalent to the equality

$$\rho = \lambda\omega.$$

It is straightforward to evaluate this equation on a Kahler toric manifold. A Kahler-Einstein toric manifold must satisfy the equation

$$\frac{\partial^2 \log(\det(h))}{\partial u^i \partial u^j} = -2\lambda h_{ij} \tag{5.14}$$

for all i and j, where λ is the Einstein constant.

Converting this equation into symplectic coordinates gives the following theorem.

Theorem 5.9. *On a Kahler toric manifold, the Einstein condition, when expressed in symplectic coordinates (x, ϕ) becomes the equation*

$$h_{ik} \frac{\partial}{\partial x^k} \left(h_{lj} \frac{\partial \log(\det(h))}{\partial x^l} \right) = -2\lambda h_{ij}. \tag{5.15}$$

for all i and j.

Proof. From equations (5.14) and (5.3), this formula follows immediately. □

This form of the Einstein condition is somewhat cumbersome and we will simplify it by making use of the adjugate matrix $\mathcal{M}$ defined above.

Corollary 5.10. *The Einstein condition on a Kahler toric manifold is equivalent to the equation*

$$\mathcal{M}_{ik} \frac{\partial}{\partial x^k} \left(\mathcal{M}_{lj} \frac{\partial \det(h)}{\partial x^l} \right) = -2\lambda \mathcal{M}_{ij} \tag{5.16}$$

for all i and j.

Proof. Recall that the matrix $\mathcal{M}$ is defined by the expression $h_{kl} = \det(h)\mathcal{M}_{kl}$. Using the equation from the previous theorem, we have

$$\det(h)\mathcal{M}_{ik} \frac{\partial}{\partial x^k} \left(\det(h)\mathcal{M}_{lj} \frac{\partial \log(\det(h))}{\partial x^l} \right) = -2\lambda \det(h)\mathcal{M}_{ij}.$$

The Corollary follows immediately.

□

Chapter 6

Einstein and Extremal Kahler Metrics on Kahler Toric Manifolds

In this chapter, we study the Einstein and extremal Kahler conditions on Kahler toric manifolds, with particular emphasis on real dimensions four and six. First, we demonstrate how to integrate the equations of the Einstein condition to obtain a Legendre transform of the Monge-Ampere equations. Next, we study the Kahler-Einstein condition on $\mathbb{C}P^2\sharp 3\overline{\mathbb{C}P^2}$ and write down explicitly the Kahler-Einstein metric on $\mathbb{P}(\mathcal{O}_{\mathbb{C}P^1\times\mathbb{C}P^1}\oplus\mathcal{O}_{\mathbb{C}P^1\times\mathbb{C}P^1}(1,-1))$ in symplectic coordinates. Finally, after reviewing the construction of extremal Kahler metrics on $\mathbb{C}P^2\sharp\overline{\mathbb{C}P^2}$, we construct a new two parameter family of extremal Kahler metrics on $\mathbb{P}(\mathcal{O}_{\mathbb{C}P^1\times\mathbb{C}P^1}\oplus\mathcal{O}_{\mathbb{C}P^1\times\mathbb{C}P^1}(1,-1))$ which contains two one-parameter families of constant scalar curvature metrics.

6.1 Kahler-Einstein Toric Manifolds

Having calculated the Einstein condition on a Kahler toric manifold, we know demonstrate how to integrate those equations, and reduce the Einstein condition to a single equation involving only the potential and its first and second derivatives.

6.1.1 Integration of the Einstein Equations

As calculated above, the Einstein condition on a Kahler toric metric in arbitrary dimension can be written as

$$\mathcal{M}_{ik}\frac{\partial}{\partial x^k}\left(\mathcal{M}_{lj}\frac{\partial\det(h)}{\partial x^l}\right)=-2\lambda\mathcal{M}_{ij}$$

where $\mathcal{M}=\frac{1}{\det(h)}h$. Note that on the interior of the polytope $\mathcal{M}$ is a positive definite matrix and therefore invertible. Multiplying both sides of this equations by $\mathcal{M}^{mi}$ gives

$$\mathcal{M}^{mi}\mathcal{M}_{ik}\frac{\partial}{\partial x^k}\left(\mathcal{M}_{lj}\frac{\partial\det(h)}{\partial x^l}\right)=-2\lambda\mathcal{M}^{mi}\mathcal{M}_{ij}.$$

This implies that

$$\delta^m_k\frac{\partial}{\partial x^k}\left(\mathcal{M}_{lj}\frac{\partial\det(h)}{\partial x^l}\right)=-2\lambda\delta^m_j.$$

Integrating these equations gives

$$\mathcal{M}_{lj}\frac{\partial \det(h)}{\partial x^l} = -2\lambda x_j + C_j$$

for all $1 \leq j \leq n$ where $2n$ is the dimension of the Kahler toric manifold and the c_j are constants. With the first integration accomplished, we attempt the second.

We begin, as before, by multiplying both sides of the equation by $\mathcal{M}^{mj}$ giving

$$\mathcal{M}^{mj}\mathcal{M}_{lj}\frac{\partial \det(h)}{\partial x^l} = \mathcal{M}^{mj}(-2\lambda x_j + C_j).$$

Since $\mathcal{M} = \frac{1}{\det(h)}h$, $\mathcal{M}^{ml} = (\det(h))h^{ml}$. Substituting this into the above equation gives

$$\delta_l^m \frac{\partial \log(\det(h))}{\partial x^l} = (-2\lambda x_j + C_j)h^{mj}.$$

We recall that $h^{mj} = \frac{\partial^2 \Phi}{\partial x^m \partial x^j}$ where Φ is the symplectic potential. This implies that

$$\frac{\partial \log(\det(h))}{\partial x^l} = (-2\lambda x_j + C_j)\frac{\partial^2 \Phi}{\partial x^l \partial x^j}$$

Integrating this equation, we have the following theorem.

Theorem 6.1. *Let (M^{2n}, g, J) be a Kahler toric manifold with symplectic potential $\Phi = \Phi(x_1, ..., x_n)$ as above. The Einstein condition is equivalent to the equation*

$$\log(\det(h)) = \sum_{j=1}^{n}(-2\lambda x_j + C_j)\frac{\partial \Phi}{\partial x^j} + 2\lambda\Phi + E \tag{6.1}$$

where E and the C_j are constants and h^{-1} is the matrix whose ijth entry is $\frac{\partial^2 \Phi}{\partial x^i \partial x^j}$.

As in the four dimensional case, we can perform these integrations using the complex coordinates $(u_1, ..., u_n)$ as well. In these coordinates, the Einstein condition can be written as

$$\frac{\partial^2 \log(\det(h))}{\partial u^i \partial u^j} = -2\lambda h_{ij}$$

for all $1 \leq i, j \leq n$. Noting that $\frac{\partial^2 \eta}{\partial x^i \partial x^j} = h_{ij}$ and integrating the above equation gives the following theorem.

Theorem 6.2. *Let (M^{2n}, g, J) be a Kahler toric manifold with complex potential $\eta = \eta(u_1, ..., u_n)$ as above. The Einstein condition is equivalent to the equation*

$$\log(\det(h)) = -2\lambda\eta + \sum_{j=1}^{n} C_j u_j + E \tag{6.2}$$

where E and the C_j are constants.

This is a well-known expression for the Einstein condition in complex coordinates on a Kahler toric manifold. This reduction of the Kahler-Einstein equations to the real Monge-Ampere equations in the Kahler toric case was studied by Calabi in [9].

Returning to equation (6.1), we want to determine the significance of the constants C_i. As a first step, we fix a scale for the metric by setting the Einstein constant, λ, equal to one. A polytope is defined in the symplectic picture by a collections of hyperplanes. Of course, translating in the coordinates x_i does not alter the metric. Neither does transforming the coordinates by an action of $SL(n,\mathbb{Z})$. We can use these two symmetries to fix the location of n sides of the polytope. Pick any vertex p of the polytope. By translation in the x_i we can set $p = (-1, ..., -1)$. Because the polytope must be Delzant to define a Kahler toric manifold we know that there are exactly n faces meeting at p. By an action of $SL(n,\mathbb{Z})$, we can set p to be the intersection of the planes $1 + x_i = 0$ for all i.

Abreu demonstrated that the potential of any Kahler metric whose polytope contains a vertex of this type can be written as

$$\Phi = \frac{1}{2}\sum_{i=1}^{n}(1+x_i)\log(1+x_i) + \Psi$$

where Ψ and its derivatives are smooth in a small open neighborhood of the point $p = (-1, ..., -1)$. Consider the a point x such that $-1 < x_i < -1+\epsilon$ for $\epsilon > 0$. Leave x_i fixed for $i > 1$. As $x_1 \longrightarrow -1$, $\frac{\partial^2\Phi}{\partial x^1 \partial x^1} \longrightarrow \frac{1}{2}\frac{1}{1+x_1}$ and $\frac{\partial^2\Phi}{\partial x^i \partial x^j}$ approaches a constant if i or j does not equal one. Furthermore, a straightforward calculation shows that $\frac{1}{\det(h)}\frac{\partial \det(h)}{\partial x_1} \longrightarrow \frac{1}{1+x_1}$ as $x_1 \to -1$.

Consider the following consequence of the Einstein condition when $\lambda = 1$

$$\frac{\partial \log(\det(h))}{\partial x^1} = (-2x_j + C_j)\frac{\partial^2\Phi}{\partial x^1 \partial x^j}.$$

As $x_1 \longrightarrow -1$, the left-hand side of this equation approaches $\frac{1}{1+x_1}$ and the right-hand side approaches $(2 + C_1)\frac{1}{2}\frac{1}{1+x_1}$. Equating these implies that $C_1 = 0$. Performing a similar calculation as $x_i \longrightarrow -1$ for $i \neq 1$, we deduce immediately that $C_i = 0$ for all i. The above proves the following proposition.

Proposition 6.3. *Let (M^{2n}, g, J) be a Kahler-Einstein toric manifold with Einstein constant $\lambda = 1$. Through translation and an $SL(n,\mathbb{Z})$ transformation, we can without loss of generality assume that the polytope has faces $x_i = -1$. Given these assumptions, the constants C_i of the Einstein condition (6.1) vanish and the Einstein condition becomes*

$$-\log\left(\det(h^{-1})\right) = \sum_{j=1}^{n} -2x_j\frac{\partial\Phi}{\partial x^j} + 2\Phi + E. \tag{6.3}$$

Below, we will assume that our Kahler Einstein metrics satisfy the conditions of the preceding proposition.

Before specializing to the compact case, we pause to mention work done in the case in which the Kahler toric manifold is also hyperkahler. Because the Einstein constant is zero, the manifold must be noncompact. In this, the real Monge-Ampere equations reduce further to a linear system. For a more detailed account, see [29].

6.1.2 Holomorphic Obstructions

Any Kahler-Einstein toric metric on a compact manifold (M, g, J) must have a positive Einstein constant. That is, M must be a Fano manifold with $c_1(M) > 0$. There are two known holomorphic obstructions to the existence of Einstein metrics on Fano manifolds. The first is due to Matsushima and involves the Lie algebra of holomorphic vector fields which we denote by $\mathfrak{h}(M)$.

Theorem 6.4. *[46] If a compact Fano manifold N admits a Kahler-Einstein metric with positive Einstein constant, then $\mathfrak{h}(N)$ is reductive.*

The second obstruction is due to Futaki. Let N^{2n} be a Fano manifold. One can find a Kahler metric whose Kahler form ω is contained in the first Chern class: $\omega \in c_1(N)$. Let ρ_ω be the Ricci form of the Kahler metric. It is well-known that $\rho_\omega \in c_1(N)$ also. Therefore, there exists a function F_ω such that

$$\rho_\omega - \omega = \partial\bar{\partial} F_\omega.$$

Futaki defined the following map $\mathcal{F} : \mathfrak{h}(N) \to \mathbb{C}$ by the equation

$$\mathcal{F}(X) = \int_N X F_\omega \omega^n$$

for all $X \in \mathfrak{h}(N)$. The map $\mathcal{F}$ is independent of the choice of $\omega \in c_1(N)$ [24]. The map $\mathcal{F}$ is known as the Futaki invariant.

The following theorem demonstrates that the Futaki invariant is an obstruction to the existence of Kahler-Einstein metrics on Fano manifolds.

Theorem 6.5. *[24] If N admits a Kahler-Einstein metric then $\mathcal{F} \equiv 0$.*

This condition can also be expressed in terms of the moment map $\mu : N \to \mathfrak{h}^*$. Let μ_X be defined by the relation $\langle \mu(p), X \rangle = \mu_X(p)$ for all $p \in N$ and $X \in \mathfrak{h}(N)$.

Corollary 6.6. *[25] The Futaki invariant satisfies*

$$\mathcal{F}(X) = -\int_N \mu_X \omega^n.$$

That is, the Futaki invariant vanishes when the barycenter of the moment map lies at zero. The above corollary allows us to calculate the Futaki invariant for a Kahler toric manifold (M^{2n}, g, J). Because the Kahler structure is T^n invariant, we can, without loss of generality, choose F_ω to be a function of the u_i only. (Here is the u_i are the real part of the complex coordinates given above.) Consider the (real) holomorphic vector field $\frac{\partial}{\partial u^i}$. As discussed above, we know that $\mu_{\frac{\partial}{\partial u^i}} = \frac{\partial \eta}{\partial u^i} = x_i$. Furthermore, we recall that the 'symplectic' coordinates (x, ϕ) are Darboux coordinates so $\omega^n = dx_1 \wedge ... \wedge dx_n \wedge d\phi_1 \wedge ... \wedge d\phi_n$. We calculate that

$$\mathcal{F}(\frac{\partial}{\partial u^i}) = -\int_M x_i \omega^n = -(2\pi)^n \int_\Delta x_i dx$$

for all i, where Δ is the polytope defined in symplectic coordinates x_i. We have the following proposition

Proposition 6.7. *[43] If (M, g, J) is a compact Kahler-Einstein toric manifold then*

$$\int_{\Delta} x_i dx = 0 \tag{6.4}$$

for all i.

6.1.3 Kahler-Einstein Toric Manifolds in Dimensions Two and Four

In dimension two, the only compact Kahler toric manifold is $\mathbb{C}P^1$. The canonical Kahler metric is defined by the potential

$$\Phi = \frac{1}{2}[(1+x_1)\log(1+x_1) + (1-x_1)\log(1-x_1)].$$

It is easily seen that this potential is Einstein. Therefore, the Kahler-Einstein condition on two-dimensional Kahler toric manifolds (and indeed all two dimensional manifolds) is completely understood.

However, there remains one open problem concerning compact Kahler-Einstein toric manifolds in dimension four. It is well-known [61] that the only Kahler toric Fano manifolds whose Lie algebra of holomorphic vector fields is reductive and whose Futaki invariant vanishes are $\mathbb{C}P^2$, $\mathbb{C}P^1 \times \mathbb{C}P^1$, and $\mathbb{C}P^2 \sharp 3\overline{\mathbb{C}P^2}$. The last is the blow-up of $\mathbb{C}P^2$ at three points in general position. The potentials of the Kahler-Einstein metrics on $\mathbb{C}P^2$ and $\mathbb{C}P^1 \times \mathbb{C}P^1$ are both known explicitly. They are

$$\Phi = \frac{1}{2}[(1+x_1)\log(1+x_1) + (1+x_2)\log(1+x_2) + (1-x_1-x_2)\log(1-x_1-x_2)]$$

and

$$\Phi = \frac{1}{2}[(1+x_1)\log(1+x_1) + (1+x_2)\log(1+x_2) + (1-x_1)\log(1-x_1) + (1-x_2)\log(1-x_2)]$$

respectively. The Kahler-Einstein potential are simply the potentials of the canonical Kahler metric on the polytopes defined by these potentials.

While $\mathbb{C}P^2 \sharp 3\overline{\mathbb{C}P^2}$ is known to admit a Kahler-Einstein toric metric (cf. [55] and [61]) the potential is not known explicitly. Much, however, is known about this metric. From the theorem of Abreu (5.2) given above, we know that the potential of the Kahler-Einstein metric can be written as $\Phi = \Phi_0 + f(x_1, x_2)$ where Φ_0 is the potential of the canonical Kahler metric and f is a function whose second derivatives are smooth on the polytope. For ease of presentation, set $x_1 = x$ and $x_2 = y$. We can write Φ_0 as

$$\Phi_0 = \frac{1}{2}[(1+x)\log(1+x) + (1+y)\log(1+y) + (1-x-y)\log(1-x-y)$$
$$+(1-x)\log(1-x) + (1-y)\log(1-y) + (1+x+y)\log(1+x+y)].$$

This potential Φ_0 is associated to polytope P_0 pictured in the figure below.

Figure 1: *Polytope associated to* Φ_0 *on* $\mathbb{C}P^2\sharp 3\overline{\mathbb{C}P^2}$



Substituting this expression for Φ into equation (6.3), the Einstein condition becomes

$$\log(1-x^2+1-y^2+1-(x+y)^2+[1-x^2+1-(x+y)^2](1-y^2)f_{yy}+[1-y^2+1-(x+y)^2](1-x^2)f_{xx} \quad (6.5)$$

$$-2(1-x^2)(1-y^2)f_{xy}+(1-x^2)(1-y^2)(1-(x+y)^2)(f_{xx}f_{yy}-f_{xy}^2))=2xf_x-2f+2yf_y-E.$$

This equation, being equivalent to the Monge-Ampere equation, is highly non-linear and cannot be integrated by any known techniques. However, one could try different types of functions in an attempt to find the solution to (6.5) that corresponds to the Kahler-Einstein metric on $\mathbb{C}P^2\sharp 3\overline{\mathbb{C}P^2}$.

Let f be a function of only one coordinate say x. We could ask whether there are any solutions to equation (6.5) such that $f=f(x)$. If f does not depend on y then (6.5) becomes

$$\log(1-x^2+1-y^2+1-(x+y)^2+[1-y^2+1-(x+y)^2](1-x^2)f_{xx})=2xf_x-2f-E.$$

The right-hand side of this equation depends only on x, therefore the same must be true of the left-hand side. The left-hand side of the equations is a function of x only if and only if

$$f_{xx}=\frac{1}{x^2-1}.$$

The function f can then be written as

$$f=\frac{1}{2}[(x-1)\log(1-x)-(1+x)\log(1+x)].$$

The resulting potential Φ becomes

$$\Phi=\frac{1}{2}[(1+y)\log(1+y)+(1-x-y)\log(1-x-y)+(1-y)\log(1-y)+(1+x+y)\log(1+x+y)]$$

which is the potential for the Kahler-Einstein metric on $S^2\times S^2$. Therefore, the Siu metric does not have potential the form $\Phi=\Phi_0+f(x)$.

Alternatively, one could look for solutions to equation (6.5) in which $f = p(x)+q(y)$. Substituting this expression for f into equation (6.5) gives

$$\log(1-x^2+1-y^2+1-(x+y)^2+[1-x^2+1-(x+y)^2](1-y^2)q_{yy}+[1-y^2+1-(x+y)^2](1-x^2)p_{xx}$$

$$+(1-x^2)(1-y^2)(1-(x+y)^2)p_{xx}q_{yy} = 2xp_x - 2p - 2q + 2yq_y - E.$$

Evaluating this equation at $y = 1$ and $y = -1$ gives

$$\log(1-x^2+1-(x+1)^2+[1-(x+1)^2](1-x^2)p_{xx}) = 2xp_x - 2p + F$$

and

$$\log(1-x^2+1-(x-1)^2+[1-(x-1)^2](1-x^2)p_{xx}) = 2xp_x - 2p + G$$

where F and G are constants. Equating these two expressions implies that

$$p = \frac{(2H-2)x^2+(2H-1)x+(1-H)}{x(x^2-1)((H-1)x+(2H+2))}$$

for some constant H. However, to define a smooth function on $\mathbb{C}P^2\sharp 3\overline{\mathbb{C}P^2}$ all of the second derivatives must be smooth on the polytope. This cannot be the case for any value of H. Therefore, the Siu metric does not have potential the form $\Phi = \Phi_0 + p(x) + q(y)$.

In a sense, it is not surprising that the function f is not a function of one variable or a sum of functions in x and y. Potentials of this form do not respect the 'hexagonal' symmetry of the polytope defining the symplectic structure on $\mathbb{C}P^2\sharp 3\overline{\mathbb{C}P^2}$. One could suppose the potential Φ corresponding to the Siu metric would respect that six-fold symmetry. One could propose that the function f which will give rise to the Siu metric will be of the form

$$f = \frac{1}{2}\sum_{j=1}^{m}[(A_i+B_ix)\log(A_i+B_ix)+(A_i+B_iy)\log(A_i+B_iy)+(A_i-B_i(x+y))\log(A_i-B_i(x+y))$$

$$+(A_i-B_ix)\log(A_i-B_ix)+(A_i-B_iy)\log(A_i-B_iy)+(A_i+B_i(x+y))\log(A_i+B_i(x+y))]$$

where A_i and B_i are constants on the function f is smooth on the polytope. Unfortunately, testing such an ansatz directly proves impossible using MAPLE as the calculations are too lengthy. We were able to show that the Siu metric potential does not have this form for $m = 1$.

Below, we write down explicitly a six-dimensional Kahler-Einstein metric in Kahler toric coordinates. We note that the potential of this metric can be written in the form given in the above ansatz.

6.1.4 A Kahler-Einstein Toric Metric in Dimension Six

In [54], Sakane proved the existence of a Kahler-Einstein metric on $\mathbb{P}(\mathcal{O}_{\mathbb{C}P^1\times\mathbb{C}P^1}\oplus\mathcal{O}_{\mathbb{C}P^1\times\mathbb{C}P^1}(1,-1))$ though he did not write the metric down explicitly. This is a six-dimensional Kahler toric manifold. In fact, it is a cohomogeneity-one manifold. Sakane's proof of the existence of this metric makes use

of the full symmetry group. (His construction will be discussed below in the section on fiberwise Kahler toric manifolds.) Here, we will express the Kahler-Einstein metric on this manifold in the 'symplectic' coordinates of Kahler toric geometry. For ease of expression, we set $x_1 = x$, $x_2 = y$, and $x_3 = z$. Following the theorem of Abreu, the symplectic potential, Φ, of the Kahler-Einstein metric can be written as

$$\Phi = \Phi_0 + f$$

where

$$\Phi_0 = \frac{1}{2}[(1+x)\log(1+x) + (1-x)\log(1-x) + (1+y)\log(1+y) + (1+z)\log(1+z)$$
$$+(1-x-y)\log(1-x-y) + (1+x-z)\log(1+x-z)]$$

and f and its derivatives are smooth on the polytope whose canonical potential is Φ_0. Because the metric is of cohomogeneity-one, $f = f(x)$. Substituting this expression into equation (6.3), the Einstein condition becomes

$$\log(f_{xx}(4-x^2)(1-x^2) + 3(2-x^2)) - \log(4) = 2xf_x - 2f - E.$$

Differentiating both sides with respect to x we have

$$(x^2-4)(x^2-1)f_{xxx} - 2x(x^2-4)(x^2-1)f_{xx}^2 + 2x(5x^2-11)f_{xx} - 6x = 0.$$

Through trial and error inputing rational functions into MAPLE for f_{xx}, we find that

$$f_{xx} = \frac{(x^2-10)}{(x^2-4)(x^2-7)}.$$

Integrating this twice, we can write f in the form

$$f = \frac{1}{2}[(-2+x)\log(-2+x) + (-2-x)\log(-2-x) + (1+\frac{\sqrt{7}}{7}x)\log(1+\frac{\sqrt{7}}{7}x) + (1-\frac{\sqrt{7}}{7}x)\log(1-\frac{\sqrt{7}}{7}x)].$$

Note that Φ is equivalent to the canonical potential obtained by adding four lines outside of the polytope Δ_0.

6.2 Extremal Kahler Metrics on Kahler Toric Manifolds

As proved above, a Kahler toric metric is extremal if and only if

$$-\frac{\partial^2 h_{ij}}{\partial x^i \partial x^j} = -\frac{1}{\det(h)} h_{ij} \frac{\partial^2 \det(h)}{\partial x^i \partial x^j} = \sum_{k=1}^{n} \alpha_k x_k + \beta \tag{6.6}$$

where $h^{ij} = \frac{\partial^2 \Phi}{\partial x^i \partial x^j}$. Clearly, any metric which is of constant scalar curvature is extremal. In this section we want to study the existence of extremal Kahler metrics on Kahler toric manifolds. After reviewing the construction of a one-parameter family of extremal Kahler metrics on $\mathbb{C}P^2 \sharp \overline{\mathbb{C}P^2}$ we will construct a two-parameter (up to scale) family of extremal Kahler metrics on a six-dimensional Kahler toric manifold: $\mathbb{P}(\mathcal{O}_{\mathbb{C}P^1 \times \mathbb{C}P^1} \oplus \mathcal{O}_{\mathbb{C}P^1 \times \mathbb{C}P^1}(1,-1))$. We shall see that within this two-parameter family of extremal Kahler metrics there exists a one-parameter subfamily of constant scalar curvature metrics, one of which is the Einstein metric described above.

6.2.1 Extremal Kahler Metrics on $\mathbb{C}P^2\sharp\overline{\mathbb{C}P^2}$

Extremal Kahler metrics on the blow-up of $\mathbb{C}P^2$ were originally found by Calabi in [10] but were not written down explicitly. These metrics were written in Kahler toric coordinates by Abreu in [2]. Abreu wrote the metrics explicitly by converting Calabi's non-explicit solutions into Kahler toric coordinates. Our presentation will provide the same results as Abreu; however, we will arrive at the solutions without reference to Calabi's method. Rather, we will obtain the solutions by solving equation (6.6) directly. This will be our method for obtaining new extremal Kahler metrics below.

Up to scale, we can write any canonical Kahler toric potential on $\mathbb{C}P^2\sharp\overline{\mathbb{C}P^2}$ as

$$\Phi_0 = \frac{1}{2}[(1+x)\log(1+x) + (1+y)\log(1+y) + (a-x-y)\log(a-x-y) + (1-x)\log(1-x)]$$

where $a > 0$ and we have set $x_1 = x$ and $x_2 = y$ [1]. Varying a is equivalent to changing the Kahler class of the metric. Any Kahler toric metric on $\mathbb{C}P^2\sharp\overline{\mathbb{C}P^2}$ can be written as $\Phi = \Phi_0 + f$ where f satisfies theorem (5.2). This manifold has more than the T^2-symmetry evident in these coordinates. There is in fact a $U(2)$-symmetry which implies that the manifold is actually of cohomogeneity-one. This translates into the conditions $f = f(x)$ and $S = \alpha x + \beta$ in the extremal Kahler case [1].

A straightforward calculation shows that

$$h^{-1} = \begin{pmatrix} \Phi_{xx} & \Phi_{xy} \\ \Phi_{xy} & \Phi_{yy} \end{pmatrix} = \begin{pmatrix} \frac{1}{1-x^2} + \frac{1}{2}\frac{1}{a-x-y} + f_{xx} & \frac{1}{2}\frac{1}{a-x-y} \\ \frac{1}{2}\frac{1}{a-x-y} & \frac{1}{2}\frac{1}{1+y} + \frac{1}{2}\frac{1}{a-x-y} \end{pmatrix}.$$

Setting $P(x) = f_{xx}(x) + \frac{1}{1-x^2}$, and calculating the inverse of the above matrix gives

$$h = \begin{pmatrix} \frac{2(a+1-x)}{2P(a+1-x)+1} & \frac{-2(1+y)}{2P(a+1-x)+1} \\ \frac{-2(1+y)}{2P(a+1-x)+1} & \frac{2(1+y)(2P(a-x-y)+1)}{2P(a+1-x)+1} \end{pmatrix}.$$

To simplify this further, we set $Q = P + \frac{1}{2}\frac{1}{a+1-x}$. After making this substitution, equation (6.6) becomes

$$-\frac{\partial^2(\frac{1}{Q})}{\partial x^2} + \frac{2}{a+1-x}\frac{\partial(\frac{1}{Q})}{\partial x} + \frac{4}{a+1-x} - \alpha x - \beta = 0. \tag{6.7}$$

We are able to solve this equation by setting

$$\frac{1}{Q} = \frac{1}{(a+1)-x}\int\left[\int\left(\frac{4}{a+1-x} - \alpha x - \beta\right)((a+1)-x)dx\right]dx.$$

A brief calculation shows that this indeed gives a solution for (6.7). Solving for f_{xx} we find that

$$f_{xx} = \frac{1}{2}\frac{1}{x-a-1} + \frac{a}{ax^2 - (3a^2+6a+2)x + 3a^3 + 9a^2 + 7a + 2}.$$

This metric has scalar curvature

$$S = \frac{6}{3a^2+6a+2}(-2ax + a^2 + 4a + 2).$$

For $a > 0$ we see that the metric is never of constant scalar curvature. It is intriguing to note that when $a = 0$ we retrieve the Fubini-Study metric on $\mathbb{C}P^2$. In a sense, the extremal Kahler metrics on $\mathbb{C}P^2\sharp\overline{\mathbb{C}P^2}$ and the Fubini-Study metric on $\mathbb{C}P^2$ are part of a single one parameter family of metrics.

Before proceeding to our construction of a new family of extremal Kahler metrics, we want to determine the form of the potential f which involves performing two more integrations. When writing down the canonical Kahler metric for a given polytope the potentials Φ_0 can be written schematically as

$$\Phi_0 = \sum_{k=1}^{l}[(a_k + \sum_{i=1}^{d} b_{ik}x_i)\log(a_k + \sum_{i=1}^{d} b_{ik}x_i)] \tag{6.8}$$

where l is the number of $(d-1)$-dimensional faces of the polytope and $2d$ is the real dimension of the Kahler toric manifold. While it is possible to construct Kahler toric metrics whose symplectic potentials cannot be written in this form one could ask whether all extremal Kahler metrics might have potentials of this form. Let us test this conjecture on the one parameter family of extremal Kahler metrics we have just constructed.

By inspection, we see that the potential can be written in the form of (6.8) if and only if the polynomial $ax^2 - (3a^2 + 6a + 2)x + 3a^3 + 9a^2 + 7a + 2$ has two real roots. This quadratic equation has two real roots when the discriminant is positive. That is $-3a^4 + 20a^2 + 16a + 4 > 0$. This is true for small values of $a > 0$. However, for a large enough the discriminant of this equation is negative. Therefore, not all extremal Kahler metrics on Kahler toric manifolds have symplectic potentials of the form (6.8).

While not every extremal Kahler metric on $\mathbb{C}P^2 \sharp \overline{\mathbb{C}P^2}$ has symplectic potential of the form (6.8), there is one privileged metric which does. When $a = 1$, the resulting extremal Kahler metric, g, is conformal to an Hermitian-Einstein metric, h, found by Page. That is,

$$h = S^{-2}g$$

where S is the scalar curvature of the extremal Kahler metric g.

When $a = 1$, we have

$$f_{xx} = \frac{1}{2}\frac{1}{x-2} + \frac{1}{x^2 - 11x + 21}.$$

This form of the metric was first found by Abreu in [1]. However, he did not write down the function f which determines the alteration in the canonical potential necessary to obtain this metric. Performing the simple double integral (which of course does not determine f uniquely) we can write f as

$$f = \frac{1}{2}(-2+x)\log(-2+x) + \frac{1}{2}\left(-1 - \frac{11}{\sqrt{37}} + \frac{2}{\sqrt{37}}x\right)\log\left(-1 - \frac{11}{\sqrt{37}} + \frac{2}{\sqrt{37}}x\right)$$
$$+\frac{1}{2}\left(-1 + \frac{11}{\sqrt{37}} - \frac{2}{\sqrt{37}}x\right)\log\left(-1 + \frac{11}{\sqrt{37}} - \frac{2}{\sqrt{37}}x\right).$$

We make a point of writing the symplectic potential down in this from because it shows that this privileged metric has potential which can be written in the form (6.8). We believe this to be interesting for the following reasoning. If one wants to describe a particular symplectic toric four manifold, one can exhibit a collection of lines in the plane which define a particular polytope. This

manifold, if Kahler toric, can be endowed with a canonical Kahler toric metric by adding together functions each of which is determined by one of the lines defining the polytope. Abreu, in theorem (5.2), then gives a method of modifying this potential to obtain all other possible Kahler toric metrics with the symplectic structure defined by the polytope. This is precisely the method we used to determine the extremal Kahler metric on $\mathbb{C}P^2\sharp\overline{\mathbb{C}P^2}$.

We could change the Kahler structure by an inverted method. We could begin with a collection of lines in the plane that define a polytope. We could first add lines which do not intersect the polytope at any point and then write down the canonical Kahler toric metric determined by the lines defining the polytope and by the 'extra' lines lying outside of the polytope. For example, to define a symplectic structure on the blow-up of $\mathbb{C}P^2$ at one-point we draw the following four lines in the xy-plane:

$$l_1 = 1 + x = 0, \quad l_2 = 1 - x = 0, \quad l_3 = 1 + y = 0, \quad l_4 = 1 - x - y = 0.$$

The canonical Kahler metric is then defined by the potential $\Phi_0 = \frac{1}{2}\sum_{j=1}^{4} l_j \log l_j$. To obtain the extremal Kahler metric conformal to the Page metric one could, instead of directly modifying the potential, first add three lines to those defining the polytope. That is, set

$$l_5 = -2 + x = 0, \quad l_6 = -1 - \frac{11}{\sqrt{37}} + \frac{2}{\sqrt{37}}x = 0, \quad l_7 = -1 + \frac{11}{\sqrt{37}} - \frac{2}{\sqrt{37}}x = 0.$$

The extremal Kahler metric conformal to the Page metric can then be viewed as the 'canonical' Kahler metric defined by the *seven* lines l_1 to l_7: $\Phi = \frac{1}{2}\sum_{j=1}^{7} l_j \log l_j$. A diagram of these lines appears in figure 2 below. Note that P corresponds to the original polytope on which the metric is defined.

Figure 2: *Lines defining the extremal Kahler metric conformal to the Page metric on* $\mathbb{C}P^2\sharp\overline{\mathbb{C}P^2}$



As demonstrated above, not all of the extremal Kahler manifolds can be constructed in this way. It would be interesting to know whether there is any significance to the metric which can be

described by a collection of lines (or hypersurfaces in higher dimensions.) One could ask whether all constant scalar curvature metrics or all Kahler-Einstein metrics on Kahler toric manifolds have potentials which can be written in this form. Finally, one could ask whether there is a deeper combinatorial significance to the location of the lines added to give this special metric. Unfortunately, we are unable at this time to provide an answer to any of these questions. Below, when discussing Hermitian-Einstein metrics in dimension four, we will revisit the Page metric.

6.2.2 New Extremal Kahler Metrics on $\mathbb{P}(\mathcal{O}_{\mathbb{C}P^1\times\mathbb{C}P^1}\oplus\mathcal{O}_{\mathbb{C}P^1\times\mathbb{C}P^1}(1,-1))$

The complex manifold $\mathbb{P}(\mathcal{O}_{\mathbb{C}P^1\times\mathbb{C}P^1}\oplus\mathcal{O}_{\mathbb{C}P^1\times\mathbb{C}P^1}(1,-1))$ is a manifold of six real dimensions which is invariant under the action of T^3. In fact, like the blow-up of $\mathbb{C}P^2$ at one point, this manifold is of cohomogeneity-one. In the previous section, we have written down the Kahler-Einstein metric first discovered by Sakane on this manifold in Kahler toric coordinates. In this section, we will find a two-parameter family of extremal Kahler metrics. Within this two-parameter family lies two one-parameter families of constant scalar curvature metrics and, of course, the Kahler-Einstein metric we identified earlier.

For ease of presentation, let $x = x_1, y = x_2$, and $z = x_3$. The potential of any extremal Kahler metric on this manifold can be written as $\Phi = \Phi_0 + f$ where

$$\Phi_0 = \frac{1}{2}[(1+x)\log(1+x) + (1-x)\log(1-x) + (1+y)\log(1+y) + (1+z)\log(1+z)$$

$$+(a-x-y)\log(a-x-y) + (c+x-z)\log(c+x-z)]$$

where $a > 0$ and $c > 0$ and the second derivatives of f are smooth on the polytope. A diagram of the polytope is given in figure 3. Note that the labels a and c give the lengths of the corresponding sides.

Figure 3: *Polytope associated to* Φ_0 *on* $\mathbb{P}(\mathcal{O}_{\mathbb{C}P^1\times\mathbb{C}P^1}\oplus\mathcal{O}_{\mathbb{C}P^1\times\mathbb{C}P^1}(1,-1))$



Because this manifold is of cohomogeneity-one, we can search for extremal Kahler metrics which are of cohomogeneity-one. That is, we can set $f = f(x)$. The presence of these extra symmetries tells us that, if the metric is extremal Kahler, then $S = \alpha x + \beta$, i.e. it is dependant on x only.

Setting $P(x) = f_{xx} + \frac{1}{1-x^2}$, a straightforward calculation shows that

$$h^{-1} = \begin{pmatrix} \Phi_{xx} & \Phi_{xy} & \Phi_{xz} \\ \Phi_{xy} & \Phi_{yy} & \Phi_{yz} \\ \Phi_{xz} & \Phi_{yz} & \Phi_{zz} \end{pmatrix} = \begin{pmatrix} P + \frac{1}{2}\frac{1}{a-x-y} + \frac{1}{2}\frac{1}{c+x-z} & \frac{1}{2}\frac{1}{a-x-y} & \frac{1}{2}\frac{1}{c+x-z} \\ \frac{1}{2}\frac{1}{a-x-y} & \frac{1}{2}\frac{1}{1+y} + \frac{1}{2}\frac{1}{a-x-y} & 0 \\ \frac{1}{2}\frac{1}{c+x-z} & 0 & \frac{1}{2}\frac{1}{1+z} + \frac{1}{2}\frac{1}{c+x-z} \end{pmatrix}.$$

As we did when constructing the extremal Kahler metrics on $\mathbb{C}P^2 \sharp \overline{\mathbb{C}P^2}$, we make one final change of variable by setting $Q = P + \frac{1}{2}\frac{1}{x+c+1} + \frac{1}{2}\frac{1}{a+1-x}$. Given this substitution a straightforward but cumbersome calculation demonstrates that equation (6.6) becomes

$$-\frac{\partial^2}{\partial x^2}\left(\frac{1}{Q}\right) + \left(\frac{2}{a+1-x} + \frac{-2}{c+1+x}\right)\frac{\partial}{\partial x}\left(\frac{1}{Q}\right) + \frac{2}{(x+c+1)(a+1-x)}\left(\frac{1}{Q}\right) \tag{6.9}$$
$$+\frac{4}{a+1-x} + \frac{4}{x+c+1} - \alpha x - \beta = 0.$$

We are able to solve this equation by, for the moment, inspection. Equation (6.9) has solution

$$\frac{1}{Q} = \frac{1}{(a+1-x)(x+c+1)} \int \left[\int \left(\frac{4}{a+1-x} + \frac{4}{c+1+x} - \alpha x - \beta \right) (a+1-x)(x+c+1) dx \right] dx$$

Performing this double integration and solving for f_{xx} we find that

$$f_{xx} = \frac{1}{2}\frac{1}{x-a-1} + \frac{1}{2}\frac{-1}{x+c+1} +$$

$$((-6c-6a^2c+6ac^2+6a)x+6c+10a^3c+70ac+40a^2c+40ac^2+10c^3a+5a^2c^2+6a)\{6(a-c)(1-ac)x^3$$

$$+(10a^3c-26c-42ac-8-20ac^2-20c^2-26a-20a^2-20a^2c+10c^3a-25a^2c^2)x^2$$

$$+(98a^2c+32a^2-20c^3-32c^2+20a^3-2c-30c^3a^2-60c^3a-98ac^2+30a^3c^2+2a+60a^3c)x$$

$$+8+30a^3c^3+34a+34c+114ac+52c^2+52a^2+184a^2c+184ac^2+257a^2c^2$$

$$+20c^3+20a^3+90a^3c^2+70c^3a+90c^3a^2+70a^3c\}^{-1}$$

While the calculations are somewhat unwieldy, it is not too difficult to check that this defines a positive definite metric with the appropriate smoothness conditions for all values of $a > 0$ and $c > 0$.

Proposition 6.8. *There exists, up to scale, a two-parameter family of extremal Kahler metrics on* $\mathbb{P}(\mathcal{O}_{\mathbb{C}P^1\times\mathbb{C}P^1} \oplus \mathcal{O}_{\mathbb{C}P^1\times\mathbb{C}P^1}(1,-1))$*: one for every value of* $a > 0$ *and* $c > 0$.

We further calculate that

$$\alpha = \frac{60(a-c)(ac-1)}{(30a^2c+15a^2c^2+10a^2+16a+56ac+30ac^2+4+16c+10c^2)}$$

and

$$\beta = \frac{6(10a^2+26a+20ac^2+10c^2+20a^2c+62ac+8+26c+5a^2c^2)}{(30a^2c+15a^2c^2+10a^2+16a+56ac+30ac^2+4+16c+10c^2)}.$$

When $\alpha = 0$, the metric is of constant scalar curvature. This occurs when $a = c$ or $a = \frac{1}{c}$. This proves the following proposition.

Proposition 6.9. *There exists, up to scale, two one-parameter families of constant scalar curvature Kahler metrics on* $\mathbb{P}(\mathcal{O}_{\mathbb{C}P^1\times\mathbb{C}P^1}\oplus\mathcal{O}_{\mathbb{C}P^1\times\mathbb{C}P^1}(1,-1))$*: one for every value of* $a>0$ *and* $c>0$ *such that* $a=c$ *or* $a=\frac{1}{c}$.

These two families of constant scalar curvature Kahler metrics intersect at the point $(a,c)=(1,1)$. This point of intersection corresponds to the Kahler-Einstein metric found by Sakane and presented in the previous section. Figure 4 below illustrate how the two families of constant scalar curvature Kahler metrics sits inside the two-parameter family of extremal Kahler metrics.

Figure 4: *The space of extremal Kahler and constant scalar curvature Kahler metrics on* $\mathbb{P}(\mathcal{O}_{\mathbb{C}P^1\times\mathbb{C}P^1}\oplus\mathcal{O}_{\mathbb{C}P^1\times\mathbb{C}P^1}(1,-1))$



We now ask what happens if we allow a and c to take zero values. We see immediately that if $a=c=0$ then we recover the Fubini-Study metric on $\mathbb{C}P^3$. If $a=0$ and $c>0$ (or equivalently $c=0$ and $a>0$) , then we recover a family of extremal Kahler metrics (of non-constant scalar curvature) on another six-dimensional Kahler toric manifold which is a $\mathbb{C}P^2$-bundle over $\mathbb{C}P^1$.

6.2.3 Hermitian-Einstein Manifolds in Dimension Four

While the question of the *existence* of Kahler-Einstein metrics with positive Einstein constant is completely understood in dimension four, less is known about the existence of Hermitian-Einstein metrics. LeBrun has used a theorem of Derdzinski to determine a list of compact manifolds which may admit positive Hermitian-Einstein metrics.

Theorem 6.10. *[40] If* M *is a four-dimensional compact manifold endowed with a non-Kahler Hermitian-Einstein metric* g*, then* M *is a blow-up of* $\mathbb{C}P^2$ *at one, two, or three points in general position. Any such metric would have a* T^2*-symmetry.*

Derdzinski showed that in dimension four, extremal Kahler metrics satisfying an additional second-order equation are conformally Einstein. This result is summarized in the following theorem.

Theorem 6.11. *[17] (M, g, J) is a four dimensional extremal Kahler manifold with nonconstant scalar curvature S such that*

$$S^3 + 6S\Delta S - 12\langle \nabla S, \nabla S\rangle = \beta, \tag{6.10}$$

where β is a constant, if and only if the metric $S^{-2}g$ is Hermitian-Einstein where it is defined.

Note that the resulting Einstein metric is not Kahler but is Hermitian.

In this section, we want to apply this theorem to the Kahler toric case in dimension four. That is, we want to express equation (6.10) in symplectic coordinates. Recall that the scalar curvature of an extremal Kahler toric manifold can be written as $S = \sum_{i=1}^{n} \alpha_i x_i + \beta$ where β and the α_i are constants. We now have the following corollary to the previous theorem.

Corollary 6.12. *Let (M, g, J) be a four-dimensional extremal Kahler toric manifold with scalar curvature $S = \sum_{i=1}^{2} \alpha_i x_i + \beta$. The metric $S^{-2}g$ is locally Einstein if and only if*

$$\left(\sum_{i=1}^{2} \alpha_i x_i + \beta\right)^3 + 6\left(\sum_{i=1}^{2} \alpha_i x_i + \beta\right)\left(\frac{1}{\det(h)} h_{ij} \frac{\partial \det(h)}{\partial x^j} \alpha_i\right) - 12 h_{ij}\alpha_i\alpha_j = \gamma \tag{6.11}$$

where γ is a constant.

Proof. To prove this theorem, we need only evaluate equation (6.10) on an extremal Kahler toric manifold. To do this let us look at the terms ΔS and $\langle \nabla S, \nabla S\rangle$. From the discussion above, we see that

$$\Delta S = h_{ij}\frac{\partial^2 S}{\partial x^i \partial x^j} + \frac{1}{\det(h)} h_{ij} \frac{\partial \det(h)}{\partial x^i}\frac{\partial S}{\partial x^j} = \frac{1}{\det(h)} h_{ij}\frac{\partial \det(h)}{\partial x^i}\alpha_j$$

in the extremal Kahler case. Furthermore,

$$\langle \nabla S, \nabla S\rangle = h_{ij}\frac{\partial S}{\partial x^i}\frac{\partial S}{\partial x^j} = h_{ij}\alpha_i\alpha_j.$$

The corollary follows immediately. □

This theorem and its corollary provide a strategy for finding the Hermitian-Einstein metrics (if they exist) on $\mathbb{C}P^2 \sharp 2\overline{\mathbb{C}P^2}$ and $\mathbb{C}P^2 \sharp 3\overline{\mathbb{C}P^2}$. First we could look for all (if they exist) extremal Kahler metrics on these manifolds, then check which (if any) satisfy the Derdzinski condition.

Alternatively, one could take a slightly different but ultimately equivalent approach. Namely, one could try to solve the four-dimensional Hermitian-Einstein condition directly. Knowing that any such manifold would be conformal (with a known conformal factor) to an extremal Kahler toric metric, we can write the Hermitian-Einstein equation in terms of the Kahler toric coordinates of the extremal Kahler metric to which it is conformal.

Let $\bar{g}$ be a four-dimensional Hermitian-Einstein toric metric on a compact manifold. The above shows that such a metric is conformal to an extremal Kahler toric metric g, such that

$$\bar{g} = S^{-2} g.$$

Now, because g is Kahler toric we can write it in symplectic coordinates. Recalling that in the extremal case one can express the scalar curvature in symplectic coordinates as $S = \alpha_1 x_1 + \alpha_2 x_2 + \beta$, the Hermitian-Einstein metric can be written as

$$\bar{g} = \begin{pmatrix} \frac{1}{(\alpha_1 x_1 + \alpha_2 x_2 + \beta)^2} h^{-1} & \\ & \frac{1}{(\alpha_1 x_1 + \alpha_2 x_2 + \beta)^2} h \end{pmatrix}$$

relative to coordinates $(x_1, x_2, \phi_1, \phi_2)$ where $h^{-1} = \frac{\partial^2 \Phi}{\partial x^i \partial x^j}$. We want to calculate the Einstein condition on $\bar{g}$ in terms of the h and the constants α_i and β. To do so we need to calculate the Ricci tensor $\bar{r}$. Once that is calculated the Einstein condition is, as usual, $\bar{r} = \lambda \bar{g}$. In [6], there is a formula for the change in Ricci tensor under a conformal transformation. Using that formula we have

$$\bar{r} = r + 2\frac{\mathrm{Dd}S}{S} + \left(\frac{\triangle S}{S} - 3\frac{\langle \nabla S, \nabla S \rangle}{S^2} \right) g$$

where r is the Ricci tensor, Dd is the Hessian, and $\triangle$ is the Laplacian of g. Because both g and $\bar{g}$ are complex, they and their Ricci tensor are by construction invariant under the action of the complex structure J. Therefore, to calculate the Ricci tensor $\bar{r}$ we need only calculate the values of $\bar{r}(\partial_{\phi_i}, \partial_{\phi_j})$.

Two of terms, $\frac{\triangle S}{S}$ and $\frac{\langle \nabla S, \nabla S \rangle}{S^2}$, we have already calculated in our proof of the above corollary. We need to calculate the terms $\mathrm{Dd}S(\partial_{\phi_i}, \partial_{\phi_j})$. By definition,

$$\mathrm{Dd}S(\partial_{\phi_i}, \partial_{\phi_j}) = \frac{\partial^2 S}{\partial \phi^i \partial \phi^j} - \left(\nabla_{\frac{\partial}{\partial \phi^i}} \frac{\partial}{\partial \phi^j} \right) S.$$

The first term is automatically because S does not depend explicitly on the coordinates ϕ_i. By calculating the Christoffel symbols directly, we have that

$$-\left(\nabla_{\frac{\partial}{\partial \phi^i}} \frac{\partial}{\partial \phi^j} \right) S = \frac{1}{2} h_{ik} \frac{\partial h_{lj}}{\partial x^k} \alpha_l$$

where we have here used the summation convention for repeated indices. Finally, we recall that

$$r(\partial_{\phi_i}, \partial_{\phi_j}) = -\frac{1}{2} h_{ik} \frac{\partial}{\partial x^k} \left(h_{lj} \frac{\partial \log(\det(h))}{\partial x^l} \right).$$

The above calculations prove the following.

Proposition 6.13. *The Hermitian metric $\bar{g} = S^{-2} g$ defined above is Einstein if and only if*

$$-\frac{1}{2} h_{ik} \frac{\partial}{\partial x^k} \left(h_{lj} \frac{\partial \log(\det(h))}{\partial x^l} \right) + \frac{1}{S} h_{ik} \frac{\partial h_{lj}}{\partial x^k} \alpha_l \tag{6.12}$$

$$+ \left(\frac{1}{S} h_{kl} \frac{\partial \log(\det(h))}{\partial x^k} \alpha_l - \frac{3}{S^2} h_{kl} \alpha_k \alpha_l \right) h_{ij} = \frac{\lambda}{S^2} h_{ij}$$

for $1 \leq i, j \leq 2$ where $S = \alpha_1 x_1 + \alpha_2 x_2 + \beta$.

Note that when all of the $\alpha_i = 0$ then equation (6.12) is equivalent to the Kahler-Einstein condition on Kahler toric four manifolds. There is only one known solution to equation (6.12) when S is not a constant, i.e. when the metric $\bar{g}$ is non-Kahler Hermitian-Einstein. That is the Page metric on $\mathbb{C}P^2 \sharp \overline{\mathbb{C}P^2}$. Above, we have already written down the extremal Kahler metric conformal to the Page metric. Note that in that case

$$S = -\frac{12}{11}x + \frac{42}{11} = -\frac{12}{11}x_1 + \frac{42}{11}.$$

Although the calculations are tedious and lengthy, this metric is a solution of equation (6.12). The only other possible compact solutions to equation (6.12) would be on the blow-up of $\mathbb{C}P^2$ at two or three points.

The Kahler-Einstein condition can be integrated to the Monge-Ampere equation as described above. It would be interesting to determine whether equation (6.12) can be integrated to give a modified version of that equation. Unfortunately, we have been unable to perform such an integration.

Chapter 7

Fiberwise Kahler Toric Manifolds

Kahler toric geometry enjoys many special features which are absent in Kahler geometry in general. In particular, the presence of a Legendre transformation from complex to symplectic coordinates allows one to fix a symplectic structure and vary the compatible complex structure via the potential Φ. This is not always possible in Kahler geometry. Furthermore, one can explicitly write down a metric in each Kahler class in terms of rational functions in the symplectic coordinates. In this section, we develop a generalization of Kahler toric geometry, which we refer to as fiberwise Kahler toric geometry, which shares many of these features.

We construct fiberwise Kahler toric manifolds in two very different ways. Construction one begins by considering a Kahler manifold of cohomogeneity-d under a semisimple group action. We impose two conditions: we assume that each principal orbit is a T^d-bundle over a coadjoint orbit G/L and we demand that the isotropy representation of $\mathfrak{g}$ is composed of *distinct* $\mathrm{Ad}(L)$-invariant summands (this is the monotypic case). We will see that the metric of such a manifold can be viewed as a Kahler toric metric along with a collection of linear functions in the symplectic coordinates of the Kahler toric metric.

Alternatively, construction two begins by first considering a d-parameter family of principal T^d-bundles over a product of m Kahler-Einstein manifolds, (M_i, g_i^*). Let $(u_1, .., u_d)$ denote the d parameters. We then construct a metric

$$g = h_{ij}(u)du_i \otimes du_j + h_{ij}(u)\theta_i \otimes \theta_j + \sum_{k=1}^{m} A_k(u)\pi^* g_k^*$$

where the θ_i are the connection forms of the T^d-bundle, the $\pi^* g_k^*$ are the lifts of the Kahler-Einstein metrics on the M_k, and the A_k are nonnegative functions. By defining a complex structure which satisfies $J\frac{\partial}{\partial u_i} = \frac{\partial}{\partial \phi_i}$ where $\frac{\partial}{\partial \phi_i}$ is the vector field dual to θ_i, we construct a Kahler metric which, in form, is identical to the one obtained by the previous construction.

Therefore, given completely different geometric assumptions, we construct metrics with identical properties. That is, while the manifolds constructed may be different, their metrics and curvature equations are of the same form. We will refer to both kinds as fiberwise Kahler toric manifolds.

This chapter will be devoted to making the above discussion more precise and to analyzing the Kahler-Einstein and extremal Kahler conditions on these manifolds.

7.1 Fiberwise Kahler Toric Geometry

We now present our two constructions of fiberwise Kahler toric manifolds.

7.1.1 Construction One

Take M to be a Kahler manifold of cohomogeneity-d under the action of a compact connected semisimple Lie group G where

$$G \to (M^n, g, J) \to (M/G, \check{g}). \tag{7.1}$$

Note that $(M/G, \check{g})$ is not a smooth manifold in general. However, on an open dense set in M the submersion onto the open interior of M/G is smooth. Furthermore, we assume that the metric is irreducible and is not hyperkahler so that the action of G preserves J and ω.

Because G is semisimple, there exists a G-equivariant moment map, μ, for the action of G on M. That is

$$\mu : M \to \mathfrak{g}^*$$

such that

$$d\mu(v)(X) = \omega(X, v) = g(JX, v)$$

where $v \in TM$ and $X \in \mathfrak{g}$. Note that we have suppressed the distinction between the element of $\mathfrak{g}$ and the Killing field on M induced by that element. We have the following lemma:

Lemma 7.1. *Each principal orbit G/K is a fiber bundle over a coadjoint orbit G/L*

$$L/K \to G/K \to G/L$$

such that $\dim(L/K) = d$.

Proof. The fiber bundle above is induced by the moment map, μ, which, when restricted to a principal orbit G/K, maps

$$\mu : G/K \to G/L$$

where G/L is a coadjoint orbit in $\mathfrak{g}^*$. We assumed that G was connected and semisimple, this implies that G/L is simply-connected. Therefore, L and L/K are connected.

For v a vector tangent to the fiber L/K, $d\mu(v) = 0$. This is true if and only if

$$d\mu(v)(X) = g(JX, v) = -g(X, Jv) = 0$$

for all $X \in \mathfrak{g}$.

Therefore, Jv is orthogonal to the principal orbits; Jv belongs to the horizontal distribution. The vector Jv is orthogonal to the principal orbits if and only if $d\mu(v) = 0$. As J is one-to-one we deduce immediately that $\dim(L/K) = \dim(M/G) = d$. □

We have seen how the complex structure, J, sends horizontal vector fields to vertical vector fields tangent to the fiber L/K. In the cohomogeneity-one case, $\dim(L/K) = 1$ and $L/K \cong S^1$. By contrast, L/K in the general cohomogeneity-d case need not be abelian. It is natural to first consider the case in which L/K is abelian.

ASSUMPTION ONE:

$$L/K \cong T^d.$$

As described above, the Lie algebra, $\mathfrak{g}$ can be decomposed as $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$. Given assumption one, we have that

$$\mathfrak{p} = \mathfrak{t}^d \oplus \mathfrak{p}_1 \oplus ... \oplus \mathfrak{p}_m$$

where $\mathfrak{t}^d$ is the space on which $Ad(K)$ acts trivially and the $\mathfrak{p}_\mathfrak{i}$ are non-trivial irreducible K-modules. Let d_i denote the real dimension of the summand $\mathfrak{p}_i$. Note that $\mathfrak{k} \oplus \mathfrak{t}^d$ is the Lie algebra of the group L. We make one more assumption.

ASSUMPTION TWO:

The $\mathfrak{p}_i$ are distinct.

On a Kahler toric manifold, the horizontal distribution of the quotient by group action

$$T^n \to (M^{2n}, g) \to (M/G, \check{h})$$

is integrable. Is this the case when G is semisimple and Assumption One holds?

Theorem 7.2. *The O'Neill invariant A of the submersion (7.1) is zero when assumption one holds. Equivalently, the action of G on M is **orthogonally transitive**.*

Proof. Take $\{\partial_{\phi_i}\}_{i=1,..,d}$ to be a basis for the Lie algebra of $T^d \cong L/K$. Clearly, $[\partial_{\phi_i}, \partial_{\phi_j}] = 0$ for all i and j. We demonstrated in the proof of the last lemma that $\{J\partial_{\phi_i}\}_{i=1,..,d}$ will form a basis for the horizontal distribution. ∂_{ϕ_i} is a Killing vector field and an automorphism of the complex structure J; therefore, Besse tells us that

$$\mathcal{L}_{\partial_{\phi_i}} J = 0 \Leftrightarrow [\partial_{\phi_i}, JX] = J[\partial_{\phi_i}, X]$$

for all $X \in TM$. From this equation we deduce immediately that

$$[\partial_{\phi_i}, J\partial_{\phi_j}] = [\partial_{\phi_i}, \partial_{\phi_j}] = 0$$

for all i and j. As J is an integrable complex structure, its Nijenhuis tensor, N_J vanishes. This implies that

$$N_J(\partial_{\phi_i}, \partial_{\phi_j}) = [J\partial_{\phi_i}, J\partial_{\phi_j}] - J[J\partial_{\phi_i}, \partial_{\phi_j}] - J[\partial_{\phi_i}, J\partial_{\phi_j}] - [\partial_{\phi_i}, \partial_{\phi_j}] = 0.$$

It follows immediately that

$$[J\partial_{\phi_i}, J\partial_{\phi_j}] = 0$$

for all i and j.

Therefore, $\{J\partial_{\phi_i}\}_{i=1,..,d}$ forms a commuting basis for the horizontal distribution. The existence of such a basis implies that the horizontal distribution is integrable. □

The above proof demonstrates that $\{-J\partial_{\phi_i}\}_{i=1,..,d}$ forms a coordinate basis for the horizontal distribution and for the quotient space M/G. Set $\partial_{u_i} = -J\partial_{\phi_i}$ for all i.

Dancer and Wang in [15] analyzed the structure of Kahler manifolds of cohomogeneity-one under the action of a semisimple Lie group such that assumption two holds, the $d = 1$ case. Our results in the rest of this section are a generalization of their work to the cohomogeneity-d case.

So far we have made no mention of the Kahler structure of the metric. Before we do, we can deduce the general form of the metric. Let Q be a bi-invariant metric on $\mathfrak{g}$ such that $Q(\mathfrak{k}, \mathfrak{p}) = 0$. Furthermore, we have that $Q(\mathfrak{p}_i, \mathfrak{p}_j)$ for all i and j. Choose an orthonormal basis $\{Y_j^i\}_{j=1,...,d_i}$ for $\mathfrak{p}_i$ relative to the background metric Q. For any metric g on M we have, by Assumption Two and Schur's Lemma, that

$$g(\cdot,\cdot)|_{\mathfrak{p}_1 \oplus ... \oplus \mathfrak{p}_m} = \overset{\perp}{\bigoplus_i} A_i Q(\cdot,\cdot)|_{\mathfrak{p}_i}$$

where the A_i are positive functions on the interior of the quotient space M/G. That is $g(Y_j^i, Y_j^i) = A_i$ for all i, j, and k. Furthermore, set $h_{ij} = g(\partial_{\phi_i}, \partial_{\phi_j})$. Because of the invariance of the metric under the complex structure J, we deduce immediately that, $h_{ij} = g(\partial_{u_i}, \partial_{u_j})$. Therefore, the metric, g, on the interior of the quotient space can be written as

$$g = \begin{pmatrix} h_{ij} & & & & \\ & h_{ij} & & & \\ & & A_1 & & \\ & & & ... & \\ & & & & A_m \end{pmatrix}$$

relative to the basis $(\partial_{u_1}, ..., \partial_{u_d}, \partial_{\phi_1}, ..., \partial_{\phi_d}, Y_1^1, ..., Y_{d_m}^m)$. The metric is determined by $\frac{d(d+1)}{2} + m$ functions of the coordinates $(u_1, ..., u_d)$. In addition to being pairwise inequivalent K-modules, the $\mathfrak{p}_\mathfrak{i}$ are also pairwise inequivalent L-modules. For each p in the interior of M/G, let g_p be the restriction

of the metric to the principal orbit that is the inverse image of the point p. Therefore g_p induces an $\mathrm{Ad}(L)$-invariant inner product on $\mathfrak{q} = \mathfrak{p}_1 \oplus ... \oplus \mathfrak{p}_m$ which can be identified with the tangent space to G/L. The induced G-invariant metric on G/L will be denoted by g_p^*.

The Kahler Condition:

In the Kahler case, $d\omega = 0$ where $\omega(\cdot,\cdot) = g(J\cdot,\cdot)$.

Theorem 7.3. *Let (M, g, J) be a Fiberwise Kahler Toric manifold. The Kahler condition is equivalent to*

(i) $\omega_p([X,Y]_\mathfrak{p}, Z) + \omega_p([Y,Z]_\mathfrak{p}, X) + \omega_p([Z,X]_\mathfrak{p}, Y) = 0$

(ii) $\partial_{u_i}\omega(X,Y) + \omega(\partial_{u_i}, [X,Y]_\mathfrak{p}) = 0$

where p is a point in the interior of M/G and $X, Y, Z \in \mathfrak{p}$ and

(iii) $\bar{g} = \begin{pmatrix} h_{ij} & \\ & h_{ij} \end{pmatrix}$ *is locally Kahler toric relative to basis $(\partial_{u_1}, ..., \partial u_d, \partial_{\phi_1}, ..., \partial_{\phi_d})$, i.e. $h_{kl} = \frac{\partial^2 \eta}{\partial u^k \partial u^l}$ for some function $\eta = \eta(u_1, ..., u_d)$.*

Proof. First, consider arbitrary vector fields X, Y, and Z in TM. The following equality always holds for any two-form ω

$$d\omega(X,Y,Z) = X(\omega(Z,Y)) + Y(\omega(X,Z)) + Z(\omega(Y,X)) + \omega([X,Y]_\mathfrak{p}, Z) + \omega([Y,Z]_\mathfrak{p}, X) + \omega([Z,X]_\mathfrak{p}, Y)$$

In the Kahler case, $d\omega = 0$. If we choose $X, Y, Z \in \mathfrak{p}$, the invariance of the Kahler form under the action of the group G immediately gives (i). Setting $Z = \partial_{u_i}$, we obtain (ii) by noting the $[\partial_{u_i}, X] = 0$ for all $X \in \mathfrak{p}$. Part (iii) can be obtained by noting that the metric $\bar{g}$ is well-defined locally relative to the basis given and all the vector fields commute. Therefore, locally, the complex structure and metric are that of a toric manifold and ω restricted to the basis given is precisely $\bar{\omega}$. If ω is closed then $\bar{\omega}$ is necessarily closed and the metric $\bar{g}$ is Kahler toric. The only remaining case to check is when $X = \partial_{u_i}$, $Z = \partial_{u_j}$, and $Y = Y_m^k \in \mathfrak{p}_k$. The Kahler condition is automatically satisfied since each ∂_{u_i} commutes with Y_m^k. Therefore, the equations listed in the theorem are the only ones not automatically satisfied. □

For every such manifold, there is a natural projection from (M, g) onto $(G/L, g^*)$. The fiber of this projection is a Kahler toric manifold $(N, \bar{g})$ such that

$$T^d \to (N, \bar{g}) \to (N/T^d, \breve{\bar{g}}) \cong (M/G, \breve{g}).$$

Moreover, the existence of this Kahler toric manifold will allow us to later make use of the 'symplectic' coordinates obtained by changing the coordinates of the base manifold M/G. Before we do that, however, let us first analyze in greater detail the action of the complex structure J.

Proposition 7.4. *Let (M, g, J) be a Kahler manifold of cohomogeneity-d under the action of a semisimple Lie group such that assumptions one and two hold. There is a coadjoint orbit G/L with $K \subset L$ such that*

(1) for each point p in the interior of the quotient space M/G, $g_p = g\,|_{\pi^{-1}(p)}$ induces a G-invariant metric g_p^* on G/L and the projection $\mu : (G/K, g_p) \to (G/L, g_p^*)$ is a Riemannian submersion with totally geodesic T_d fibers.

(2) there is a complex structure J^*, which does not depend on the choice of p, on G/L and relative to this complex structure g_p^* is Kahler.

Proof. Part (1) follows directly from the discussion above. To obtain (2), note that J_p is $\mathrm{Ad}(K)$-invariant and skew-symmetric with respect to the metric g_p. Therefore, we deduce that J_p leaves each $\mathfrak{p}_i$ invariant because of assumption two. The above discussion also demonstrates that J_p is independent of the choice of p. Because each summand $\mathfrak{p}_i$ is irreducible, all invariant metrics differ only by a constant. This implies that $J\,|_q$ is skew-symmetric with respect to the Killing form.

From Lemma (7.1) we see that $\mathrm{ad}(\partial_{\phi_i})$ is $\mathrm{Ad}(K)$-invariant for every $i = 1, ..., d$ and if, of course, skew with respect to the Killing form. On each $\mathfrak{p}_i$, J is proportional to $\mathrm{ad}(\partial_{\phi_i})$ and also commutes with that operator. This implies that $J\,|_q$ is in fact $\mathrm{Ad}(L)$-invariant and gives rise to an almost-complex structure J^* on G/L. To complete the proof of this proposition, we need to show that this complex structure is integrable and that the induced invariant metric on G/L is Kahler with respect to this complex structure.

Integrability:

The integrability of J^* is equivalent to the vanishing of the Nijenhuis tensor which is equivalent to

$$[J^*X, J^*Y]_{\mathfrak{q}} - [X, Y]_{\mathfrak{q}} - J^*[X, J^*Y]_{\mathfrak{q}} - J^*[J^*X, Y]_{\mathfrak{q}} = 0$$

for all $X, Y \in \mathfrak{q}$. Now, the submersion $\mu : G/K \times \{p\} \to G/L$ is independent of the point p as discussed above. Take $\check{X}$ and $\check{Y}$ to be basic vector fields of this submersion. We have that $d\mu[\check{X}, \check{Y}] = [d\mu(\check{X}), d\mu(\check{Y})]$. Also, we see that $d\mu(J\check{X}) = J^*(d\mu(\check{X}))$ because of the $\mathrm{Ad}(L)$ invariance of J. This together with the integrability of J implies the integrability of J^*.

Kahler:

We now must demonstrate that the metric (g_p^*, J^*) is Kahler. The Kahler form, ω_p^*, is closed by part (i) of (7.3). □

Instead of the construction above, we could construct manifolds of this type by the following process. Let G/L be a coadjoint orbit, whose isotropy representation is composed of inequivalent summands, with complex structure J^*. Take g_p^* to be a d-parameter family of Kahler metrics on G/L. The first Chern class of $(G/L, J^*)$ is positive and therefore all class in $H^2(G/L;\mathbb{Z})$ are of type $(1,1)$. Every T^d-bundle over G/L is of the form G/K where $L/K \cong T^d$.

Any principal T^d-bundle, G/K, over G/L ($\mu : G/K \to G/L$) is characterized by d invariant S^1 connections θ_i, $1 \leq i \leq d$. We can construct a metric

$$g = h_{ij} du_i \otimes du_j + h_{ij}\theta_i \otimes \theta_j + \mu^* g_p^*$$

locally on $G/K \times \mathbb{R}^d$ where $h_{ij} = h_{ij}(u_1, ..., u_d)$. We define the complex structure J to be the J^* when restricted to vector field tangent to G/L and to otherwise satisfy the equations $J\partial_{u_i} = \partial_{\phi_i}$ where ∂_{ϕ_i} is the dual vector field to the connection θ_i. Furthermore, we demand that $h_{ij} = \frac{\partial^2 \eta}{\partial_{u_i}\partial_{u_j}}$. It follows that this defines a Kahler metric which is invariant under the action of G.

We return now to the Kahler condition. For $X, Y \in \mathfrak{q}$, the Kahler condition implies that

$$\partial_{u_i}(\omega(X,Y)) + \omega(\partial_{u_i}, [X,Y]_{\mathfrak{p}}) = 0$$

for all $1 \leq i \leq d$. As mentioned above, the moment map induces a fibration $\mu : G/K \to G/L$ on each principal orbit. This is a principal T^d-bundle over a coadjoint orbit. Since, as proved above, each fiber is totally geodesic, the O'Neill invariant T^μ of this submersion vanishes. However, the invariant A^μ of the submersion does not vanish. From the definition of

$$A^\mu_X Y = \frac{1}{2}\mathcal{V}[X,Y]_{\mathfrak{p}}$$

for $X, Y \in \mathfrak{q}$ where $\mathcal{V}$ is the vertical projection onto the T^d fibers. Letting $\{\partial_{\phi_i}\}$ be a basis from $\mathfrak{t}^d$ letting $\omega_j^* = \omega^*_{\mathfrak{p}_j}$, we have that

$$A^\mu_X Y = -\frac{1}{2}\sum_{k=1}^{d}\sum_{j=1}^{m} b_{kj}\omega_j^*(X,Y)\partial_{\phi_k}$$

where the b_{jk} are the eigenvalues of the curvature form of the connection θ_j.

Take $X_j, Y_j \in \mathfrak{p}_j$, the Kahler condition implies that

$$\partial_{u_i}(\omega(X_i, Y_i)) + \omega(\partial_{u_i}, [X_j, Y_j]_{\mathfrak{p}_j}) = 0$$

which implies that

$$\partial_{u_i}(g(X_i, Y_i)) = g(\partial_{\phi_i}, \sum_{k=1}^{d} b_{kj}\omega_j^*(X_j, Y_j)\partial_{\phi_k}).$$

Using the form of the metric g, it is straightforward to calculate that the Kahler condition implies that

$$\partial_{u_i} A_j = \sum_{k=1}^{d} h_{ik} b_{kj}.$$

multiplying both sides by h^{-1} and making use of the summation convention, we have that

$$h^{ik}\partial_{u_i}A_j = b_{kj} \tag{7.2}$$

for all $1 \leq k \leq d$ and all $1 \leq j \leq m$. Therefore, up to constants, the Kahler condition completely determines the functions $A_j = A_j(u_1, ..., u_d)$ in terms of the h_{ij} and the b_{ij}. Although the A_j satisfy (7.2), this equation is somewhat difficult to use directly as determining an explicit formula for the A_j would involve integrating the functions h_{ij}. If, however, we change coordinates on the quotient space, we can place equation (7.2) in a form which can be explicitly integrated.

We have demonstrated that within every manifold of this type is a Riemannian submersion over coadjoint orbit G/L such that the fiber of this submersion is a Kahler toric manifold with metric

$$\bar{g} = \begin{pmatrix} h_{ij} & \\ & h_{ij} \end{pmatrix}$$

relative to vector fields $(\partial_{u_1}, ..., \partial_{\phi_d})$ such that $h_{ij} = \frac{\partial^2 \eta}{\partial_{u_i}\partial_{u_j}}$. Instead, of these complex coordinates, we could express the Kahler toric metric in symplectic coordinates by setting $x_i = \frac{\partial \eta}{\partial u_i}$. With this change, we have

$$\bar{g} = \begin{pmatrix} h^{ij} & \\ & h_{ij} \end{pmatrix}$$

relative to basis $(\partial_{x_1}, ..., \partial_{x_d}, \partial_{\phi_1}, ..., \partial_{\phi_d})$.

We can now view the functions A_j as functions of the x_i. After expressing equation (7.2) in these coordinates, we obtain the following lemma.

Lemma 7.5. *The Kahler condition implies that the functions $A_j(x_1, ..., x_d)$ are of the form*

$$A_j = \sum_{k=1}^{d} b_{kj}x_k + a_j \tag{7.3}$$

where the a_j are constants.

Proof. From calculations given in the section on Kahler toric manifolds, $h^{kl}\frac{\partial}{\partial u^l} = \frac{\partial}{\partial x_k}$. Equation (7.2) implies that

$$h^{ik}\partial_{u_i}A_j = \frac{\partial}{\partial x^k}A_j = b_{kj}$$

for all $1 \leq j \leq m$ and all $1 \leq k \leq d$. The lemma follows immediately. □

We pause here to discuss the significance of the constants b_{kj} and a_j. Each principal orbit G/K is a principal T^d-bundle over coadjoint orbit G/L. Given a particular G/L, a principal T^d-bundle over that orbit is determined by the constants b_{kj}. While changing the constants a_j alters the metric is does not change the principal orbit type.

These above results are summarized in the following theorem:

Theorem 7.6. *The metric on a Kahler manifold (M, g, J) of cohomogeneity-d under the action of a semisimple Lie group such that assumptions one and two hold can be written locally as*

$$g = \begin{pmatrix} h^{ij} & & & & \\ & h_{ij} & & & \\ & & (\sum_{k=1}^{d} b_{k1}x_k + a_1)Id_{d_1} & & \\ & & & ... & \\ & & & & (\sum_{k=1}^{d} b_{km}x_k + a_m)Id_{d_m} \end{pmatrix}$$

relative to basis $(\partial_{x_1}, ..., \partial_{x_d}, \partial_{\phi_1}, ..., \partial_{\phi_d}, Y_1^1, ..., Y_{d_m}^m)$ *and*

$$h^{kl} = \frac{\partial^2 \Phi}{\partial x^k \partial x^l}$$

for $\Phi = \Phi(x_1, ..., x_d)$.

7.1.2 Construction Two

We now construct Kahler manifolds with metrics of the same form using different geometric assumptions. Let (M_k, J_k^*, g_k^*) be compact Kahler-Einstein manifolds of real dimension d_k with positive Einstein constant for all $1 \leq k \leq m$. We normalize each of these manifolds so that the Einstein constant is equal to one. Furthermore, let P be a principal T^d-bundle over $M_1 \times ... \times M_m$. Manifolds of this type were studied by Wang and Ziller in [58] and we rely on their construction here.

A principal T^d-bundle $\pi : P \to M_1 \times ... \times M_m$ is characterized by d connections θ_i which satisfy $d\theta_i = \pi^*(\sum_{k=1}^{m} b_{ik}\omega_k^*)$ for constants b_{ij}. We can construct a metric on such a torus bundle by writing

$$h_{ij}\theta_i \otimes \theta_j + \sum_{k=1}^{m} A_k \pi^* g_k^*$$

where the A_k are, for the moment, constants and h_{ij} is a positive-definite matrix of constants. (Note that here we are summing over i and j.) It is important to note the product manifold $(M_1 \times ... \times M_m, \sum_{k=1}^{m} A_k g_k^*)$, is itself not necessarily Einstein, though it is of constant scalar curvature. The map π is a Riemannian submersion with totally geodesic fibers. Let A^π denote the O'Neill invariant of this submersion. By definition,

$$A_X^\pi Y = \frac{1}{2}\mathcal{V}[X, Y]$$

for X and Y vector fields in the horizontal distribution of the submersion π. If we let ∂_{ϕ_i} be the dual vector field to the connection θ_i, it is straightforward to calculate (see [58] for details) that

$$A_X^\pi Y = -\frac{1}{2}\sum_{j=1}^{d}\sum_{k=1}^{m} b_{jk}\omega_k^*(X, Y)\partial_{\phi_j}.$$

Next, we consider a d-parameter family of such metrics, i.e. we set $h_{ij} = h_{ij}(u_1, ..., u_d)$ and $A_k = A_k(u_1, ..., u_d)$. We now construct what is to be our fiberwise Kahler metric by setting

$$g = h_{ij}du_i \otimes du_j + h_{ij}\theta_i \otimes \theta_j + \sum_{k=1}^{m} A_k \pi^* g_k^*.$$

Locally, we now have a Riemannian manifold (M, g) of dimension $n = 2d + \sum_{k=1}^{m} d_k$. To turn this into a Kahler manifold, we first need to define a complex structure J which is compatible with this metric. If X is the lift of a vector field on the product of the M_i then set $JX = \sum_{k=1}^{m} \pi^* J_k^* X$. Also, let $J\partial_{u_i} = \partial_{\phi_i}$ for all i. These requirements completely determine a compatible almost complex structure and it is straightforward to see that this almost complex structure is in fact integrable. Thus, locally, (M, g, J) is a complex manifold. We note that the Einstein condition on metrics of this type were studied in both the Hermitian and the Kahler case for $d = 1$ by Wang and Wang in [57].

We define the two form ω in the usual way by the equation $\omega(\cdot, \cdot) = g(J\cdot, \cdot)$. The Kahler condition is equivalent to $d\omega = 0$. Using a well-known identity, this condition is equivalent to

$$X(\omega(Z,Y)) + Y(\omega(X,Z)) + Z(\omega(Y,X)) + \omega([X,Y],Z) + \omega([Y,Z],X) + \omega([Z,X],Y) = 0$$

for all X, Y, and Z in TM. Consider the induced metric

$$\bar{g} = h_{ij} du_i \otimes du_j + h_{ij} \theta_i \otimes \theta_j \tag{7.4}$$

with induced complex structure $\bar{J} = J|_{\text{span}\{\partial_{u_1}, \ldots, \partial_{\phi_d}\}}$. Let $\bar{\omega}(\cdot, \cdot) = \bar{g}(\bar{J}, \cdot, \cdot)$. From equation (7.4), it is clear that $d\omega = 0$ implies that $d\bar{\omega} = 0$. Put another way, if g is Kahler then $\bar{g}$ is Kahler toric, i.e. $h_{kl} = \frac{\partial^2 \eta}{\partial u^k \partial u^l}$ for some function $\eta = \eta(u_1, ..., u_d)$.

Next, let X, Y and Z be the lifts of vector fields on the product of the Kahler-Einstein manifolds $(M_k, g_k^*$. For such vector fields $d\omega(X, Y, Z) = 0$ because each of the ω_k^* are closed. Lastly, we see that, given X and Y as above, $d\omega$ implies that

$$\partial_{u_i} \omega(X, Y) + \omega(\partial_{u_i}, [X, Y]) = 0$$

for all $1 \leq i \leq d$. This equation implies that

$$\partial_{u_i}(g(X,Y)) = g(\partial_{\phi_i}, \sum_{j=1}^{d} b_{jk} \sum_{k=1}^{m} \omega_k^*(X,Y) \partial_{\phi_j}).$$

From the form of the metric determined above, this becomes

$$\partial_{u_i} A_k = \sum_{j=1}^{d} h_{ij} b_{jk} \tag{7.5}$$

for all $1 \leq j \leq d$ and all $1 \leq k \leq m$. Up to a constant, the Kahler condition completely determines the function $A_i(u)$. We can express this equation in a simplified form if we express the Kahler toric metric $\bar{g}$ in terms of symplectic coordinates. That is, let $x_i = \frac{\partial \eta}{\partial_{u_i}}$ for all $\leq i \leq d$. Given this coordinate change

$$\bar{g} = \begin{pmatrix} h^{ij} & \\ & h_{ij} \end{pmatrix}$$

relative to basis $(\partial_{x_1}, ..., \partial_{\phi_d})$. Given this coordinate change, equation (7.5) implies that

$$A_k = \sum_{j=1}^{d} b_{jk} x_j + a_j.$$

This completes the calculation of the Kahler condition.

As in construction one, we wish to describe more precisely the significance of the constants b_{jk} and a_j. Each point in the interior of M/G is a principal T^d-bundle over the product of Kahler-Einstein $M_1 \times ... \times M_m$. Given a product manifold, the various principal T^d-bundles are described by the constants b_{jk}. Varying the constants a_j alters the metric but does not alter these principal torus bundles. However, the a_j in part determine the zero sets of the functions A_j. Whether or not some of the A_j vanish at special orbits will modify the manifold on which the metric is defined. While the a_j do not affect the principal orbit type, they do affect the topology of the total manifold M.

The above results are summarized in the following theorem:

Theorem 7.7. *The metric on Kahler manifold (M, g, J) obtained via construction two can be written locally as*

$$g = \begin{pmatrix} h^{ij} & & & & \\ & h_{ij} & & & \\ & & (\sum_{k=1}^{d} b_{k1} x_k + a_1) Id_{d_1} & & \\ & & & ... & \\ & & & & (\sum_{k=1}^{d} b_{km} x_k + a_m) Id_{d_m} \end{pmatrix}$$

relative to basis $(\partial_{x_1}, ..., \partial_{x_d}, \partial_{\phi_1}, ..., \partial_{\phi_d}, Y_1^1, ..., Y_{d_m}^m)$ and

$$h^{kl} = \frac{\partial^2 \Phi}{\partial x^k \partial x^l}$$

for $\Phi = \Phi(x_1, ..., x_d)$ where $(Y_1^k, ..., Y_{d_k}^k)$ is an orthonormal basis for the Kahler-Einstein manifold (M_k, g_k^*).

Note that this metric has exactly the same form as the metric determined by construction one. This justifies the following definition:

Definition 9. *We refer to a Kahler manifold (M, g, J) obtained via either construction one or construction two as a **fiberwise Kahler toric manifold**.*

When calculating the curvature equations on a fiberwise Kahler toric manifold, we will not distinguish between to two types of manifolds determined by the above constructions. Since the curvature equations are identical, it is not necessary to distinguish between them for the purpose of calculations. Finally, we state the following definition:

Definition 10. *A fiberwise Kahler toric manifold is of **cohomogeneity-**d if h_{ij} is a $d \times d$ matrix.*

In the case of manifolds obtained via construction one, the appellation is strictly true as the manifolds are of cohomogeneity-d. In the case of manifolds obtained via construction two, the metrics need only have a T^d symmetry and will therefore not necessarily be of cohomogeneity-d. As the form of the metric on the two types of manifolds are the same, we believe this abuse will not cause confusion.

Before discussing various curvature conditions on fiberwise Kahler toric manifolds, we make a few comments about the coordinates we are using. The coordinates on the induced Kahler toric metric, $\bar{g}$, are the symplectic Darboux coordinates. However, the metric on the fiberwise Kahler manifold presented here is not written in Darboux coordinates at all. Furthermore, these metrics are not written in complex coordinates. We believe that it is best to think of fiberwise Kahler toric manifolds, and these coordinates, as Kahler toric manifolds with metrics written in symplectic coordinates along with additional structure. The connection with Kahler toric geometry will be central in our work below.

7.2 Curvature Equations and Conditions

In this section, we calculate the scalar curvature equation and the equations of the Einstein condition on a fiberwise Kahler toric manifold. After presenting these equations, we will examine the extremal Kahler condition on manifolds of this type.

7.2.1 The Einstein Equations

We demonstrated above that the action of the group G is orthogonally transitive in the case of a fiberwise Kahler toric manifold. The full Einstein equations on a manifold of cohomogeneity-d under an orthogonally transitive group action are given in the section on Riemannian submersions. Not surprisingly, the equations given in theorem (2.5) simply greatly in the fiberwise Kahler toric case. We begin with the following lemma.

Lemma 7.8. *If (M, g, J) is a fiberwise Kahler toric manifold, then*

$$r(\partial_{\phi_i}, \partial_{u_j}) = 0$$

for all i, j. This implies that the second equation of the Einstein condition in theorem (2.5) is automatically satisfied.

Proof. Recall the second equation of the Einstein condition given in theorem (2.5). The form of a fiberwise Kahler toric metric implies that this equation becomes

$$h^{ij}\breve{\nabla}\langle\partial_{\phi_i}, [\partial_{\phi_j}, X]\rangle = 0$$

for all $X \in \mathfrak{p}$. However, the vector fields $\partial_{\phi i}$ commute with all other vector fields in $\mathfrak{p}$. Therefore, the left-hand side of this equation is automatically equal to zero. □

Next we recall that $\partial_{u_i} = -J\partial_{\phi_i}$. Because the metric is Kahler it is invariant under the action of the complex structure J. Furthermore, on a Kahler manifold, the Ricci tensor is also invariant under the action of the complex structure. That is, $r(\partial_{u_i}, \partial_{u_j}) = r(J\partial_{\phi_i}, J\partial_{\phi_j}) = r(\partial_{\phi_i}, \partial_{\phi_j})$. The invariance of the metric and the Ricci tensor under the action of the complex structure proves the following lemma.

Lemma 7.9. *On a fiberwise Kahler toric manifold, if the first equation of the Einstein condition in (2.5) is satisfied then the third equation is satisfied automatically.*

We further note that $r(\partial_{\phi_i}, Y_i^j) = 0$ for all i and j and $r(Y_k^l, Y_i^j) = 0$ for all $k \neq i$ and $l \neq j$. Therefore, the Einstein condition on a fiberwise Kahler toric manifold contains two types of equations: $r(\partial_{\phi_i}, \partial_{\phi_j}) = \lambda g(\partial_{\phi_i}, \partial_{\phi_j})$ and $r(Y_k^i, Y_k^i) = \lambda g(Y_k^i, Y_k^i)$. To calculate these equations we will need to determine the quantities $\hat{r}(\partial_{\phi_i}, \partial_{\phi_j})$ and $\hat{r}(Y_k^i, Y_k^i)$.

Noting that each principal orbit is a principal torus bundle over a coadjoint orbit and using Wang and Ziller's work in [58] we calculate that

$$\hat{r}(Y_k^i, Y_k^i) = r^*(Y_k^i, Y_k^i) - \frac{1}{2}\frac{h_{kl}b_{ki}b_{li}}{A_i}$$

where we have summed over the repeated indices k and l. Dancer and Wang proved that $r^*(Y_k^i, Y_k^i) = 1$ in the cohomogeneity-one case. However, their proof works without alteration in the cohomogeneity-d case. Therefore,

$$\hat{r}(Y_k^i, Y_k^i) = 1 - \frac{1}{2}\frac{h_{kl}b_{ki}b_{li}}{A_i}.$$

Similarly, we calculate that

$$\hat{r}(\partial_{\phi_i}, \partial_{\phi_j}) = \frac{1}{4}h_{ik}h_{jl}\sum_{r=1}^{m}\frac{d_r b_{kr}b_{lr}}{A_r^2}$$

Using the first equation in (2.5) we calculate that

$$r(Y_k^i, Y_k^i) = -\frac{1}{2}\check{\triangle}A_i - \frac{1}{4}\frac{\langle\check{\nabla}\det(h), \check{\nabla}A_i\rangle}{\det(h)} - \frac{1}{4}\frac{\langle\check{\nabla}V, \check{\nabla}A_i\rangle}{V} - \frac{1}{2}A_i\langle\check{\nabla}(\frac{1}{A_i}), \check{\nabla}A_i\rangle + 1 - \frac{1}{2}\frac{h_{kl}b_{ki}b_{li}}{A_i}$$

where $V = \prod_{r=1}^{m} A_r^{d_r}$. Noting that, just as in the Kahler toric case, $\check{\triangle}(\cdot) = h_{kl}\frac{\partial^2(\cdot)}{\partial x^k \partial x^l} + \frac{1}{2}\frac{1}{\det(h)}h_{kl}\frac{\partial\det(h)}{\partial x^k}\frac{\partial(\cdot)}{\partial x^l}$. Since the A_j are linear in the x_i, we see that $\frac{\partial^2 A_j}{\partial x^k \partial x^l} = 0$ for all j, k, l. We calculate that

$$-\frac{1}{2}A_i\langle\check{\nabla}(\frac{1}{A_i}), \check{\nabla}A_i\rangle = \frac{1}{2}\frac{1}{A_i}h_{kl}\frac{\partial A_i}{\partial x^k}\frac{\partial A_i}{\partial x^l} = \frac{1}{2}\frac{h_{kl}b_{ki}b_{li}}{A_i}.$$

Equation $r(Y_k^i, Y_k^i) = \lambda g(Y_k^i, Y_k^i)$ of the Einstein condition becomes

$$-\frac{1}{2}\frac{\langle\check{\nabla}\det(h), \check{\nabla}A_i\rangle}{\det(h)} - \frac{1}{4}\frac{\langle\check{\nabla}V, \check{\nabla}A_i\rangle}{V} + 1 = \lambda A_i$$

for all $1 \leq i \leq m$ where $A_i = \sum_{r=1}^{d} b_{ri}x_r + a_i$. Expanding this equation gives

$$\frac{1}{\det(h)}h_{kl}\frac{\partial\det(h)}{\partial x^l}\frac{\partial A_i}{\partial x^k} + \frac{1}{2}h_{kl}\frac{\partial\log V}{\partial x^l}\frac{\partial A_i}{\partial x^k} - 2 = -2\lambda\sum_{k=1}^{d} b_{ki}x_k + a_i.$$

This becomes

$$(h_{kl}\frac{\partial \log(\det(h)V^{\frac{1}{2}})}{\partial x^l} + 2\lambda x_k)b_{ki} = 2(1-\lambda a_i)$$

for all $1 \leq i \leq m$.

Next, we need to express equation $r(\partial_{\phi_i}, \partial_{\phi_j}) = \lambda g(\partial_{\phi_i}, \partial_{\phi_j})$ in terms of the h_{kl} and the A_i. The first equation in (2.5) becomes

$$-\frac{1}{2}\check{\triangle}h_{ij} - \frac{1}{4}\frac{\langle \check{\nabla}\det(h), \check{\nabla}h_{ij}\rangle}{\det(h)} - \frac{1}{4}\frac{\langle \check{\nabla}V, \check{\nabla}h_{ij}\rangle}{V} - \frac{1}{2}h_{ik}\langle \check{\nabla}h^{kl}, \check{\nabla}h_{lj}\rangle + \frac{1}{4}h_{ik}h_{jl}\sum_{r=1}^{m}\frac{d_r b_{kr}b_{lr}}{A_r^2} = \lambda h_{ij}.$$

Using the calculations for the Einstein condition of a Kahler toric manifold performed in the previous section, we see that the above equation becomes

$$-\frac{1}{2}h_{ik}\frac{\partial}{\partial x^k}(\frac{1}{\det(h)}h_{lj}\frac{\partial \det(h)}{\partial x^l}) - \frac{1}{4}\frac{\langle \check{\nabla}V, \check{\nabla}h_{ij}\rangle}{V} + \frac{1}{4}h_{ik}h_{jl}\sum_{r=1}^{m}\frac{d_r b_{kr}b_{lr}}{A_r^2} = \lambda h_{ij}$$

for all $1 \leq i, j \leq d$.

Expanding this equation we see that

$$h_{ik}\frac{\partial}{\partial x^k}(\frac{1}{\det(h)}h_{lj}\frac{\partial \det(h)}{\partial x^l}) + h_{ik}\frac{\partial h_{lj}}{\partial x^k}\frac{\partial \log(V^{\frac{1}{2}})}{\partial x^l} + h_{ik}h_{jl}\frac{\partial^2 \log(V^{\frac{1}{2}})}{\partial x^k \partial x^l} = -2\lambda h_{ij}$$

which is equivalent to

$$h_{ik}\frac{\partial}{\partial x^k}(h_{lj}\frac{\partial \log(\det(h)V^{\frac{1}{2}})}{\partial x^l}) = -2\lambda h_{ij}.$$

We collect the above calculations in the following theorem

Theorem 7.10. *The Einstein condition on a fiberwise Kahler toric manifold is given by*

$$h_{ik}\frac{\partial}{\partial x^k}\left(h_{lj}\frac{\partial \log(\det(h)V^{\frac{1}{2}})}{\partial x^l}\right) = -2\lambda h_{ij} \tag{7.6}$$

for all $1 \leq i, j \leq d$ *and*

$$\left(h_{kl}\frac{\partial \log(\det(h)V^{\frac{1}{2}})}{\partial x^l} + 2\lambda x_k\right)b_{ki} = 2(1-\lambda a_i) \tag{7.7}$$

for all $1 \leq i \leq m$, *where* λ *is the Einstein constant,* $h^{kl} = \frac{\partial^2 \Phi}{\partial x^k \partial x^l}$, $A_j = \sum_{r=1}^{d} b_{ri}x_r + a_j$ *for all* $1 \leq j \leq m$, *and* $V = \prod_{r=1}^{m} A_r^{d_r}$.

7.2.2 The Scalar Curvature Equation

Having calculated the equations of the Einstein condition, it is a simple matter to calculate the scalar curvature equation (the scalar curvature being the trace of the Ricci tensor).

Theorem 7.11. *The scalar curvature, S, of a fiberwise Kahler toric manifold can be written as*

$$S = -\frac{1}{\det(h)}h_{kl}\frac{\partial^2 \det(h)}{\partial x^k \partial x^l} - \sum_{r=1}^{m} d_r \frac{h_{kl}\frac{\partial \det(h)}{\partial x^k}b_{lr}}{\det(h)A_r} + \frac{1}{2}\sum_{r=1}^{m} d_r \frac{h_{kl}b_{kr}b_{lr}}{A_r^2} \tag{7.8}$$

$$-\frac{1}{4}\sum_{r=1}^{m}\sum_{s=1}^{m} d_r d_s \frac{h_{kl}b_{kr}b_{ls}}{A_r A_s} + \sum_{r=1}^{m}\frac{d_r}{A_r}$$

Proof. From the equations of the Einstein condition, we calculated that

$$S = \bar{S} - \sum_{r=1}^{m} d_r \frac{\langle \check{\nabla} \det(h), \check{\nabla} A_r \rangle}{\det(h) A_r} + \frac{1}{2} \sum_{r=1}^{m} d_r \frac{\langle \check{\nabla} A_r, \check{\nabla} A_r \rangle}{A_r^2} - \frac{1}{4} \sum_{r=1}^{m} \sum_{s=1}^{m} d_r d_s \frac{\langle \check{\nabla} A_r, \check{\nabla} A_s \rangle}{A_r A_s} + \sum_{r=1}^{m} \frac{d_r}{A_r}$$

where $\bar{S}$ is the scalar curvature of the associated Kahler toric metric $\bar{g}$. The theorem follows immediately from the form of the A_r and the scalar curvature of a Kahler toric metric given in the previous section. □

7.2.3 The Extremal Kahler Condition

As discussed in the previous section, a Kahler metric is extremal if and only if the gradient of the scalar curvature, ∇S is a holomorphic vector field. Expressed in the complex coordinates $\partial_{z_i} = \partial_{u_i} + i\partial_{\phi_i}$, the gradient of the scalar curvature of a fiberwise Kahler toric metric is

$$\nabla S = \left(h^{kl} \frac{\partial S}{\partial u^l} \right) \frac{\partial}{\partial z^k}.$$

Because the coefficients, $h^{kl} \frac{\partial S}{\partial u^l}$, are real, the vector field ∇S is holomorphic if and only if these coefficients are constant. The extremal Kahler condition is equivalent to

$$h^{kl} \frac{\partial S}{\partial u^k} = \frac{\partial S}{\partial x^k} = \alpha_k$$

where the α_k are constants. This gives the following theorem.

Theorem 7.12. *A fiberwise Kahler toric metric is extremal Kahler if and only if its scalar curvature satisfies*

$$S = \sum_{k=1}^{d} \alpha_k x_k + \beta$$

where β and the α_k are constants.

Note the striking parallel between the extremal Kahler condition on a fiberwise Kahler toric metric and the extremal Kahler condition on a Kahler toric metric given in the previous section. This provides further support for the idea that fiberwise Kahler toric metrics can be viewed as Kahler toric metrics with additional structure.

Chapter 8

Cohomogeneity-One Fiberwise Kahler Toric Manifolds

The simplest class of fiberwise Kahler toric manifolds are those of cohomogeneity-one, i.e. $d = 1$. If (M, g, J) is a cohomogeneity-one manifold under the action of a compact connected semisimple Lie group G, then $\dim(L/K) = 1$. Therefore, $L/K \cong S^1$ and assumption one is automatically satisfied. Assumption two, however, is not automatically satisfied, although it is satisfied for a generic S^1-bundle over a coadjoint orbit G/L. The explanation for this can be found in [15]. Assumption two holds by definition in the fiberwise Kahler toric case and we will from now on assume that the summands in the isotropy representation are distinct.

From the calculations preformed above, the metric on a cohomogeneity-one fiberwise Kahler toric metric is of the form

$$g = \begin{pmatrix} \frac{1}{h} & & & & \\ & h & & & \\ & & A_1 Id_{d_1} & & \\ & & & \ldots & \\ & & & & A_m Id_{d_m} \end{pmatrix}$$

relative to basis $(\partial_x, \partial_\phi, Y_1^1, ..., Y_{d_m}^m)$ where we have set $h_{11} = h$, $x = x_1$, and $\partial_\phi = \partial_{\phi_1}$. Recall that $h = h(x)$ and $A_i = A_i(x)$ for all $1 \leq i \leq m$.

Furthermore, from the Kahler condition, we know that $A_i = b_i x + a_i$ for all i where b_i and a_i are constants. The metric can be written

$$g = \begin{pmatrix} \frac{1}{h} & & & & \\ & h & & & \\ & & (b_1 x + a_1) Id_{d_1} & & \\ & & & \ldots & \\ & & & & (b_m x + a_m) Id_{d_m} \end{pmatrix}.$$

The metric on a manifold of this type is determined by one function, h, in one variable, x, and $2m$ constants. Note that m of the $2m$ constants, the b_i determine the principal orbit types.

We now want to examine the constant scalar curvature, extremal Kahler, and Einstein conditions on a cohomogeneity-one fiberwise Kahler toric manifold After deriving explicit local solutions for each of these conditions, we will look at the global structure of such metrics. The Einstein case was already solved by Dancer and Wang in [15]. However, they did not, nor to the author's knowledge has anyone else, derive explicit solutions in the constant scalar curvature or the extremal Kahler case. Before, proceeding to new results, we first review the work of Dancer and Wang.

8.1 Explicit Integration of the Einstein Equations

Let λ be the Einstein constant. Following the calculations of the previous section, there are two types of equations in the Einstein condition. The first such equation arise from the condition $r(\partial_\phi, \partial_\phi) = \lambda g(\partial_\phi, \partial_\phi))$. Because the Ricci tensor, like the metric, is invariant under the action of complex structure, this equation contains the same information as the equation $r(\partial_u, \partial_u) = \lambda g(\partial_u, \partial_u))$ and therefore $r(\partial_x, \partial_x) = \lambda g(\partial_x, \partial_x))$. The second type of equation comes from $r(Y_k^i, Y_k^i) = \lambda g(Y_k^i, Y_k^i)$. The Einstein condition is equivalent to the following system of $m+1$ equations

$$-\frac{1}{2}h'' - \frac{1}{4}h'\left(\sum_{j=1}^{m}\frac{d_j b_j}{b_j x + a_j}\right) + \frac{1}{4}h\left(\sum_{j=1}^{m}\frac{d_j b_j^2}{(b_j x + a_j)^2}\right) = \lambda \tag{8.1}$$

and

$$-\frac{1}{2}h'\frac{b_i}{b_i x + a_i} - \frac{1}{4}\frac{b_i}{b_i x + a_i}h\left(\sum_{j=1}^{m}\frac{d_j b_j}{b_j x + a_j}\right) + \frac{1}{b_i x + a_i} = \lambda \tag{8.2}$$

for all $1 \leq i \leq m$, where $h' = \frac{dh}{dx}$. Equation (8.2) is equivalent to $1 = \lambda a_i$ if $b_i = 0$ and

$$-\frac{1}{2}h' - \frac{1}{4}h\left(\sum_{j=1}^{m}\frac{d_j b_j}{b_j x + a_j}\right) = \lambda x + \frac{a_i\lambda - 1}{b_i} \tag{8.3}$$

when $b \neq 0$. We noted above that not all of the b_i are equal to zero.

We see immediately that differentiating equation (8.3) gives equation (8.1). Therefore, equation (8.1) of the Einstein condition is automatically satisfied if equation (8.2) is. We now have the following theorem.

Theorem 8.1. *[15] A cohomogeneity-one fiberwise Kahler toric metric is Einstein if and only if*

- $1 = \lambda a_i$, *if* $b_i = 0$
- $\frac{1-\lambda a_i}{b_i} = \frac{1-\lambda a_k}{b_k} = D$, *if* $b_i b_k \neq 0$
- $h' + \frac{1}{2}h(\sum_{j=1}^{m}\frac{d_j b_j}{b_j x + a_j}) + 2\lambda x - 2D = 0$.

Proof. The first equation follows immediately from equation (8.3) when $b_i = 0$. The second equation follows from equation (8.3) when $i = k$. The third equation follows from the second. □

Dancer and Wang were able to integrate the Einstein equation to obtain an explicit solution for h

Theorem 8.2. *[15] If (M, g, J) is a cohomogeneity-one fiberwise Kahler toric metric, then*

$$h = \frac{2}{\prod_{j=1}^{m}(b_j x + a_j)^{\frac{d_j}{2}}} \int (D - \lambda x) \prod_{j=1}^{m} (b_j x + a_j)^{\frac{d_j}{2}} dx. \tag{8.4}$$

Note that the integrand of the expression for h is always a polynomial as each d_j is even and greater than or equal to two. Therefore, one can always, in principle, perform this integration and express h as a rational function in x.

8.2 Explicit Integration of the Scalar Curvature Equation

From the equations of the Einstein condition, it is straightforward to calculate the scalar curvature of (M, g, J)

Theorem 8.3. *If (M, g, J) is a cohomogeneity-one fiberwise Kahler toric manifold, the scalar curvature, $S = S(x)$, is given by*

$$S = -h'' - h' \left(\sum_{j=1}^{m} \frac{d_j b_j}{b_j x + a_j} \right) + h \left(\frac{1}{2} \sum_{j=1}^{m} \frac{d_j b_j^2}{(b_j x + a_j)^2} - \frac{1}{4} \left(\sum_{j=1}^{m} \frac{d_j b_j}{b_j x + a_j} \right)^2 \right) + \sum_{j=1}^{m} \frac{d_j}{b_j x + a_j}. \tag{8.5}$$

Note that $-h''$ is the scalar curvature $\bar{S}$ of the embedded two-dimensional Kahler toric metric.

We have been able to integrate this equation for *any* scalar curvature S. Moreover, we can obtain *explicit* local solutions in the constant scalar curvature and extremal Kahler cases.

Theorem 8.4. *If (M, g) is a cohomogeneity-one fiberwise Kahler toric manifold the function h satisfies*

$$h = \frac{1}{\prod_{j=1}^{m}(b_j x + a_j)^{\frac{d_j}{2}}} \int \left[\int \left(\sum_{j=1}^{m} \frac{d_j}{b_j x + a_j} - S \right) \prod_{j=1}^{m} (b_j x + a_j)^{\frac{d_j}{2}} dx \right] dx \tag{8.6}$$

where S is the scalar curvature of the manifold.

Proof. The explicit solution in the Einstein case suggests that h is of the form

$$h = \frac{1}{\prod_{j=1}^{m}(b_j x + a_j)^{\frac{d_j}{2}}} \int \left(\prod_{j=1}^{m} (b_j x + a_j)^{\frac{d_j}{2}} \right) \Theta dx$$

for some function $\Theta = \Theta(x)$. With this assumption the scalar curvature equation (8.5) becomes a first-order equation in Θ.

$$S = -\Theta' - \frac{1}{2}\Theta \left(\sum_{j=1}^{m} d_j \frac{b_j}{b_j x + a_j} \right) + \sum_{j=1}^{m} \frac{d_j}{b_j x + a_j}.$$

This equation is a simple first-order linear ordinary differential equation which we can integrate explicitly. We see that

$$\Theta = \frac{1}{\prod_{j=1}^{m}(b_j x + a_j)^{\frac{d_j}{2}}} \int \left(\sum_{j=1}^{m} \frac{d_j}{b_j x + a_j} - S \right) \prod_{j=1}^{m} (b_j x + a_j)^{\frac{d_j}{2}} dx.$$

Substituting this expression for Θ into our expression for h in terms of theta above completes the proof. □

Note that if we are looking for metrics of constant scalar curvature $S = \beta$, the we can perform the double integration in (8.6) explicitly and thereby obtain an explicit description of *any* cohomogeneity-one fiberwise Kahler toric metric of constant scalar curvature in terms of *rational* functions in x.

Furthermore, if we are looking instead for extremal Kahler metrics then, as proved above, $S = \alpha x + \beta$ where α and β are constants. Once again we can perform explicitly the double integration in (8.6), and express any extremal Kahler cohomogeneity-one fiberwise Kahler toric metric in terms of rational functions in x.

8.3 Special Orbits and Completeness

A general cohomogeneity-one metric must have zero, one, or two special orbits. If the manifold is compact, then any extremal Kahler metric will have two special orbits. This follows from the fact that the quotient space of any compact cohomogeneity-one manifold is either a closed interval or a copy of S^1 [5]. If the quotient space is S^1 then there are no special orbits and the metric must be periodic. However, this implies that all of the functions $A_i = b_i + a_i$ are periodic which can only occur if all of the b_i vanish. In this case the metric is a product metric. We have assumed above that not all of the b_i are zero so the metric must have two special orbits. (Below we will present some examples in which all of the b_i are zero but in general we assume that at least one is non-zero.) We will see below that a complete cohomogeneity-one fiberwise Kahler toric metric must have at least one. After determining the possible special orbit types, we will determine the smoothness conditions which h and the constants a_i and b_i must satisfy for the metric to be extended smoothly over the principal orbits.

Let G/H be a special orbit with $K \subset H$. Via translation, we can assume that the special orbit is located at $x = 0$.

Lemma 8.5. *[15] The moment map μ gives us a fiber bundle over*

$$L^*/H \to G/H \to G/L^*$$

where G/L^* is a coadjoint orbit. Furthermore, if V denotes the normal slice to the special orbit G/H then either $H = L^*$ or the complex structure, J, induces an H-equivariant isomorphism $ker d\mu \cong V$.

Proof. We know that V is an irreducible H-module and that if $v \in \ker(d\mu)$ then $Jv \in V$. These imply that J acts as an H-equivariant monomorphism from $\ker(d\mu)$ to V. If $\ker(d\mu)$ is trivial at the special orbit then $H = L^*$. If not, then the irreducibility of V requires that the map is an isomorphism. Finally, we note that $\pi(G/L^*) = \{1\}$ and therefore L^*/H is not a discrete set of more than one point. □

Dancer and Wang were able to demonstrate that, at a special orbit in the cohomogeneity-one fiberwise Kahler toric case, $H = L^*$. As we will attempt to extend this argument to the cohomogeneity-d case below, we will review their proof.

We begin by noting the following important facts: $L \subset L^*$ by semicontinuity and $H/K \cong S^k$ for some non-negative odd integer k for smoothness.

At the special orbits, we have the following Lie algebra decomposition of K-modules

$$\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{m}_1 \oplus \mathfrak{m}_2 \oplus \mathfrak{m}_3$$

where $\mathfrak{h} = \mathfrak{k} \oplus \mathfrak{m}_1$ and $\mathfrak{m}_3 = \ker(d\mu)$. If $H = L^*$ then $\ker(d\mu) = 0$. If not, the preceding lemma proves that J maps $\mathfrak{m}_3$ isomorphically onto $1 \oplus \mathfrak{m}_1 \approx V$. This would mean that $\mathfrak{m}_3 \cong 1 \oplus \mathfrak{m}_1$ as K-modules. However, this contradicts Assumption Two as $\mathfrak{m}_1$ would then appear twice in the isotropy representation. We conclude that $\mathfrak{m}_1 = 0$. This is equivalent to saying the collapsing sphere H/K has dimension zero. For the metric to be extended smoothly over this special orbit we require that $h(0) > 0$ while $A_i > 0$ and $A_i' = 0$. This contradicts the equations of the Kahler condition. Since $H = L^*$ and $\ker(d\mu) = 0$, the S^1 fiber must collapse at any special orbit and $h(0) = 0$.

The above discussion can be summarized as follows. Each special orbit occurs at a boundary point of the one-dimensional quotient space. At that point the S^1 fiber of the principal orbit *must* collapse. A portion of the coadjoint orbit G/L may or may not collapse along with it. That is, $H/K \cong S^1$ or $H/K \cong S^k$ for $k > 1$. This sphere admits the fibration

$$S^1 \cong L/K \to S^k \to H/L.$$

It remains to determine the possible H/L. It is straightforward to see that $H/L \cong \mathbb{C}P^{\frac{k-1}{2}}$.

Next, we look at the necessary conditions on the functions h A_i for the metric to be extended smoothly over a special orbit. Because of the invariance of the metric under translation in x, we can fix the location of any one special orbit to be $x = 0$ where the principal orbits occur at $x > 0$. We can write $\mathfrak{m}_1$ as

$$\mathfrak{m}_1 = \mathfrak{p}_1 \oplus ... \oplus \mathfrak{p}_l.$$

It is, of course, possible that $l = 0$; in fact, by the classification of coadjoint orbits, $l \leq 2$ [15].

For the metric to be smooth at $x = 0$, we must have

$$h(0) = 0, \quad \frac{dh}{dx}(0) = 2.$$

Before determining what the smoothness conditions on the A_i are, we note that the smoothness conditions on h at the special orbit are identical to the smoothness conditions at a special orbit of the toric Kahler manifold (see [1]) for smoothness conditions on special orbits of Kahler toric manifolds). The implicit manifold $(N, \bar{g})$ mentioned above is in fact a global Kahler toric manifold of cohomogeneity-one. That is, every fiberwise Kahler toric manifold (M, g, J) can be viewed as a toric Kahler manifold $(N, \bar{g}, \bar{J})$ along with a collection of $3m$ constants (the a_i, b_i, and d_i) along with conditions on these constants to insure that the metric is positive definite and smooth at the special orbits.

We return now to the conditions on the functions A_i at the special orbit $x = 0$ necessary for smoothness. Clearly,

$$A_i(0) = 0$$

for $1 \leq i \leq l$. Since $A_j = b_j x + a_j$, we see that $a_i = 0$ for $1 \leq i \leq l$. Since the principal orbits are located at positive values of x, $b_i > 0$ for $1 \leq i \leq l$.

There is one more condition on the b_i for $1 \leq i \leq l$ necessary for the metric to be extended smoothly over the special orbit at $x = 0$. To obtain this condition, we evaluate the scalar curvature equation (8.5) at $x = 0$. The scalar curvature is constant at $x = 0$ if the metric is smooth there so the left-hand side of this equation is smooth. However, the right-hand side appears to contain poles of order one at $x = 0$. By inspection, the right-hand side will be smooth at $x = 0$ if and only if

$$\sum_{j=1}^{l} \frac{d_j}{b_j} = \frac{k^2 - 1}{2}$$

where $k - 1 = \sum_{j=1}^{l} d_j$.

Let us consider in more detail the case in which each principal orbit is an S^1-bundle over a product of Kahler-Einstein manifolds $M_1 \times ... \times M_m$ as in construction two. Again, two things can happen at the special orbit $x = 0$. The S^1-fiber can collapse alone or the S^1-fiber can collapse along with, say, $M_1 \cong \mathbb{C}P^{\frac{d_1}{2}}$. In the second case, $l = 1$ and $k - 1 = d_1$ and we deduce that

$$b_1 = \frac{2}{d_1 + 2} \tag{8.7}$$

for the metric to be extended smoothly over the special orbit. We take this opportunity to be more precise about the meaning of the b_i from construction two. Take, say, M_i to be a Kahler-Einstein manifold with positive Einstein constant. The first Chern class of M_i can be written as

$$c_1(M_i) = p_i \alpha_i$$

where p_i is a positive integer and α_i is an indivisible integral cohomology class. Furthermore, if d_i is the real dimension of M_i, then $p \leq \frac{d_i+2}{2}$ with equality if and only if $M_i \cong \mathbb{C}P^{\frac{d_i}{2}}$ [38]. If ω_i is the Kahler form of the Kahler-Einstein metric on M_i then we have that $[\omega_i] = 2\pi p_i \alpha_i$. Finally, the

curvature form, Ω_i, of the connection θ_i of the S^1-bundle over M_i can be written as $\Omega_i = -2\pi q_i \alpha_i$ where q_i is some integer. From the definition of b_i given above we deduce that

$$b_i = -\frac{q_i}{p_i}.$$

In the case being considered $M_1 \cong \mathbb{C}P^{\frac{d_1}{2}}$ so $p_1 = \frac{d_1+2}{2}$. Therefore, equation (8.7) implies that $q = -1$.

Having determined the smoothness conditions for the metric to be extended over a special orbit, we now address the problem of finding *complete* constant scalar curvature and extremal Kahler metrics. Cohomogeneity-one metrics are defined on intervals in $\mathbb{R}$. Endpoints can, a priori, occur at poles of h, zeros of h, zeros of the A_i, or at $\pm\infty$.

Let p be an endpoint of the interval on which the metric is defined (of course p could be $\pm\infty$). Given the form of the metric g, the geodesic distance to an endpoint p can be written as

$$\int_*^p h^{-\frac{1}{2}} dx.$$

If p is a pole of h, then the geodesic distance is finite and there are no complete metrics. If h is nonzero and $A_i = 0$ for some i then again the geodesic distance is finite. We must have a special orbit at the endpoint which is impossible because $h = 0$ at all special orbits. Therefore, if the metric is to be complete, each endpoint must be either a zero of h (possibly also a zero of one or more of the A_i) or $\pm\infty$. Furthermore, if the manifold is noncompact, the above discussion proves that one of the endpoints must be $\pm\infty$.

For g an extremal Kahler metric, the scalar curvature is linear in x: $S = \alpha x + \beta$. In equation (8.6), we express h as a double integral involving S. If $\alpha = 0$, then $h = O(x^2)$ or $O(x)$ as $x \to \pm\infty$ and the geodesic distance is infinite if $p = \pm\infty$. A constant scalar curvature metric will be complete as we approach infinity. On the other hand, if $\alpha \neq 0$, then $h = O(x^3)$ as $x \to \pm\infty$ and the geodesic distance is *finite* and the metric is not complete at $\pm\infty$. This proves the following theorem.

Theorem 8.6. *Let (M, g, J) be a non-compact cohomogeneity-one fiberwise Kahler toric manifold. There are no complete extremal Kahler metrics on M which are not of constant scalar curvature*

8.4 Non-Compact Metrics of Constant Scalar Curvature

As proved above, any complete extremal Kahler metric on a non-compact cohomogeneity-one fiberwise Kahler toric manifold must have constant scalar curvature. Therefore, we can restrict our attention to the search for constant scalar curvature metrics. From the form of the metric we deduce that g must have at least one special orbit. This can be seen by noting that each A_i vanishes at some value of x. Moreover, if the manifold is non-compact, then there are no other special orbits.

By translating in x if necessary, we take the location of the special orbit to be $x = 0$. We construct the metric so that the principal orbits occur at $x > 0$ and that the metric extends to $+\infty$.

Let $S = \beta$, a constant. For the metric to be smooth and positive definite, the following conditions must be satisfied

- $h(0) = 0$ and $h'(0) = 2$,
- $A_i = b_i x + a_i > 0$ for $x \in (0, \infty)$ which implies that $b_i \geq 0$ and $a_i \geq 0$ for all i, and
- $h(x) > 0$ for $x \in (0, \infty)$.

If $A_i(0) = 0$ for some i (i.e. if $a_i = 0$ for some i) then an additional smoothness condition must be satisfied. Let us first consider the case in which $A_i = 0$ for $x \in (0, \infty)$ (that is $a_i > 0$). In this case the manifold is a complex line bundle over a coadjoint orbit (or a product of Kahler-Einstein manifolds).

The function h must satisfy equation (8.6). We set $h = \frac{P(x)}{Q(x)}$ where

$$P(x) = \int \left[\int \left(\sum_{j=1}^{m} \frac{d_j}{b_j x + a_j} - \beta \right) \prod_{j=1}^{m} (b_j x + a_j)^{\frac{d_j}{2}} dx \right] dx$$

and

$$Q(x) = \prod_{j=1}^{m} (b_j x + a_j)^{\frac{d_j}{2}}.$$

Note that because we have assumed that $b_j x + a_j > 0$ for $x \in [0, \infty)$, we have that $Q(x) > 0$ for $x \in [0, \infty)$. For the metric to be positive, $P > 0$ for $x \in (0, \infty)$. Both P and Q are polynomials in x because each of the d_i is even and is greater than or equal to two. After integration, P can be written as

$$P(x) = f + ex + ... - \beta \frac{\left(\prod_{j=1}^{m} b_j^{\frac{d_j}{2}} \right)}{(l+2)(l+1)} x^{l+2}$$

where e and f are the constants of integration and $l = \sum_{j=1}^{m} \frac{d_j}{2}$. The smoothness conditions at $x = 0$ determine the constants. To see this we note that $h(0) = 0$ implies that $P(0) = 0$ as $Q(0) > 0$ by assumption. Therefore

$$h(0) = 0 \Rightarrow P(0) = 0 \Rightarrow f = 0.$$

Next, we note that $h'(0) = \frac{P'(0)}{Q(0)}$ since $P(0) = 0$. We see that

$$h'(0) = 2 \Rightarrow \frac{P'(0)}{Q(0)} = 2 \Rightarrow \frac{e}{\prod_{j=1}^{m} a_j^{\frac{d_j}{2}}} = 2.$$

This implies that

$$P(x) = 2 \prod_{j=1}^{m} a_j^{\frac{d_j}{2}} x + ... - \beta \frac{\left(\prod_{j=1}^{m} b_j^{\frac{d_j}{2}} \right)}{(l+2)(l+1)} x^{l+2}.$$

It remains only to check that $h(x) > 0$ for $x \in (0, \infty)$. This is equivalent to demanding that $P(x) > 0$ for $x \in (0, \infty)$. Now, as $x \to \infty$,

$$P(x) \sim -\beta \frac{\left(\prod_{j=1}^{m} b_j^{\frac{d_j}{2}}\right)}{(l+2)(l+1)} x^{l+2}.$$

If $\beta > 0$, then $P(x) \to -\infty$ as $x \to \infty$ and the metric will fail to be positive for large values of x. Therefore, $\beta \leq 0$.

If $\beta \leq 0$, then we see that in the polynomial $P(x)$, all of the coefficients of the x^k are positive as $b_i \geq 0$ and $a_i > 0$ for all i. Therefore, $P(x) > 0$ for $x \in (0, \infty)$. All of the smoothness conditions and the positivity condition are satisfied for all values of $a_i > 0$. As discussed above, it is the values of the b_i's which determine the principal orbit type. So given a collection of $b_i > 0$, we see that there exists a scalar flat metric for all values of a_i as well as a metric with scalar curvature, say, -1. We have the following theorem.

Theorem 8.7. *Let (M, g) be a noncompact cohomogeneity-one fiberwise Kahler toric manifold defined on the interval $x \in [0, \infty)$ such that $b_i \geq 0$ for all i (i.e. M is a complex line bundle) as described above. Every such manifold admits a scalar-flat metric ($\beta = 0$) as well as a negative scalar curvature metric ($\beta = -1$) for every value of $a_i > 0$. That is, each such manifold admits an m-parameter family of scalar-flat metrics and an m-parameter family of metrics with scalar curvature equal to -1.*

On the other hand, if $a_i = 0$ when $1 \leq i \leq l$ for some l then there is the additional smoothness condition that $\sum_{i=1}^{l} \frac{d_i}{b_i} = \frac{k^2-1}{2}$.

Consider the case in which each principal orbit is an S^1-bundle over the product of Kahler-Einstein manifolds $M_1 \times ... \times M_m$. Consider a special orbit in which $A_1(0) = 0$ and $A_i(0) > 0$ for all $2 \leq i \leq m$. We demonstrated above that $M_1 \cong \mathbb{C}P^{\frac{d_1}{2}}$, $a_1 = 0$, and $b_1 = \frac{2}{d_1+2}$. In the constant scalar curvature case the function h can be written as $h = \frac{P(x)}{Q(x)}$ where

$$P = \int \left[\int \left(\frac{d_1(d_1+2)}{2x} + \sum_{j=2}^{m} \frac{d_j}{b_j x + a_j} - \beta \right) x^{\frac{d_1}{2}} \prod_{j=2}^{m} (b_j x + a_j)^{\frac{d_j}{2}} dx \right] dx$$

and

$$Q = x^{\frac{d_1}{2}} \prod_{j=2}^{m} (b_j x + a_j)^{\frac{d_j}{2}}.$$

Both P and Q are polynomials and we can write P schematically as

$$P = -\beta \frac{\prod_{j=2}^{m} b_j^{\frac{d_j}{2}}}{(l+2)(l+1)} x^{l+2} + ... + 2 \left(\prod_{j=2}^{m} a_j^{\frac{d_j}{2}} \right) x^{\frac{d_1+2}{2}} + ex + f$$

where $l = \sum_{j=1}^{m} \frac{d_j}{2} = \frac{n-2}{2}$ and e and f are the constants of integration. The polynomial Q can be written as

$$Q = \left(\prod_{j=2}^{m} b_j^{\frac{d_j}{2}} \right) x^l + ... + \left(\prod_{j=2}^{m} a_j^{\frac{d_j}{2}} \right) x^{\frac{d_1}{2}}.$$

It is straightforward to check that $h(0) = 0$ and $h'(0) = 2$ if and only if $e = f = 0$.

All that remains is to determine when $h(x) > 0$ for all $x \in (0, \infty)$. Recalling that $b_i > 0$ and $a_i > 0$ for all $2 \leq i \leq m$, we see that we obtain a smooth positive definite metric if $\beta \leq 0$. We see, as above, that h will not be positive for large values of x if β is positive. Finally, we note that the constant scalar curvature metrics we have here constructed are on $\mathbb{C}^{\frac{d_1+2}{2}}$-bundles over $M_2 \times ... \times M_m$.

Theorem 8.8. *Let (M, g) be a fiberwise Kahler toric manifold in which each principal orbit is an S^1-bundle over $M_1 \times ... \times M_m$ such that $M_1 \cong \mathbb{C}P^{\frac{d_1}{2}}$ and $b_1 = \frac{2}{d_1+2}$. Let g be defined on the interval $x \in [x, \infty)$ such that $b_i \geq 0$ for all $1 \leq i \geq m$. In this case M is a $\mathbb{C}^{\frac{d_1+2}{2}}$-bundle over $M_2 \times ... \times M_m$. Every such manifold admits a scalar-flat metric and a metric with scalar curvature equation to -1 for all values of $a_i > 0$ for $2 \leq i \leq m$.*

8.5 Compact Extremal Kahler Metrics

Having found all the non-positive constant scalar curvature metrics on non-compact cohomogeneity-one fiberwise Kahler toric metrics, we now turn our attention to the case of extremal Kahler metrics on compact manifolds. In the compact case, the metric must have two special orbits at which h and possibly some A_i's vanish. The presence of a second special orbit makes the problem of finding constant scalar curvature Kahler metrics substantially more difficult in the compact case than it was in the non-compact case. In fact, there are known examples of compact cohomogeneity-one fiberwise Kahler toric manifolds which do not admit any constant scalar curvature metrics (e.g. any Hirzebruch surface other that $S^2 \times S^2$ with its product metric [6]). However, it is reasonable to suppose that every compact cohomogeneity-one fiberwise Kahler toric metric could admit an extremal Kahler metric.

We begin by looking at the case of $\mathbb{C}P^1$-bundles.

8.5.1 Extremal Kahler Metrics on $\mathbb{C}P^1$-Bundles

Let (M, g, J) be a compact fiberwise Kahler toric manifold of cohomogeneity-one. If both of the special orbits occur when the S^1-fiber collapses and $A_i \neq 0$ for all i, then the total manifold M is a $\mathbb{C}P^1$-bundle over a coadjoint orbit G/L or the product of Kahler-Einstein manifolds $M_1 \times ... \times M_m$ each with positive Einstein constant (though of course the product metric need not be Einstein).

By translation, we can fix the location of the first special orbit to be at $x = 0$. By rescaling if necessary, we can fix the location of the second special orbit to be located at, say, $x = 1$. We have

assumed that at both of the special orbits $A_i \neq 0$ for all i. For the metric to be smooth on the special orbits and positive for $x \in (0,1)$ the following conditions must be satisfied:

- $h(0) = 0$ and $h'(0) = 2$
- $h(1) = 0$ and $h'(1) = -2$
- $A_i = b_i x + a_i > 0$ for $x \in [0,1]$ which implies that $a_i > 0$ for all i and $-\frac{a_i}{b_i} < 0$ or $-\frac{a_i}{b_i} > 1$ for all i
- $h(x) > 0$ for $x \in (0,1)$.

We ask which manifolds of this type admit extremal Kahler metrics. In the extremal Kahler case, $S = \alpha x + \beta$ and

$$h = \frac{1}{\prod_{j=1}^{m}(b_j x + a_j)^{\frac{d_j}{2}}} \int \left[\int \left(\sum_{j=1}^{m} \frac{d_j}{b_j x + a_j} - \alpha x - \beta \right) \prod_{j=1}^{m} (b_j x + a_j)^{\frac{d_j}{2}} dx \right] dx.$$

We write h as $h(x) = \frac{P(x)}{Q(x)}$ where $P(x) = \int \left[\int (\sum_{j=1}^{m} \frac{d_j}{b_j x + a_j} - \alpha x - \beta) \prod_{j=1}^{m} (b_j x + a_j)^{\frac{d_j}{2}} dx \right] dx$ and $Q(x) = \prod_{j=1}^{m} (b_j x + a_j)^{\frac{d_j}{2}}$. By construction $Q(x) > 0$ for $x \in [0,1]$. Schematically,

$$P(x) = f + ex + ... - \alpha \frac{\prod_{j=1}^{m} b_j^{\frac{d_j}{2}}}{(l+3)(l+2)} x^{l+3}$$

where e and f are the constants of integration and $l = \sum_{j=1}^{m} \frac{d_j}{2}$. The condition $h(0) = 0$ holds if and only if $P(0) = 0$. This implies that $f = 0$. Next, consider the condition $h'(0) = 2$. Since $P(0) = 0$, we have that $h'(0) = \frac{P'(0)}{Q(0)}$ which implies that $e = 2 \prod_{j=1}^{m} a_j^{\frac{d_j}{2}}$. The two constants of integration are then completely determined by the smoothness conditions at the special orbit located at $x = 0$. Moreover,

$$P(x) = 2 \prod_{j=1}^{m} a_j^{\frac{d_j}{2}} x + ... - \alpha \frac{\prod_{j=1}^{m} b_j^{\frac{d_j}{2}}}{(l+3)(l+2)} x^{l+3}.$$

There are two remaining smoothness conditions at the second special orbit. The condition $h(1) = 0$ is equivalent to $P(1) = 0$. This can be satisfied by fixing the value of, say, the free variable β. The condition $h'(1) = -2$ is equivalent to

$$P'(1) = -2Q(1) = -2 \prod_{j=1}^{m} (b_j + a_j)^{\frac{d_j}{2}}$$

since $P(1) = 0$. We can always choose α so that this equation is satisfied. The resulting function h has zeroes of order one at the points $x = 0$ and $x = 1$ and its first derivative satisfies the smoothness conditions at those points. However, it is not immediately obvious that the resulting function h will be greater than zero on the open interval $(0,1)$. That is, the function could have zeroes in the interval $(0,1)$.

Before proving this sections main theorem, we pause to recall the significance of the constants b_i and a_i. The b_i determine the principal orbit type; changing the values of the b_i changes the connection of the S^1-bundle. It can therefore be said that it is the b_i that determine the complex manifold. On the other hand, changing the a_i (so long as one does not violate smoothness) does not change the manifold. It does however change the Kahler class of the Kahler form.

Therefore, we consider the b_i to be fixed, and we take the a_i to be free parameters. Above, we have shown that for every value of a_i we can construct a function $h_{a_1 \dots a_m}$ which satisfies the extremal Kahler condition at each point and also satisfies all of the smoothness conditions. If $h_{a_1 \dots a_m}$ is positive for $x \in (0,1)$, then we have a global extremal Kahler metric. What remains is to determine at what values of a_i the function $h_{a_1 \dots a_m} > 0$ on that open interval.

There are two possible questions that we could ask. First, we could ask which manifolds admit at least one extremal Kahler metric. Second, we could ask which Kahler classes admit an extremal Kahler metric. In the following theorem, we give a complete answer to the first question for manifolds of the type being considered.

Theorem 8.9. *Let (M, g, J) be a fiberwise Kahler toric manifold of cohomogeneity-one such that M is a $\mathbb{C}P^1$-bundle over a coadjoint orbit G/L whose isotropy representation has m distinct summands or the product of m Kahler- Einstein manifolds M_k. **Every** such manifold admits, up to scaling, an m-parameter family of extremal Kahler metrics.*

Proof. The a_i are free parameters and for each value of the a_i we can construct a function $h_{a_1 \dots a_m}$ with zeros of order one at $x = 0, 1$ which satisfies the smoothness conditions. The function $h_{a_1 \dots a_m}$ defines an extremal Kahler metric on M if and only if

$$h_{a_1 \dots a_m} > 0 \quad \text{for} \quad x \in (0,1).$$

In this proof, by looking at the limit of the functions $h_{a_1 \dots a_m}$ as $a_i \to \infty$ for *all* i, we will demonstrate that the function $h_{a_1 \dots a_m}$ is positive on $x \in (0,1)$ for the a_i sufficiently large.

Let $h_{a_1 \dots a_m} = \frac{P_{a_1 \dots a_m}}{Q_{a_1 \dots a_m}}$ as above. Grouping the terms by the order of the a_i we write $P_{a_1 \dots a_m}$ as

$$P_{a_1 \dots a_m} = \prod_{j=1}^{m} a_j^{\frac{d_j}{2}} (2x - \frac{\alpha x^3}{6} - \frac{\beta x^2}{2}) + H$$

where H consists of lower order terms in the a_i. Also,

$$Q_{a_1 \dots a_m} = \prod_{j=1}^{m} a_j^{\frac{d_j}{2}} + G$$

where G consists of lower order terms in the a_i.

For a_i sufficiently large for all i,

$$P_{a_1 \dots a_m} \sim \prod_{j=1}^{m} a_j^{\frac{d_j}{2}} (2x - \frac{\alpha x^3}{6} - \frac{\beta x^2}{2})$$

and

$$Q_{a_1 \ldots a_m} \sim \prod_{j=1}^{m} a_j^{\frac{d_j}{2}}.$$

Therefore,

$$h_{a_1 \ldots a_m} \sim 2x - \frac{\alpha x^3}{6} - \frac{\beta x^2}{2}.$$

The conditions $h(1) = 0$ and $h'(1) = -2$ imply that for the a_i sufficiently large

$$\alpha \sim 0,$$

$$\beta \sim 4,$$

and

$$h_{a_1 \ldots a_m} \sim 2x(1-x) \equiv h_\infty.$$

Recall that each of the $h_{a_1 \ldots a_m}$ is a smooth function on $[0,1]$ with zeros of order one at $x = 0, 1$. As the a_i tend to infinity for all i, these functions approach a smooth function, h_∞, on $[0,1]$ with zeros of order one at $x = 0, 1$. Since h_∞ is positive on the interval $(0,1)$, we deduce that, for the a_i sufficiently large for all i, the function $h_{a_1 \ldots a_m}$ must be positive on the interval $(0,1)$ also. Therefore, for the a_i sufficiently large, $h_{a_1 \ldots a_m}$ defines an extremal Kahler metric on M. We therefore have the desired m parameter family of extremal Kahler metrics.

Note that while $\alpha \to 0$ as the a_i approach infinity, α may be non-zero and we cannot deduce that the metrics are of constant scalar curvature for the a_i sufficiently large. In fact, there are known examples of manifolds of this type which admit no constant scalar curvature Kahler metrics. □

This theorem provides a wealth of new examples of extremal Kahler metrics. Furthermore, these metrics can be written down *explicitly* in terms of rational functions. Previously, a one-parameter family of extremal Kahler metrics were found on $\mathbb{C}P^1$-bundles over $\mathbb{C}P^l$ [6]. The extremal Kahler metrics we have found can be viewed as a generalization of those metrics.

8.5.2 Extremal Kahler Metrics on $\mathbb{C}P^{\frac{d_1+2}{2}}$-Bundles

In the previous section, we assumed that only the S^1-fiber collapsed at the special orbits. In this section, we set

$$A_1(0) = 0$$

while keeping $A_i(x) > 0$ when $x \in [0,1]$ for all $2 \leq i \leq m$. This condition implies that $a_1 = 0$ and $b_1 > 0$.

For simplicity of presentation, we restrict ourselves to the case in which each principal orbit is an S^1-bundle over $M_1 \times \ldots \times M_m$. As described above, for the metric to be extended smoothly over the

special orbit, we must have $M_1 \cong \mathbb{C}P^{\frac{d_1}{2}}$ and $b_1 = \frac{2}{d_1+2}$. In the extremal Kahler case, the function h can be written as $h(x) = \frac{P(x)}{Q(x)}$ where

$$P = \int \left[\int \left(\frac{d_1(d_1+2)}{2x} + \sum_{j=2}^{m} \frac{d_j}{b_j x + a_j} - \alpha x - \beta \right) x^{\frac{d_1}{2}} \prod_{j=2}^{m} (b_j x + a_j)^{\frac{d_j}{2}} dx \right] dx$$

and

$$Q = x^{\frac{d_1}{2}} \prod_{j=2}^{m} (b_j x + a_j)^{\frac{d_j}{2}}.$$

Both P and Q are polynomials in x; they can be written schematically as

$$P = -\alpha \frac{\prod_{j=2}^{m} b_j^{\frac{d_j}{2}}}{(l+3)(l+2)} x^{l+3} + ... + 2 \left(\prod_{j=2}^{m} a_j^{\frac{d_j}{2}} \right) x^{\frac{d_1+2}{2}} + ex + f$$

where $l = \sum_{j=1}^{m} \frac{d_j}{2} = \frac{n-2}{2}$ and e and f are the constants of integration. The polynomial Q can be written as

$$Q = \left(\prod_{j=2}^{m} b_j^{\frac{d_j}{2}} \right) x^l + ... + \left(\prod_{j=2}^{m} a_j^{\frac{d_j}{2}} \right) x^{\frac{d_1}{2}}.$$

It is straightforward to check that $h(0) = 0$ and $h'(0) = 2$ if and only if $e = f = 0$. There are two remaining smoothness conditions are $h(1) = 0$ and $h'(1) = -2$. As in the previous section, these two conditions will determine the values of the constants α and β. After selecting α and β to satisfy those conditions, we need only check that the resulting function h is positive for $x \in (0, 1)$.

We distinguish between two cases: $m = 1$ and $m > 1$. If $m = 1$, then it is straightforward to calculate that

$$h = -\frac{4\alpha}{(d_1+6)(d_1+4)} x^3 - \frac{4\beta}{(d_1+4)(d_1+2)} x^2 + 2x.$$

Solving the equations $h(1) = 0$ and $h'(0) = -2$, we see that $\alpha = 0$, $\beta = \frac{1}{2}(d_1+4)(d_1+2)$, and

$$h = 2x(1-x).$$

This gives the Fubini-Study metric on $M \cong \mathbb{C}P^{\frac{d_1+2}{2}}$.

The case $m > 1$ is the more interesting. Since the conditions $h(1) = 0$ and $h'(1) = -2$ determine the values of α and β, we have shown that for every value of a_i we can construct a function $h_{a_2 \ldots a_m}$ which satisfies the extremal Kahler condition at each point and also satisfies all of the smoothness conditions. If $h_{a_2 \ldots a_m} > 0$ for $x \in (0, 1)$, then we have constructed a global extremal Kahler metric on a $\mathbb{C}P^{\frac{d_1+2}{2}}$-bundle over $M_2 \times ... \times M_m$.

Theorem 8.10. *Let (M, g, J) be a fiberwise Kahler toric metric of cohomogeneity-one such that M is a $\mathbb{C}P^{\frac{d_1+2}{2}}$-bundle over a product of Kahler-Einstein manifolds $M_2 \times ... \times M_m$ as described above. **Every** such manifold admits, up to scaling, an $(m-1)$ parameter family of extremal Kahler metrics.*

Proof. We have already considered the case in which $m = 1$. When $m > 1$, the function $h_{a_2 \ldots a_m}$ defines an extremal Kahler metric on M if and only if

$$h_{a_2 \ldots a_m} > 0 \text{ for } x \in (0, 1).$$

Letting $h_{a_2 \ldots a_m} = \frac{P_{a_2 \ldots a_m}}{Q_{a_2 \ldots a_m}}$ as above, we see that for a_i sufficiently large for all i,

$$P_{a_2 \ldots a_m} \sim \left(\prod_{j=2}^{m} a_j^{\frac{d_j}{2}} \right) \left(-\frac{4\alpha}{(d_1+6)(d_1+4)} x^{\frac{d_1}{2}+3} - \frac{4\beta}{(d_1+4)(d_1+2)} x^{\frac{d_1}{2}+2} + 2x^{\frac{d_1}{2}} \right)$$

and

$$Q_{a_2 \ldots a_m} \sim \left(\prod_{j=2}^{m} a_j^{\frac{d_j}{2}} \right) x^{\frac{d_1}{2}}.$$

Therefore,

$$h_{a_2 \ldots a_m} \sim -\frac{4\alpha}{(d_1+6)(d_1+4)} x^3 - \frac{4\beta}{(d_1+4)(d_1+2)} x^2 + 2x.$$

The conditions $h(1) = 0$ and $h'(0) = -2$ imply that for the a_i sufficiently large

$$\alpha \sim 0,$$

$$\beta \sim \frac{1}{2}(d_1+4)(d_1+2),$$

and

$$h_{a_2 \ldots a_m} \sim 2x(1-x) \equiv h_\infty.$$

The function h_∞ is positive on the interval $x \in (0, 1)$. Therefore, by the same reasoning used in the proof of the theorem of the previous section, we deduce that $h_{a_2 \ldots a_m} > 0$ for $x \in (0, 1)$ for the a_i sufficiently large. This gives the $(m-1)$-parameter family of extremal Kahler metrics. □

8.5.3 Constant Scalar Curvature Kahler Metrics on $\mathbb{C}P^1$-Bundles

We have just demonstrated the existence of extremal Kahler metrics on compact codimension-one fiberwise Kahler toric manifolds in which only the S^1 fiber collapses on the special orbits. Globally, these manifolds are $\mathbb{C}P^1$-bundles. We found that all such manifolds admit at least one family of extremal Kahler metrics. In this section, we ask which if these $\mathbb{C}P^1$-bundles admit at least one constant scalar curvature metrics. Unlike in the general extremal Kahler case, we will see below that there are manifolds of this type which admit no constant scalar curvature metric.

In the compact constant scalar curvature case,

$$h = \frac{1}{\prod_{j=1}^{m} (b_j x + a_j)^{\frac{d_j}{2}}} \int \left[\int \left(\sum_{j=1}^{m} \frac{d_j}{b_j x + a_j} - \beta \right) \prod_{j=1}^{m} (b_j x + a_j)^{\frac{d_j}{2}} dx \right] dx$$

where β is a positive constant. By translation and rescaling, we again set the location of the special orbits to be $x = 0$ and $x = 1$. We again assume that at both special orbits $A_i \neq 0$ for all $1 \leq i \leq m$.

The same four smoothness conditions hold in the constant scalar curvature case as did in the extremal Kahler case.

If one of the b_i vanishes, say b_1, then the function h becomes

$$h = \frac{1}{\prod_{j=2}^{m}(b_j x + a_j)^{\frac{d_j}{2}}} \int \left[\int \left(\sum_{j=2}^{m} \frac{d_j}{b_j x + a_j} - (\beta - \frac{d_1}{a_1}) \right) \prod_{j=2}^{m} (b_j x + a_j)^{\frac{d_j}{2}} dx \right] dx.$$

Setting $\bar{\beta} = \beta - \frac{d_1}{a_1}$, we see that the problem of finding a constant scalar curvature metric is unaltered by the presence of absence of terms with $b_i = 0$. Therefore, we can without loss of generality assume that $b_i \neq 0$ for all i. As it is the b_i's which determine the manifold, the question we ask is: Given a collection of b_i, when is it possible to choose the a_i so that the metric is of constant scalar curvature and all of the smoothness conditions can be satisfied?

There are two general cases: either $b_i > 0$ (equivalently $b_i < 0$) for all $1 \leq i \leq m$ or some of the b_i are positive and some are negative. We consider the second case in the following theorem:

Theorem 8.11. *Let (M, g, J) be a fiberwise Kahler toric manifold of cohomogeneity-one such that M is a $\mathbb{C}P^1$-bundle as above. If there exists $b_i > 0$ for some $1 \leq i \leq m$ and $b_j < 0$ for some $1 \leq j \leq m$, then M admits at least a 1-parameter family of constant positive scalar curvature Kahler metric.*

Proof. By reordering if necessary, let $b_1, ..., b_l > 0$ and $b_{l+1}, ..., b_m < 0$ for some $2 \leq l \leq m$. Let $c_i = \frac{a_i}{|b_i|}$. We have

$$V^{\frac{1}{2}} = \prod_{j=1}^{m} (b_j x + a_j)^{\frac{d_j}{2}} = \prod_{j=1}^{m} |b_j|^{\frac{d_j}{2}} \prod_{j=1}^{l} (x + c_j)^{\frac{d_j}{2}} \prod_{j=l+1}^{m} (c_j - x)^{\frac{d_j}{2}}$$

and

$$\sum_{j=1}^{m} \frac{d_j}{b_j x + a_j} = \sum_{j=1}^{l} \frac{\frac{d_j}{|b_j|}}{c_j + x} + \sum_{j=l+1}^{m} \frac{\frac{d_j}{|b_j|}}{c_j - x}.$$

To prove this theorem, we need only find constant scalar curvature metrics for some 1-parameter family of the a_i. To this end, we make the following assumption which simplifies the calculations.

Set

$$c_j = c > 0$$

for all $1 \leq j \leq l$ and set

$$c_i = e > 1$$

for all $l + 1 \leq i \leq m$.

This implies that

$$h = \frac{1}{(x + c)^p (e - x)^q} \int \left[\int \left(\frac{r}{x + c} + \frac{t}{e - x} - \beta \right) (x + c)^p (e - x)^q dx \right] dx$$

where $p = \sum_{j=1}^{l} \frac{d_j}{2}$, $q = \sum_{j=l+1}^{m} \frac{d_j}{2}$, $r = \sum_{j=1}^{l} \frac{d_j}{|b_j|}$, and $t = \sum_{j=l+1}^{m} \frac{d_j}{|b_j|}$ which are all greater than zero and independent of the c_j.

Set $h(x) = \frac{T(x)}{L(x)}$ where

$$T(x) = \int \left[\int \left(\frac{r}{x+c} + \frac{t}{e-x} - \beta \right) (x+c)^p (e-x)^q dx \right] dx$$

and

$$L(x) = (x+c)^p (e-x)^q.$$

Schematically, we can write T as

$$T = F_{p+q+2} x^{p+q+2} + ... + F_w x^w + ... + Ex + F$$

where E and F are the constants of integration and the F_w are polynomials in c, e, r, t, p, q, and β. For the metric to be smooth at the special orbit $x = 0$, we must have $h(0) = 0$ and $h'(0) = 2$. These conditions determine the constants E and F. Because $L(x) > 0$ for $x \in [0, 1]$, $h(0) = 0$ implies that $T(0) = 0$ which is equivalent to saying that $F = 0$. Furthermore, we see that $h'(0) = 2$ implies that $\frac{T'(0)}{L(0)} = 2$. This is equivalent to $E = 2c^p e^q$.

The two remaining conditions $h(1) = 0$ and $h'(1) = -2$ determine β and place a condition on c and e.

To obtain a constant scalar curvature metric, it will be helpful to group the terms of T and L by the order of c and e. It is straightforward to see that

$$T(x) = [c^{p-1} e^{q-1} (re + tc) \frac{x^2}{2} + ...] - \beta [c^p e^q \frac{x^2}{2} + c^{p-1} e^{q-1} (pe - qc) \frac{x^3}{6} + ...] + 2c^p e^q x.$$

Here the $'...'$ refers to lower order terms in c and e. The condition $h(1) = 0$ implies that $T(1) = 0$. This condition determines the value of $S = \beta$,

$$\beta = \frac{2c^p e^q + \frac{1}{2} c^{p-1} e^{q-1} (re + tc) + ...}{\frac{1}{2} c^p e^q + \frac{1}{6} c^{p-1} e^{q-1} (pe - qc) + ...}. \tag{8.8}$$

As $c \to \infty$ and $e \to \infty$ we see that $\beta \to 4$. The next condition at the special orbit $x = 1$ is $h'(1) = -2$ which is equivalent to $\frac{T'(1)}{L(1)} = -2$. This condition can be interpreted as a condition on c and e. To calculate this condition, we first note that

$$L(1) = c^p e^q + c^{p-1} e^{q-1} (pe - qc) + ...$$

and

$$T'(1) = [2c^p e^q + c^{p-1} e^{q-1} (re + tc) + ...] - \beta [c^p e^q + \frac{1}{2} c^{p-1} e^{q-1} (pe - qc) + ...].$$

The condition $h'(1) = -2$ becomes

$$\beta [c^p e^q + \frac{1}{2} c^{p-1} e^{q-1} (pe - qc) + ...] = 4c^p e^q + c^{p-1} e^{q-1} (re + tc) + 2c^{p-1} e^{q-1} (pe - qc) + ...$$

Substituting equation (8.8) into this equation we have the condition

$$\frac{2}{3}c^{2p-1}e^{2q-1}(pe-qc)-\frac{1}{12}c^{2p-2}e^{2q-2}(pe-qc)(re+tc)+\frac{1}{3}c^{2p-2}e^{2q-2}(pe-qc)^2+...=0$$

This condition is equivalent to

$$\frac{2}{3}c^{2p-1}e^{2q-1}(pe-qc)+K(c,e)=0$$

where $K=K(c,e)$ is a polynomial in c and e in which each of the terms in K is of the form $K_{\eta\xi}c^{\eta}e^{\xi}$ such that $\eta\leq 2p$, $\xi\leq 2q$, $\eta+\xi<2p+2q-1$, and the $K_{\eta\xi}$ are constants depending on the fixed values of r, t, p, and q. Dividing by $c^{2p-1}e^{2q-1}$ implies that

$$B(c,e)\equiv pe-qc+\frac{3}{2}\frac{K(c,e)}{c^{2p-1}e^{2q-1}}=0. \tag{8.9}$$

As e and c go to infinity,

$$\frac{3}{2}\frac{K(c,e)}{c^{2p-1}e^{2q-1}}\to\bar{K}$$

for some constant $\bar{K}$. For all $0<\epsilon\ll 1$, there exists $N_\epsilon\gg 0$ such that $|\frac{3}{2}\frac{K(c,e)}{c^{2p-1}e^{2q-1}}-\bar{K}|<\epsilon$ whenever $pe, qc>N_\epsilon$. Set $pe=2N_\epsilon+2\epsilon$. If $qc=2N_\epsilon+\bar{K}+\epsilon$ then the quantity $B=B(c,e)>0$. If, on the other hand, $qc=2N_\epsilon+\bar{K}+3\epsilon$, then $B<0$. By the intermediate value theorem, there exists $qc\in(2N_\epsilon+\bar{K}+\epsilon,2N_\epsilon+\bar{K}+3\epsilon)$ such that $B=0$ and equation (8.9) is satisfied.

Therefore as $qc\to\infty$ the condition $h'(1)=-2$ can be solved by setting pe equal to a value increasingly close to $qc-\bar{K}$. Since q and p are positive constants, we have a family of solutions to the smoothness conditions in which c and e increase together. These solutions have zeros of order one at $x=0,1$. We must show that the solutions are positive for $x\in(0,1)$ for e and c sufficiently large. It is straightforward to see that as c and e approach infinity

$$h\sim 2x(1-x).$$

The family of solutions to the smoothness conditions tend to a function which has zeros of order one at $x=0,1$ and is positive for $x\in(0,1)$. By the same argument used in the proof of the existence of extremal Kahler metrics above, we see that for c and e sufficiently large, the function h will be positive for $x\in(0,1)$ and will define a constant scalar curvature metric.

To summarize, we have shown the existence of a one-parameter family of constant scalar curvature metrics for c and e and hence the a_i sufficiently large.

□

This existence proof fails when $b_i>0$ (or $b_i<0$) for all $1\leq i\leq m$. Indeed, there are known examples where there is no constant scalar curvature metric in which all of the b_i's have the same sign. The best known examples are the Hirzebruch surfaces which will be discussed below.

8.5.4 Constant Scalar Curvature Metrics on $\mathbb{C}P^{\frac{d_1+2}{2}}$-Bundles

In this section, we set

$$A_1(0) = 0$$

while continuing to set $A_i(x) > 0$ when $x \in [0,1]$ for all $2 \leq i \leq m$. This condition is equivalent to setting $a_1 = 0$. In this case, the manifold M is a $\mathbb{C}P^{\frac{d_1+2}{2}}$-bundle over $M_2 \times ... \times M_m$.

As in the extremal Kahler situation discussed above, we restrict to the case to the fiberwise Kahler manifolds of Construction Two in which each principal orbit is an S^1-bundle over $M_1 \times ... \times M_m$. For the metric to be extended smoothly over the special orbit at $x = 0$, we must have $M_1 \cong \mathbb{C}P^{\frac{d_1}{2}}$ and $b_1 = \frac{2}{d_1+2}$.

We have already seen that we retrieve the Fubini-Study metric on $M \cong \mathbb{C}P^{\frac{d_1+2}{2}}$ when $m = 1$. Therefore, we need only consider the case in which $m > 1$.

As above, we can without loss of generality assume that none of the b_i are zero. We distinguish between three general cases: 1. $b_i > 0$ for all $2 \leq i \leq m$; 2. $b_i < 0$ for all $2 \leq i \leq m$; and 3. for $2 \leq i \leq m$ some of the b_i are positive and some are negative. Obviously, in the third case $m \geq 3$. We consider the third case in the following theorem:

Theorem 8.12. *Let (M, g, J) be a fiberwise Kahler toric manifold obtained from Construction Two such that each principal orbit is an S^1-bundle over $M_1 \times ... \times M_m$ such that $M_1 \cong \mathbb{C}P^{\frac{d_1}{2}}$ and M is a $\mathbb{C}P^{\frac{d_1+2}{2}}$-bundle over $M_2 \times ... \times M_m$. If there exist $b_i > 0$ for some $2 \leq i \leq m$ and $b_j < 0$ for some $2 \leq j \leq m$, then M admits at least a 1-parameter family of constant positive scalar curvature Kahler metrics.*

Proof. By reordering if necessary, let $b_2, ..., b_l > 0$ and $b_{l+1}, ..., b_m > 0$ for some $3 \leq l \leq m$. Let $c_i = \frac{a_i}{|b_i|}$. We see that

$$\prod_{j=1}^{m}(b_j x + a_j)^{\frac{d_j}{2}} = \frac{2}{d_1+2}x^{\frac{d_1}{2}}\prod_{j=2}^{m}|b_j|^{\frac{d_j}{2}}\prod_{j=2}^{l}(x+c_j)^{\frac{d_j}{2}}\prod_{j=l+1}^{m}(c_j - x)^{\frac{d_j}{2}}$$

and

$$\sum_{j=1}^{m}\frac{d_j}{b_j x + a_j} = \frac{d_1(d_1+2)}{2x} + \sum_{j=2}^{l}\frac{\frac{d_j}{|b_j|}}{c_j + x} + \sum_{j=l+1}^{m}\frac{\frac{d_j}{|b_j|}}{c_j - x}.$$

To find our 1-parameter family of constant scalar curvature metrics, we make the following assumption to simplify calculations. Set

$$c_j = c > 0$$

for all $2 \leq j \leq l$ and set

$$c_i = e > 1$$

for all $l + 1 \leq i \leq m$.

This implies that

$$h = \frac{1}{x^k(x+c)^p(e-x)^q}\int\left[\int\left(\frac{2k(k+1)}{x}+\frac{r}{x+c}+\frac{t}{e-x}-\beta\right)x^k(x+c)^p(e-x)^q dx\right]dx$$

where $k=\frac{d_1}{2}$, $p=\sum_{j=2}^{l}$, $q=\sum_{j=l+1}^{m}$, $r=\sum_{j=2}^{l}\frac{d_j}{|b_j|}$, and $t=\sum_{j=l+1}^{m}\frac{d_j}{|b_j|}$ which are all greater than zero and independent of the a_i.

Set $h(x)=\frac{T(x)}{L(x)}$ where

$$T(x)=\int\left[\int\left(\frac{2k(k+1)}{x}+\frac{r}{x+c}+\frac{t}{e-x}-\beta\right)x^k(x+c)^p(e-x)^q dx\right]dx$$

and

$$L(x)=x^k(x+c)^p(e-x)^q.$$

Schematically, we can write T as

$$T=F_{k+p+q+2}x^{k+p+q+2}+...+F_w x^w+...+F_{k+2}x^{k+2}+2c^pe^qx^{k+1}+Ex+F$$

where E and F are the two constant of integration and the F_w are polynomials in c, e, r, t, p, q, and β. For the metric to be smooth at the special orbit $x=0$, we must have $h(0)=0$ and $h'(0)=2$. It is straightforward to see that these two conditions hold if and only if $E=F=0$. The two remaining conditions $h(1)=0$ and $h'(1)=-2$ determine β and place a condition on c and e.

It will be helpful to group the terms of T and L by the order of c and e. We see, after factoring out x^k from T and L, that

$$W=\frac{T(x)}{x^k}=[2c^pe^qx+\frac{c^{p-1}e^{q-1}}{(k+2)(k+1)}(re+tc)x^2+...]$$

$$-\beta[\frac{c^pe^q}{(k+2)(k+1)}x^2+\frac{c^{p-1}e^{q-1}}{(k+3)(k+2)}(pe-qc)x^3+...]$$

and

$$Z=\frac{L(x)}{x^k}=c^pe^q+c^{p-1}e^{q-1}(pe-qc)x+...$$

The condition $h(1)=0$ holds if and only if $W(1)=T(1)=0$. This conditions determines the value of β,

$$\beta=\frac{2c^pe^q+\frac{c^{p-1}e^{q-1}}{(k+2)(k+1)}(re+tc)+...}{\frac{c^pe^q}{(k+2)(k+1)}+\frac{c^{p-1}e^{q-1}}{(k+3)(k+2)}(pe-qc)+...}.$$

As $c\to\infty$ and $e\to\infty$, we see that $\beta\to 2(k+2)(k+1)$. The second condition as the special orbit $x=1$, $h'(1)=-2$, is equivalent to $\frac{W'(1)}{Z(1)}=-2$. Having found an expression for β, this condition becomes an equation in c and e. We note that

$$W'(1)=[2c^pe^q+\frac{2c^{p-1}e^{q-1}}{(k+2)(k+1)}(re+tc)+...]-\beta[\frac{2c^pe^q}{(k+2)(k+1)}+\frac{3c^{p-1}e^{q-1}}{(k+3)(k+2)}(pe-qc)+...]$$

and

$$Z(1)=c^pe^q+c^{p-1}e^{q-1}(pe-qc)+...$$

The condition $h'(1) = -2$ becomes

$$\beta[\frac{2c^p e^q}{(k+2)(k+1)} + \frac{3c^{p-1}e^{q-1}}{(k+3)(k+2)}(pe-qc)+...] = 4c^p e^q + 2c^{p-1}e^{q-1}(pe-qc) + \frac{2c^{p-1}e^{q-1}}{(k+2)(k+1)}(re+tc).$$

Combining this with our expression for β above implies that

$$\frac{4c^{2p-1}e^{2q-1}}{(k+3)(k+2)(k+1)}(pe-qc) + K(c,e) = 0 \tag{8.10}$$

where $K = K(c,e)$ is a polynomial is c and e in which each of the terms is of the form $K_{\eta\xi}c^\eta e^\xi$ such that $\eta \leq 2p$, $\xi \leq 2q$, $\eta + \xi < 2p + 2q - 1$, and the $K_{\eta\xi}$ are constants depending on the fixed values of r, t, p, and q.

Just as in our construction of constant scalar curvature metrics on $\mathbb{C}P^1$-bundles this equation can always be solved with e and c both arbitrarily large. We do not write the details here as they are almost identical to those of the proof of the last theorem in the previous section. We note that, as c and e approach infinity

$$h \sim 2x(1-x).$$

Therefore, for c and e arbitrarily large and satisfying (8.10), $h(x)$ will be positive for $x \in (0,1)$. This results gives a one-parameter family of constant scalar curvature metrics and completes the proof. □

8.5.5 Four-Dimensional Extremal Kahler Manifolds

On a four dimensional Kahler toric manifold, the number of summands, m, is equal to one and the dimension of that summand is $d_1 = d = 2$. Therefore, each principal orbit must be an S^1-bundle over $\mathbb{C}P^1$. To simplify the presentation, we set $b_1 = b$ and $a_1 = a$. From the definition of b, we see that

$$b = -\frac{q}{2}$$

where q is an integer and each value of q determines a distinct principal orbit type. In dimension four, the fiberwise Kahler toric metric can be written as

$$g = \begin{pmatrix} \frac{1}{h} & & \\ & h & \\ & & (bx+a)Id_2 \end{pmatrix}.$$

Because the manifold is compact, there must be two special orbits. Via translation in x, fix the first special orbit at $x = -1$. By rescaling, we can set the second special orbit at $x = 1$. For the metric to be positive on $[-1,1]$, $a > 0$ and $bx + a > 0$ when $x \in (-1,1)$.

From the discussion above, we notice two possibilities for the special orbits. Either both special orbits are copies of $\mathbb{C}P^1$ or one of the special orbits is $\mathbb{C}P^1$ and the other is a point. In the first case, $bx + a > 0$ on $x \in [-1,1]$ and in the second case, $a = b$ or $a = -b$. We will address these two case separately.

Case 1: One Special Orbit is a Point:

If the special orbit at $x = -1$ is a point, then $a = b$. The dimension of the sphere vanishing as that point is three: $k = 3$. The smoothness condition demands that $\frac{d}{b} = \frac{k^2-1}{2} = \frac{3^2-1}{2}$ which implies that $b = a = \frac{1}{2}$. Plugging this into the equation and integrating, we retrieve the Fubini-Study metric on $\mathbb{C}P^2$.

Case 2: Both Special Orbits are Copies of $\mathbb{C}P^1$:

When both the special orbits are two-spheres, the manifold M will be a Hirzebruch surface indexed by the integer q. Calabi found that each Hirzebruch surface admits, up to scaling, a one-parameter family of extremal Kahler metrics [6]. However, these metrics were not written down explicitly. In fact, Hwang and Simanca prove that within each of these one-parameter families is an extremal Kahler metric locally equivalent to an Hermitian-Einstein metric [31]. This correspondence, however, defines globally a smooth Hermitian-Einstein four manifold when $q = 0, 1$.

For this metric to be extremal Kahler, we have shown that the function h must satisfy the equation

$$h = \frac{1}{bx+a} \int \left[\int \left(\frac{2}{bx+a} - (\alpha x + \beta) \right) (bx+a) dx \right] dx \tag{8.11}$$

for $x \in (-1, 1)$. For the metric to be smooth at the special orbits $h(-1) = h(1) = 0$, $h'(-1) = 2$, and $h'(1) = -2$. Performing this simple double integral and imposing the boundary conditions we find that

$$\begin{aligned} \alpha &= \frac{6b(2a-1)}{3a^2-b^2}, \\ \beta &= \frac{6(a-b^2+a^2)}{3a^2-b^2}, \end{aligned} \tag{8.12}$$

and

$$h = \frac{(1-x)(1+x)((2a-1)b^2x^2 + 2(3a^2-b^2)bx + 6a^3 + b^2 - 4b^2a)}{2(3a^3-b^2)(bx+a)}. \tag{8.13}$$

For a sufficiently large, this function is positive for $x \in (-1, 1)$. Therefore, for each value of b (i.e. on each Hirzebruch surface), there exists a one-parameter family of extremal Kahler metrics.

It remains to determine when these extremal Kahler metrics are of constant scalar curvature. That is, when do we get smooth metrics with $\alpha = 0$? From the form of α in the extremal Kahler case, $\alpha = 0$ if and only if $b = 0$ or $a = \frac{1}{2}$ by equation (8.12). If $b = 0$ then the S^1 bundle over $\mathbb{C}P^1$ is trivial and the manifold $\mathbb{C}P^1 \times \mathbb{C}P^1$ with its product metric. Moreover, the only extremal Kahler metrics on $\mathbb{C}P^1 \times \mathbb{C}P^1$ are of constant scalar curvature. If $a = \frac{1}{2}$, then $|q| < \frac{1}{2}$. This follows from the fact that $a + b > 0$ and $a - b > 0$ for the metric to be positive at the special orbits. However, q

is an integer and this is satisfied if and only if $q = 0$. Therefore, b must equal zero and the manifold is a again a product.

Note: Because each $\mathbb{C}P^1$ in the fibers is S^1-invariant, these manifolds are Kahler toric manifolds whose polytopes are four-sided. In fact, these are the only Kahler toric four manifolds whose polytopes have four sides. Transforming the metric from fiberwise Kahler toric coordinates to Kahler toric coordinates is straightforward. The extremal Kahler metrics on the blow-up of $\mathbb{C}P^2$ at one point (the Hirzebruch surface with $q = 1$ or -1) was written by Abreu in Kahler toric coordinates [1].

8.5.6 Six-Dimensional Extremal Kahler Manifolds

While the classification of four-dimensional fiberwise Kahler toric manifolds admitting was already known, there remain open questions in dimension six. Here we will classify all six-dimensional fiber Kahler toric manifolds admitting extremal Kahler metrics and write those metrics down explicitly in terms of rational functions. In dimension six, each principal orbit is an S^1-bundle over a four-dimensional Kahler-Einstein manifold. The only four manifolds admitting a Kahler-Einstein metric with positive Einstein constant are $\mathbb{C}P^2$, $\mathbb{C}P^1 \times \mathbb{C}P^1$, or the blow-up of $\mathbb{C}P^2$ at k distinct points where $3 \leq k \leq 8$ such that no three points lie on a line and no six points lie on a conic [61].

When the base manifold of the S^1-bundle is any manifold other than $\mathbb{C}P^1 \times \mathbb{C}P^1$, the metric can be written as

$$g = \begin{pmatrix} \frac{1}{h} & & \\ & h & \\ & & (bx+a)Id_4 \end{pmatrix}$$

since the Kahler-Einstein metrics on $\mathbb{C}P^2$ and the blow-ups are irreducible.

On the other hand, if each principal orbit is an S^1-bundle over $\mathbb{C}P^1 \times \mathbb{C}P^1$, the metric looks like

$$g = \begin{pmatrix} \frac{1}{h} & & & \\ & h & & \\ & & (b_1x+a_1)Id_2 & \\ & & & (b_2x+a_2)Id_2 \end{pmatrix}.$$

To the author's knowledge, it had not been determined before whether or not manifolds of this type admit extremal Kahler metrics.

For ease of presentation, we set $b_1 = b$, $a_1 = a$, $b_2 = d$, and $a_2 = c$. The values of b and d determine the principal orbit type. As each principal orbit is an S^1 bundle over $\mathbb{C}P^1 \times \mathbb{C}P^1$, $b = -\frac{p}{2}$ and $d = -\frac{q}{2}$ where p and q are integers [15]. Let $P(p, q)$ denote the principal orbit type. Note that $P(p, q) \cong P(-p, -q)$.

Again, by translation in x and rescaling of the metric, we can set the location of the two special orbits to be $x = -1$ and $x = 1$. First, we assume that both the special orbits are $S^2 \times S^2$, i.e.

$$M = \mathbb{P}(\mathcal{O}_{\mathbb{C}P^1 \times \mathbb{C}P^1} \oplus \mathcal{O}_{\mathbb{C}P^1 \times \mathbb{C}P^1}(p, q))$$

a $\mathbb{C}P^1$-bundle over $\mathbb{C}P^1 \times \mathbb{C}P^1$.

The metric is positive if and only if $a > 0$, $c > 0$, $|\frac{a}{b}| > 1$, $|\frac{c}{d}| > 1$, and $h > 0$ when $x \in (-1, 1)$. Finally, smoothness at the special orbits requires that $h(-1) = h(1) = 0$, $h'(-1) = 2$, and $h'(1) = -2$.

In the extremal Kahler case, h satisfies (8.6) with $S = \alpha x + \beta$. Integrating this equation and imposing smoothness conditions we find that

$h = \frac{1}{2}(1-x)(x+1)((6a^2d^2bc + 6db^2ac^2 - 3ba^2d^2 - 3b^2c^2d + b^3d^2 + b^2d^3)x^3 + (3cb^2d^2 + 3ab^2d^2 + 25dba^2c^2 - 5a^3d^2 + 10c^3b^2a + 10a^3d^2c - 5a^2d^3b - 5c^2b^3d - 5c^3b^2 + 3d^3b^3 - 2d^2b^2ac)x^2 + (-8a^2d^2bc - 8db^2ac^2 + 30a^2c^3b - b^3d^2 + 6cd^2b^3 - 10c^3b^3 - 10a^3d^3 - b^2d^3 + 3ba^2d^2 + 3b^2c^2d + 6ad^3b^2 + 30a^3c^2d)x - 5c^2b^3d + 13dba^2c^2 - 5a^2d^3b + 5a^3d^2 + 5c^3b^2 + 30a^3c^3 + 3d^3b^3 - 20a^3d^2c - 20c^3b^2a + 16d^2b^2ac - 3ab^2d^2 - 3cb^2d^2)/((-5c^2b^2 + 15a^2c^2 + 4dbac - 5a^2d^2 + 3d^2b^2)(bx + a)(dx + c))$.

The constants α and β in the scalar curvature satisfy

$$\alpha = \frac{10(6a^2cd + 6abc^2 + bd^2 + db^2 - 3c^2b - 3a^2d)}{-5c^2b^2 + 15a^2c^2 + 4dbac - 5a^2d^2 + 3d^2b^2} \tag{8.14}$$

and

$$\beta = \frac{2(15c^2a - 5cb^2 + 4dbc + 15a^2c - 5ad^2 + 4bad - 6dbac + 15a^2c^2 - 15c^2b^2 - 15a^2d^2 + 9d^2b^2)}{-5c^2b^2 + 15a^2c^2 + 4dbac - 5a^2d^2 + 3d^2b^2}.$$

All of the smoothness conditions are satisfied for all values of a, b, c, and d. However, the function h is not necessarily positive for all values of these variable. It is positive for a and c sufficiently large. Therefore, for every choice of b and d (that is for every principal orbit type) there exists, up to scale, a two parameter family of extremal Kahler metrics.

Theorem 8.13. *Let M be $\mathbb{P}(\mathcal{O}_{\mathbb{C}P^1\times\mathbb{C}P^1} \oplus \mathcal{O}_{\mathbb{C}P^1\times\mathbb{C}P^1}(p,q))$ endowed with a cohomogeneity-one fiberwise Kahler toric metric as described above. Every such manifold admits, up to scaling, a two parameter family of extremal Kahler metrics.*

It remains to determine which of these manifolds admit metrics of constant scalar curvature. To determine this, we need only determine when $\alpha = 0$ while the smoothness and positiveness conditions are satisfied.

Theorem 8.14. *Let $M = \mathbb{P}(\mathcal{O}_{\mathbb{C}P^1\times\mathbb{C}P^1} \oplus \mathcal{O}_{\mathbb{C}P^1\times\mathbb{C}P^1}(p,q))$, with p and q as above. For $(p,q) = (0,0)$, the manifold is the product of three copies of $\mathbb{C}P^1$ and admits, up to scale, a two parameter family of constant scalar curvature metrics. If $p > 0$ and $q < 0$, then the manifold admits at least a one-parameter family of constant scalar curvature metrics within its two-parameter family of extremal Kahler metrics. For $p \geq 0$ and $q > 0$, the manifold admits no constant scalar curvature Kahler metrics.*

Proof. Recall that $b = -\frac{p}{2}$ and $d = -\frac{q}{2}$ for $p, q \in \mathbb{Z}$ and that $a > \frac{|p|}{2}$ and $c > \frac{|q|}{2}$. The metric has constant scalar curvature if $\alpha = 0$ which is equivalent to

$$3a^2d(2c-1) + 3c^2b(2a-1) + bd(b+d) = 0.$$

If $b = d = 0$, then $\alpha = 0$ automatically and there is a two-parameter family of extremal Kahler metrics. On the other hand if $b \geq 0$ and $d > 0$ then $\alpha \neq 0$ because a and c are both greater then $\frac{1}{2}$.

Finally, if $b > 0$ and $d < 0$, then we can solve for c in terms of a. The metric has constant scalar curvature if

$$c = \frac{-6a^2d \pm \sqrt{9a^4d^2 + (3b^3d + 3b^2d^2 - 9bda^2)(1-2a)}}{3b(2a-1)}. \tag{8.15}$$

Consider the first of these solutions (taking '+') as $a \to \infty$, $c \to \infty$. Therefore, for sufficiently large a we are certain to have a constant scalar curvature metric with h positive on $(-1, 1)$. This gives the one-parameter family of constant scalar curvature metrics.

□

Note that this theorem determines which of these *manifolds* admit a constant scalar curvature metric. It further demonstrates that those manifolds which do admit a constant scalar curvature Kahler metric in fact admit at least a one-parameter family. They may admit more. In fact, $\mathbb{C}P^1 \times \mathbb{C}P^1 \times \mathbb{C}P^1$ trivially admits a full two-parameter family of such metrics. What is more interesting is that $\mathbb{P}(\mathcal{O}_{\mathbb{C}P^1 \times \mathbb{C}P^1} \oplus \mathcal{O}_{\mathbb{C}P^1 \times \mathbb{C}P^1}(1, -1))$, as demonstrated in a previous section, admits *two* one-parameter families of constant scalar curvature metrics. This occurs because both solutions for the constant c in terms of the constant a can be solved without violating the smoothness and positive definiteness conditions.

Proposition 8.15. *For $M = \mathbb{P}(\mathcal{O}_{\mathbb{C}P^1 \times \mathbb{C}P^1} \oplus \mathcal{O}_{\mathbb{C}P^1 \times \mathbb{C}P^1}(p, q))$, M admits, up to scale, more than a one-parameter family of constant scalar curvature Kahler metrics if and only if $(p, q) = (0, 0)$ or $(1, -1)$.*

Proof. The proof consists of demonstrating that the '$-$' solution of (8.15) only gives a valid solution for those two values of p and q. This is trivially done but awkward to present. We omit the details. □

The existence of Kahler-Einstein metrics on manifolds of this type was already answered by Sakane.

Theorem 8.16. *[54] For $M = \mathbb{P}(\mathcal{O}_{\mathbb{C}P^1 \times \mathbb{C}P^1} \oplus \mathcal{O}_{\mathbb{C}P^1 \times \mathbb{C}P^1}(p, q))$, M admits a Kahler-Einstein metric if and only if $(p, q) = (0, 0)$ or $(1, -1)$.*

We note that the manifolds admitting Kahler-Einstein metrics are precisely those that admit more than a one-parameter of constant scalar curvature metrics. It would be interesting to study whether there is a connection between the existence of Kahler-Einstein metrics and the 'number' of constant scalar curvature Kahler metrics.

Chapter 9

Cohomogeneity-d Fiberwise Kahler Toric Manifolds

In the cohomogeneity-one case, the Einstein and extremal Kahler conditions were seen to be second-order ordinary differential equations of one function in one variable, h, which we could integrate explicitly. The Einstein and extremal Kahler conditions in the cohomogeneity-d case are partial differential equations and we must therefore use a slightly different strategy of integration. Our integration procedure is similar to that employed to integrate the Einstein equations in the Kahler toric case.

9.1 Explicit Integration of the Einstein Condition

Above, we calculate that the Einstein condition on a fiberwise Kahler toric metric is equivalent to equations (7.6) and (7.7). We begin our integration of the Einstein equations by equation (7.6):

$$h_{ik}\frac{\partial}{\partial x^k}\left(h_{jl}\frac{\partial \log(\det(h)V^{\frac{1}{2}})}{\partial x^l}\right) = -2\lambda h_{ij}.$$

Multiplying both sides of this equation by h^{im} we find that

$$\frac{\partial}{\partial x^k}\left(h_{lj}\frac{\partial \log(\det(h)V^{\frac{1}{2}})}{\partial x^l}\right) = -2\lambda \delta^k_j.$$

Integrating these equations we find that

$$h_{lj}\frac{\partial \log(\det(h)V^{\frac{1}{2}})}{\partial x^l} = -2\lambda x_j + C_j \tag{9.1}$$

for all $1 \leq j \leq d$ where the C_j are constants. We can integrate these equations by first multiplying both sides of equation (9.1) by h^{mj} to obtain

$$\frac{\partial \log(\det(h)V^{\frac{1}{2}})}{\partial x^l} = h^{lj}(-2\lambda x_j + C_j).$$

Recall that $h^{lj} = \frac{\partial^2 \Phi}{\partial x^l \partial x^j}$ and therefore

$$\frac{\partial \log(\det(h)V^{\frac{1}{2}})}{\partial x^l} = (-2\lambda x_j + C_j)\frac{\partial^2 \Phi}{\partial x^l \partial x^j}.$$

These equations integrate to

$$\log(\det(h)V^{\frac{1}{2}}) = \sum_{j=1}^{d}(-2\lambda x_j + C_j)\frac{\partial \Phi}{\partial x^j} + 2\lambda\Phi + E$$

where E is a constant.

In light of equation (9.1), equation (7.7) of the Einstein condition becomes

$$\left(h_{kl}\frac{\partial \log(\det(h)V^{\frac{1}{2}})}{\partial x^l} + 2\lambda x_k\right) b_{ki} = \sum_{k=1}^{d} C_k b_{ki} = 2(1-\lambda a_i)$$

for all $1 \leq i \leq m$. These results prove the following theorem.

Theorem 9.1. *Let (M^n, g, J) be a fiberwise Kahler toric manifold of cohomogeneity-d. The Einstein condition is equivalent to equation*

$$\log(\det(h)V^{\frac{1}{2}}) = \sum_{j=1}^{d}(-2\lambda x_j + C_j)\frac{\partial \Phi}{\partial x^j} + 2\lambda\Phi + E \tag{9.2}$$

and the relation

$$\begin{pmatrix} b_{11} & \cdot & \cdot & \cdot & b_{d1} \\ \cdot & \cdot & & & \cdot \\ \cdot & & \cdot & & \cdot \\ \cdot & & & \cdot & \cdot \\ b_{1m} & \cdot & \cdot & \cdot & b_{dm} \end{pmatrix} \begin{pmatrix} C_1 \\ \cdot \\ \cdot \\ \cdot \\ C_d \end{pmatrix} = \begin{pmatrix} 2(1-\lambda a_1) \\ \cdot \\ \cdot \\ \cdot \\ 2(1-\lambda a_m) \end{pmatrix} \tag{9.3}$$

where the a_r, the b_{jr}, the C_j and E are constants and V, Φ, and deth *are as defined above.*

Expressed in complex coordinates, this theorem becomes the following.

Theorem 9.2. *Let (M, g, J) be a fiberwise Kahler toric manifold with complex coordinates $\eta = \eta_{u_1,\ldots,u_d}$ as above. The Einstein condition is equivalent to the equation*

$$\log(\det(h)V^{\frac{1}{2}}) = -2\lambda\eta + \sum_{j=1}^{d} C_j u_j + E$$

and the relation

$$\begin{pmatrix} b_{11} & \cdot & \cdot & \cdot & b_{d1} \\ \cdot & \cdot & & & \cdot \\ \cdot & & \cdot & & \cdot \\ \cdot & & & \cdot & \cdot \\ b_{1m} & \cdot & \cdot & \cdot & b_{dm} \end{pmatrix} \begin{pmatrix} C_1 \\ \cdot \\ \cdot \\ \cdot \\ C_d \end{pmatrix} = \begin{pmatrix} 2(1-\lambda a_1) \\ \cdot \\ \cdot \\ \cdot \\ 2(1-\lambda a_m) \end{pmatrix}. \tag{9.4}$$

In complex coordinates, the functions A_i satisfy

$$A_i = \sum_{j=1}^{d} b_{ji}\frac{\partial \eta}{\partial u_j} + a_i.$$

From the Einstein condition, we see that much of the structure of the induced Kahler toric metric $\bar{g}$ determines much of the structure of the total fiberwise Kahler toric metric in the Einstein case. For example, we have the following theorem.

Theorem 9.3. *Let (M, g) be a cohomogeneity-d fiberwise Kahler toric metric such that g is Kahler-Einstein. If, locally, $(N, \bar{g})$ can be written as a product $(N_1 \times N_2, \bar{g}_1 \oplus \bar{g}_2)$ of two Kahler toric metrics where $\mathrm{dimN}_1 = 2\mathrm{l}_1$ and $\mathrm{dimN}_2 = 2\mathrm{l}_2$ then (M, g) splits as $(M_1 \times M_2, g_1 \oplus g_2)$ where M_1 and M_2 are cohomogeneity-l_1 and codimension-l_2 fiberwise Kahler toric manifolds.*

Proof. If $(N, \bar{g})$ can be written as a product metric then the potential $\Phi(x_1, ..., x_d)$ can be written as $\Phi = \Phi_1(x_1, ..., x_{l_1}) + \Phi_2(x_{l_1+1}, ..., x_d)$ (here we might have transformed by $SL(d, \mathbb{Z})$ and reordered the coordinates). The right-hand side of equation (9.2) splits as the sum of two functions: one in the variables $x_1, ..., x_{l_1}$ and one in the coordinates $x_{l_1}, ..., x_d$. On the left-hand side, the $\log(\det(h))$ term similarly splits. Therefore, the $\log(V^{\frac{1}{2}})$ term must split in the same way which occurs if and only if the total manifold is a product.

□

9.2 The Futaki Invariant on Fiberwise Kahler Toric Manifolds

Above, we reviewed the known holomorphic obstructions to the existence of Kahler-Einstein metrics with positive Einstein constant. More specifically, we demonstrated how to calculate the Futaki invariant on Kahler toric manifolds with positive first Chern class in terms of the symplectic coordinates. In this section we want to calculate the Futaki invariant on fiberwise Kahler toric manifolds in terms of the coordinates introduced above . We will see that the formulae we obtain are very similar to those obtained in the Kahler toric case.

Let (M^{2n}, g, J) be a compact fiberwise Kahler toric manifold with positive first Chern class: $c_1(M) > 0$. If a manifold of this type admits a Kahler-Einstein metric then the Futaki invariant must be zero. Recall that for $X \in \mathfrak{h}(M)$ a holomorphic vector field, the Futaki invariant satisfies

$$\mathcal{F}(X) = -\int_N \mu_X \omega^n.$$

To calculate the Futaki invariant on a fiberwise Kahler toric manifold, consider the (real) holomorphic vector fields $\frac{\partial}{\partial u^i}$. As shown above, we know that $\mu_{\frac{\partial}{\partial u^i}} = \frac{\partial \eta}{\partial u^i} = x_i$. Furthermore, given the form of the metric of a fiberwise Kahler toric manifold, we see that

$$\omega^n = \prod_{j=1}^{m} A_j^{\frac{d_j}{2}} dx_1 \wedge .. \wedge dx^d \wedge ... \wedge d\phi_1 \wedge ... \wedge d\phi_d \wedge d\mathrm{vol}^*$$

where $d\mathrm{vol}^*$ is the volume form of the metric g^* associated to either the coadjoint orbit G/L or the product of Kahler-Einstein manifolds $M_1 \times ... \times M_m$. If vol^* is the volume associated $d\mathrm{vol}^*$ then we calculate that

$$\mathcal{F}\left(\frac{\partial}{\partial u^i}\right) = -\int_M x_i \omega^n = -(\mathrm{vol}^*)(2\pi)^d \int_{\triangle} x_i \prod_{j=1}^{m} A_j^{\frac{d_j}{2}} dx$$

for all i where $\triangle$ is the polytope defined in symplectic coordinates.

We have the following proposition.

Proposition 9.4. *If (M, g, J) is a compact fiberwise Kahler toric manifold with a Kahler-Einstein metric then*

$$\int_{\triangle} x_i V^{\frac{1}{2}} dx = 0$$

where $V^{\frac{1}{2}} = \prod_{j=1}^{m} A_j^{\frac{d_j}{2}} = \prod_{j=1}^{m} (\sum_{r=1}^{d} b_{rj} x_r + a_j)^{\frac{d_j}{2}}$.

In the case when $d = 1$, Koiso and Sakane proved that this condition was in fact sufficient. It would be interesting to determine whether the vanishing of the Futaki invariant is sufficient for the existence of a Kahler-Einstein metric for higher values of d.

Chapter 10

Hermitian Metrics of Constant Scalar Curvature

In our study of fiberwise Kahler toric manifolds with $d = 1$, we were able to integrate the constant scalar curvature equation and express the resulting metrics explicitly in terms of rational functions. We will now perform this analysis on manifolds of a very similar structure which are Hermitian and non-Kahler. Manifolds of this type were first studied by Wang and Wang in [57] who provide solutions to the Einstein condition on such manifolds. After reviewing their construction of these manifolds, we will calculate the scalar curvature equation.

We begin, as we did in constructing fiberwise Kahler toric manifolds, by letting (M_j, g_j^*) be Kahler-Einstein manifolds of real dimension d_j with positive Einstein constant for $1 \leq j \leq m$. Let P be a principal S^1-bundle over the product manifold $M_1 \times ... \times M_m$. The first Chern class can be written as $c_1(M_j) = p_j \alpha_j$ where p_j is a positive integer and α_j is an indivisible class in $H^2(M_j, \mathbb{Z})$ [38]. Let $\chi = \sum_{j=1}^m b_j \pi_j^* \alpha_j$, where π_j is the projection from P onto the manifold M_j, be the Euler class which determines the principal S^1-bundle as in [58]. Finally, let θ be the connection of this S^1-bundle whose curvature represents the Euler class.

We define a metric g on $P \times I$ where I is some open interval in $\mathbb{R}$ by the equation

$$g = dt^2 + h(t)\theta \otimes \theta + \sum_{j=1}^{m} A_j(t) \pi_j^* g_j^*$$

where h and the A_j are positive and smooth functions on I. Let M be the manifold that we will construct by extending g over special orbits.

We perform a change of variable by letting $dx = \sqrt{h} dt$. Under this transformation, the metric becomes

$$g = \frac{1}{h} dx^2 + h(x)\theta \otimes \theta + \sum_{j=1}^{m} A_j(x) \pi_j^* g_j^*.$$

This metric contains, trivially, a two-dimensional Kahler toric metric $\bar{g} = \frac{1}{h} dx^2 + h(t)\theta \otimes \theta$ with complex structure $\bar{J}$ which satisfies $\bar{J}(\partial_\phi) = h^2 \partial_x$ where ∂_ϕ is the dual vector field to the connection

θ. By combining $\bar{J}$ and the lifts of the complex operators J_i in the natural way, the manifold (M, g) can be endowed with a complex structure J.

In these variables the Einstein condition becomes (see M. Wang's report in [41]) the following system of equations

$$-\frac{1}{2}h'' - h'\left(\sum_{j=1}^{m}\frac{d_j}{4}\frac{A_j'}{A_j}\right) + h\left[\sum_{j=1}^{m}\frac{d_j}{4}\left(\frac{A_j'}{A_j}\right)^2 - \sum_{j=1}^{m}\frac{d_j}{2}\frac{A_j''}{A_j}\right] = \lambda \tag{10.1}$$

$$-\frac{1}{2}h'' - h'\left(\sum_{j=1}^{m}\frac{d_j}{4}\frac{A_j'}{A_j}\right) + h\left(\sum_{j=1}^{m}\frac{d_j}{4}\frac{b_j^2}{A_j^2}\right) = \lambda \tag{10.2}$$

and

$$-\frac{1}{2}h'\frac{A_i'}{A_i} + h\left[-\frac{1}{2}\frac{A_i''}{A_i} + \frac{1}{2}\left(\frac{A_i'}{A_i}\right)^2 - \frac{A_i'}{A_i}\left(\sum_{j=1}^{m}\frac{d_j}{4}\frac{A_j'}{A_j}\right) - \frac{1}{2}\frac{b_i^2}{A_i^2}\right] + \frac{1}{A_i} = \lambda \tag{10.3}$$

for all $1 \leq i \leq m$. (Note that $f' = \frac{df}{dx}$ for all functions $f(x)$.)

Before proceeding to the question of finding constant scalar curvature metrics we first review some of the results found in [57]. By subtracting equation (10.1) from (10.2) we find that

$$\sum_{j=1}^{m} d_j\left[2\frac{A_j''}{A_j} - \left(\frac{A_j'}{A_j}\right)^2 + \frac{b_j^2}{A_j^2}\right] = 0. \tag{10.4}$$

Following Wang and Wang we set

$$\mu_j = \left[2\frac{A_j''}{A_j} - \left(\frac{A_j'}{A_j}\right)^2 + \frac{b_j^2}{A_j^2}\right]$$

Although unable to solve the Einstein condition in full generality, Wang and Wang were able to obtain explicit solutions when $\mu_j = 0$ for all $1 \leq i \leq m$. They were able to assign geometric meaning to this condition. They say that demanding that the $\mu_j = 0$ for all j was equivalent to demanding that the Riemannian curvature tensor, R, exhibit a particular kind of J invariance.

Assumption:

$$R(X, Y, Z, W) = R(JX, JY, JZ, JW)$$

for all X, Y, Z, and W in TM where R is the curvature tensor of g. Hermitian metrics of this type were studied in [27] and [20].

Let h' denote differentiation with respect to the new coordinate x. We have the following proposition.

Proposition 10.1. *[57] The above assumption is equivalent to the condition that*

$$\mu_j = 0 \tag{10.5}$$

for all $1 \leq j \leq m$.

Proof. The proof is a straightforward calculation. We refer the reader to [57] which gives this equation in slightly different coordinates. □

Equation (10.5) admits two types of solutions. Namely,

$$A_j = b_j x + a_j \tag{10.6}$$

or

$$A_j = e_j(x + c_j)^2 - \frac{1}{4}\frac{b_j^2}{e_j} \tag{10.7}$$

where a_j, c_j, and e_j are the constants of integration.

Note that solution (10.6) is equivalent to the Kahler condition if it holds for all j. If, however, solution (10.6) does not hold for some j, the manifold will be Hermitian and non-Kahler. We will not discuss their work further except to note that in the Kahler case, the resulting metric is fiberwise Kahler toric.

We now return to the question of integrating the constant scalar curvature equation on manifolds of this type. From equations (10.1)-(10.3), it is straightforward to retrieve the equation for the scalar curvature, S, of this kind of metric. We have

$$S = -h'' - h'\left(\sum_{j=1}^{m} d_j \frac{A_j'}{A_j}\right) \tag{10.8}$$

$$+h\left[-\sum_{j=1}^{m} d_j \frac{A_j''}{A_j} + \frac{3}{4}\sum_{j=1}^{m} d_j \left(\frac{A_j'}{A_j}\right)^2 - \frac{1}{4}\left(\sum_{j=1}^{m} d_j \frac{A_j'}{A_j}\right)^2 - \sum_{j=1}^{m} \frac{d_j}{4}\frac{b_j^2}{A_j^2}\right] + \sum_{j=1}^{m} \frac{d_j}{A_j}.$$

Although this equation is only second-order in h we are unable to integrate it in the general case. However, we can integrate this equation explicitly for a large subclass. To see how this is done, let us recall our method of integration in the Kahler case. The Kahler condition requires that $A_j = b_j x + a_j$ for all $1 \leq j \leq m$. We were able to integrate this equation to find that

$$h = \frac{1}{\prod_{j=1}^{m}(b_j x + a_j)^{\frac{d_j}{2}}} \int \left[\int \left(\sum_{j=1}^{m} \frac{d_j}{b_j x + a_j} - S\right) \prod_{j=1}^{m} (b_j x + a_j)^{\frac{d_j}{2}} dx\right] dx.$$

This suggests a possible form for h in the general case. We ask when h is of this form with A_i a general function instead of $b_i x + a_i$.

The answer is provided by the following theorem.

Theorem 10.2. *The scalar curvature equation (10.8) is satisfied for*

$$h = \frac{1}{\prod_{j=1}^{m} A_j^{\frac{d_j}{2}}} \int \left[\int \left(\sum_{j=1}^{m} \frac{d_j}{A_j} - S\right) \prod_{j=1}^{m} A_j^{\frac{d_j}{2}} dx\right] dx. \tag{10.9}$$

if and only if equation (10.4),

$$\sum_{j=1}^{m} d_j \mu_j \equiv \sum_{j=1}^{m} d_j \left[2\frac{A_j''}{A_j} - \left(\frac{A_j'}{A_j}\right)^2 + \frac{b_j^2}{A_j^2} \right] = 0,$$

holds.

Proof. First we calculate the first and second derivatives of h. They are

$$h' = \left(-\sum_{j=1}^{m} \frac{d_j}{2} \frac{A_j'}{A_j} \right) h - \frac{1}{\prod_{j=1}^{m} A_j^{\frac{d_j}{2}}} \int \left(\sum_{j=1}^{m} \frac{d_j}{A_j} - S \right) \prod_{j=1}^{m} A_j^{\frac{d_j}{2}} dx$$

and

$$h'' = h' \left(-\sum_{j=1}^{m} \frac{d_j}{2} \frac{A_j'}{A_j} \right) + h \left[\sum_{j=1}^{m} \frac{d_j}{2} \left(\frac{A_j'}{A_j}\right)^2 - \sum_{j=1}^{m} \frac{d_j}{2} \frac{A_j''}{A_j} \right]$$
$$+ \left(\sum_{j=1}^{m} \frac{d_j}{2} \frac{A_j'}{A_j} \right) \left[\frac{1}{\prod_{j=1}^{m} A_j^{\frac{d_j}{2}}} \int \left(\sum_{j=1}^{m} \frac{d_j}{A_j} - S \right) \prod_{j=1}^{m} A_j^{\frac{d_j}{2}} dx \right] + S - \sum_{j=1}^{m} \frac{d_j}{A_j}.$$

Substituting these equations into equation (10.8) gives equation (10.4). This completes the proof. □

While equation (10.9) holds in the Einstein case, it also holds more generally.

One may ask what the geometric significance of equation (10.4) is. We demonstrated above that the manifold can be endowed naturally with a complex structure J. Equation (10.1) of the Einstein condition is equivalent to the equation

$$r\left(\frac{\partial}{\partial x}, \frac{\partial}{\partial x}\right) = \lambda g\left(\frac{\partial}{\partial x}, \frac{\partial}{\partial x}\right)$$

and equation (10.2) is equivalent to

$$r\left(\frac{\partial}{\partial \phi}, \frac{\partial}{\partial \phi}\right) = \lambda g\left(\frac{\partial}{\partial \phi}, \frac{\partial}{\partial \phi}\right).$$

Equation (10.4) states that these two equations are equivalent. Since the metric is already J-invariant, equation (10.4) is equivalent to the condition

$$r\left(J\frac{\partial}{\partial \phi}, J\frac{\partial}{\partial \phi}\right) = r\left(\frac{\partial}{\partial \phi}, \frac{\partial}{\partial \phi}\right). \tag{10.10}$$

Note that if (10.10) holds, then the Ricci tensor is completely J-invariant. The preceding theorem states that when the Ricci tensor is J-invariant, the scalar curvature equation can be integrated to (10.9).

10.1 Constant Scalar Curvature Hermitian Metrics with J-Invariant Ricci Tensor

In the previous section, we were able to integrate the scalar curvature equation when the Ricci tensor of the Hermitian manifold was J-invariant. Before using this integration to construct new non-Kahler Hermitian metrics with J-invariant Ricci tensor and constant scalar curvature, we wish to discuss briefly why we believe this condition to be both natural and of interest, particularly in dimension four. To that end, we will try to motivate the condition as a natural generalization of both the Hermitian-Einstein and constant scalar curvature Kahler conditions.

Let (M, g, J) be a compact Hermitian manifold which is not necessarily Kahler. By the definition of an Hermitian manifold, the metric, g, is automatically J-invariant,

$$g(J\cdot, J\cdot) = g(\cdot, \cdot).$$

The Ricci tensor, r, is *not* automatically J-invariant. It is, however, invariant if either of two important conditions hold. If the metric is Kahler (i.e. for $\omega(\cdot, \cdot) = g(J\cdot, \cdot)$, $d\omega = 0$), it is well-known that the Ricci tensor is automatically J-invariant:

$$\text{Kahler} \Rightarrow r(J\cdot, J\cdot) = r(\cdot, \cdot).$$

If the metric, g, is Hermitian-Einstein, then $r(\cdot, \cdot) = \lambda g(\cdot, \cdot)$ where λ is a constant. Since g is J-invariant, the Einstein condition implies that the Ricci tensor is also J-invariant,

$$\text{Einstein} \Rightarrow r(J\cdot, J\cdot) = r(\cdot, \cdot).$$

Hermitian manifolds with J-invariant Ricci tensor provide a natural generalization of both the Einstein and the Kahler conditions. However, Kahler metrics, and by extension Hermitian metrics with J-invariant Ricci tensor, are very plentiful (though not every complex manifold admits a Kahler metric). We would like to place an additional condition on Hermitian metrics with J-invariant Ricci tensor. We propose that an interesting additional condition is to demand that the metrics be of constant scalar curvature.

Hermitian metrics of constant scalar curvature with J-invariant Ricci tensor include both Hermitian-Einstein metrics (Einstein metrics are automatically of constant scalar curvature) and constant scalar curvature Kahler metrics, both of which are of great interest to differential geometers. Manifolds of this type are also a subclass of the space of Hermitian manifolds with constant scalar curvature. While the classification of these types of manifolds is an interesting problem in all dimensions, it is particularly intriguing in dimension four.

In dimension four, much is known about constant scalar curvature Kahler metrics and Hermitian-Einstein metrics. The compact four-manifolds admitting Kahler-Einstein metrics have been completely classified by the work of Yau, Tian, and others (see LeBrun's article in [41] for a discussion).

Many open problems still exist in the classification of constant scalar curvature Kahler metrics. Much work has been done on this topic by Tian, Donaldson, and others relating the existence of constant scalar curvature metrics to various notions of stability, but no classification as yet exists in dimension four or in higher dimensions. More is known about the existence of non-Kahler Hermitian-Einstein metrics. As mentioned above LeBrun demonstrated that, in addition to the manifold $\mathbb{C}P^2\sharp\overline{\mathbb{C}P^2}$ which is known to admit a non-Kahler Hermitian-Einstein metric, the only other compact four manifolds which *could* admit a non-Kahler Hermitian-Einstein metric are $\mathbb{C}P^2\sharp 2\overline{\mathbb{C}P^2}$ and $\mathbb{C}P^2\sharp 3\overline{\mathbb{C}P^2}$.

We now turn our attention to the classification of constant scalar curvature Hermitian metrics with J-invariant Ricci tensor in dimension four. As yet no classification exists; in fact, to the author's knowledge this particular condition has not been studied. However, some work has been done on Hermitian metrics with J-invariant Ricci tensor on complex surfaces. For example, see [3]. We now review some of that work, beginning with the following theorem due to Apostolov and Gauduchon:

Theorem 10.3. *[4] Suppose (M, g, J) is a compact Hermitian four manifold with J-invariant Ricci tensor. The metric g is locally conformal to a Kahler metric. It is globally conformal to a Kahler metric $\bar{g} = f^2 g$ if and only if the first Betti number $b_1(M)$ is even. In that case, the vector field $J(\nabla_g f)$ is a Killing vector field of both g and $\bar{g}$.*

In the case of b_1 even, finding an Hermitian metric with J-invariant Ricci tensor is equivalent to finding a Kahler metric with a Killing vector field. In the case of b_1 odd, there are no known examples in four dimensions of a non-Kahler Hermitian metric with J-invariant Ricci tensor (Apostolov, personal communication).

To the author's knowledge, other than the Page metric, no non-Kahler Hermitian metrics of constant scalar curvature with J-invariant Ricci tensor on a compact four manifold have been constructed previously. In the following section, we construct a one-parameter family of such metrics on $\mathbb{C}P^1 \times \mathbb{C}P^1$ and $\mathbb{C}P^2\sharp\overline{\mathbb{C}P^2}$. It would be interesting to classify all of the metrics of this type in dimension four. In addition, one could ask whether *all* constant scalar curvature Hermitian metrics *must* have J-invariant Ricci tensor. Unfortunately, we are unable to give an answer to either of these questions.

10.2 New Constant Scalar Curvature Hermitian Metrics in Dimension Four

Consider the case in which $m = 1$ and $d_1 = 2$. That is, each principal orbit is an S^1-bundle over $\mathbb{C}P^1$ endowed with its canonical Kahler-Einstein metric. In this case,

$$g = \begin{pmatrix} \frac{1}{h} & & \\ & h & \\ & & A_1 Id_2 \end{pmatrix}.$$

The Kahler case was discussed above in the section on four-dimensional extremal Kahler metrics. In that section, we showed that, setting $b_1 = b$,

$$b = -\frac{q}{2}$$

where q is an integer and each value of q determines a distinct principal orbit type. Up to a change in orientation q and $-q$ give the same principal orbit type. Without loss of generality, we can assume that $b \geq 0$ and $q \leq 0$.

If the metric has two special orbits, which via translation and rescaling we can fix at $x = -1$ and $x = 1$, and that $A_1 > 0$ for $x \in [-1, 1]$ then the resulting manifold will be a Hirzebruch surface

$$\mathbb{P}(\mathcal{O}_{\mathbb{C}P^1} \oplus \mathcal{O}_{\mathbb{C}P^1}(q)).$$

We want to find constant scalar curvature metrics when the Ricci tensor is J-invariant, i.e. when equation (10.4) holds. Since $m = 1$, equation (10.4) is equivalent to

$$2A_1A_1'' - (A_1')^2 + b_j^2 = 0.$$

As seen above this equation admits two solutions; either

$$A_1 = bx + a$$

or

$$A_1 = e(x + c)^2 - \frac{1}{4}\frac{b^2}{e}.$$

If the first solution holds then the metric is Kahler. If the second solution holds then the metric will be Hermitian non-Kahler which is the case we will be concerned with in this section.

To simplify the calculations we make a slight change in variables and set $l = 2ec$ and $k = ec^2 - \frac{1}{4}\frac{b_1^2}{e} = \frac{1}{4e}(l^2 - b^2)$. We now have

$$A_1 = ex^2 + lx + k.$$

For the metric to have scalar curvature $S = \beta$ where β is a constant, the function h must satisfy

$$h = \frac{1}{ex^2 + lx + k} \int \left[\int \left(\frac{2}{ex^2 + lx + k} - \beta \right) (ex^2 + lx + k) dx \right] dx.$$

Furthermore, for the metric to be smooth at the special orbits $x = -1$ and $x = 1$, we require that $h(-1) = h(1) = 0$, $h'(-1) = 2$, and $h'(1) = -2$. Performing this double integral and imposing the boundary conditions implies that

$$S = 6,$$

$$k = \frac{1}{2} = \frac{1}{4e}(l^2 - b^2),$$

and

$$h = \frac{(1 - x)(1 + x)(ex^2 + 2lx + 1 + e)}{2ex^2 + 2lx + 1}.$$

The variable e is completely determined by b_1 and l, we can therefore write h as

$$h = \frac{(1-x)(1+x)((l^2-b^2)x^2+4lx+2+l^2-b^2)}{2((l+b)x+1)((l-b)x+1)}.$$

Furthermore, we find that

$$A_1 = \frac{l^2-b^2}{2}x^2 + lx + \frac{1}{2} = \frac{1}{2}((l+b)x+1)((l-b)x+1).$$

For the metric to be a positive definite metric on a Hirzebruch surface, we must have $h(x) > 0$ for $x \in (-1,1)$ and $A_i > 0$ for $x \in [-1,1]$. The condition $A_i > 0$ implies that

$$((l+b)x+1)((l-b)x+1) > 0.$$

For this to hold on the entire interval $x \in [-1,1]$ it is clear that both

$$(l+b)x+1 > 0 \tag{10.11}$$

and

$$(l-b)x+1 > 0. \tag{10.12}$$

Adding these two conditions together implies that $2lx+2 > 0$ for $x \in [-1,1]$. We deduce that

$$|l| < 1.$$

As mentioned at the beginning of this section, we can assume that $b \geq 0$. Evaluating inequality (10.11) at $x = -1$ and inequality (10.12) at $x = 1$ implies that

$$1 - l > b$$

and

$$1 + l > b$$

respectively. Adding these two equations together implies that

$$b < 1$$

which implies that

$$-q < 2.$$

As q is assumed to be a nonpositive integer, this leaves only two possibilities: $q = 0$, or -1.

Case 1: $q = 0$

If $q = 0$ then $b = 0$ as well and the manifold is $\mathbb{C}P^1 \times \mathbb{C}P^1$. The function h reduces to

$$h = \frac{(1-x)(1+x)(l^2x^2+4lx+2+l^2)}{2(lx+1)^2}$$

and A_1 becomes

$$A_1 = \frac{1}{2}(lx+1)^2.$$

This gives a constant scalar curvature Hermitian metric for all values of $|l| < 1$. Note that when $l = 0$, we recover a product metric on $\mathbb{C}P^1 \times \mathbb{C}P^1$. However, it is interesting to note that this is not the Kahler-Einstein metric as the two spheres are of different volumes.

Case 2: $q = -1$

If $q = -1$ then $b = \frac{1}{2}$ and the manifold is the blow-up of $\mathbb{C}P^2$ at one point, i.e. $\mathbb{C}P^2 \sharp \overline{\mathbb{C}P^2}$. The function h reduces to

$$h = \frac{(1-x)(1+x)((4l^2-1)x^2+16lx+7+4l^2)}{2((2l+1)x+2)((2l-1)x+2)}$$

and A_1 becomes

$$A_1 = \frac{1}{8}((2l+1)x+2)((2l-1)x+2).$$

By inspection, we see that we obtain a constant scalar curvature Hermitian metric on $\mathbb{C}P^2 \sharp \overline{\mathbb{C}P^2}$ when $|l| < \frac{1}{2}$.

When $l = 0$, we recover the Hermitian-Einstein metric (the Page metric) on $\mathbb{C}P^2 \sharp \overline{\mathbb{C}P^2}$ [57].

The above results prove the following proposition.

Proposition 10.4. *Let (M, J) be a Hirzebruch surface, $\mathbb{P}(\mathcal{O}_{\mathbb{C}P^1} \oplus \mathcal{O}_{\mathbb{C}P^1}(q))$ then M admits a $U(2)$-invariant compatible Hermitian metric, g, whose Ricci tensor is J-invariant and whose scalar curvature is constant if and only if $q = 0$ or -1. In both cases, the manifold admits a one-parameter family of such metrics. Furthermore, only one of these metrics, the product metric on $\mathbb{C}P^1 \times \mathbb{C}P^1$, is Kahler.*

10.3 A Curiosity

We end this section by noting an intriguing similarity between a cohomogeneity-one Hermitian-Einstein metric in dimension four and a cohomogeneity-one Kahler-Einstein metric in dimension six. Above, in our discussion of Extremal Kahler Fiberwise Toric manifolds in dimension six, we saw that the Kahler-Einstein metric found by Sakane on the manifold $\mathbb{P}(\mathcal{O}_{\mathbb{C}P^1 \times \mathbb{C}P^1} \oplus \mathcal{O}_{\mathbb{C}P^1 \times \mathbb{C}P^1}(1,-1))$ sits inside a two-parameter family of Extremal Kahler metrics on that manifold. In the previous section, we proved that the non-Kahler Hermitian-Einstein found by Page on $\mathbb{C}P^2 \sharp \overline{\mathbb{C}P^2} = \mathbb{P}(\mathcal{O}_{\mathbb{C}P^1} \oplus \mathcal{O}_{\mathbb{C}P^1}(1))$ sits inside a one-parameter family of non-Kahler constant scalar curvature Hermitian metrics with J-invariant Ricci tensor. Here, we want to note the striking similarity between these two metrics.

The Kahler-Einstein metric on $\mathbb{P}(\mathcal{O}_{\mathbb{C}P^1\times\mathbb{C}P^1}\oplus\mathcal{O}_{\mathbb{C}P^1\times\mathbb{C}P^1}(1,-1))$ can be written as

$$g=\begin{pmatrix}\frac{1}{h} & & & \\ & h & & \\ & & \frac{1}{2}(2+x)Id_2 & \\ & & & \frac{1}{2}(2-x)Id_2\end{pmatrix}$$

where

$$h=\frac{(1-x^2)(7-x^2)}{2(4-x^2)}.$$

The non-Kahler Hermitian-Einstein metric on $\mathbb{P}(\mathcal{O}_{\mathbb{C}P^1}\oplus\mathcal{O}_{\mathbb{C}P^1}(1))$ can be written as

$$g=\begin{pmatrix}\frac{1}{h} & & \\ & h & \\ & & \frac{1}{8}(2+x)(2-x)Id_2\end{pmatrix}$$

where

$$h=\frac{(1-x^2)(7-x^2)}{2(4-x^2)}.$$

It would be interesting to know if this is merely a coincidence or whether it is indicative of a more widespread correspondence between Kahler-Einstein and non-Kahler Hermitian-Einstein metrics.

10.4 Non-Kahler Hermitian Metrics with Constant Scalar Curvature in Higher Dimension

Above, we demonstrated that if the Ricci tensor of the complex manifolds is J-invariant on the manifolds being considered, i.e. if equation (10.4) holds then the scalar curvature equation can be integrated to

$$h=\frac{1}{\prod_{j=1}^m A_j^{\frac{d_j}{2}}}\int\left[\int\left(\sum_{j=1}^m\frac{d_j}{A_j}-S\right)\prod_{j=1}^m A_j^{\frac{d_j}{2}}dx\right]dx.$$

We want to find solutions to this equation in the case when $S=\beta$ where β is a constant. Unfortunately, we are not able to find all of the solutions to this problem because we do not know *all* of the solutions to equation (10.4). We can, however, give explicit solution for a subfamily of solutions to equation (10.4) obtained by adding the additional condition that the Riemannian curvature tensor is J-invariant in the following way: $R(J\cdot,J\cdot,J\cdot,J\cdot)=R(\cdot,\cdot,\cdot,\cdot)$. When the Riemannian curvature exhibits this invariance the functions A_i must satisfy $A_i=b_ix+a_i$ or $A_i=e_j(x+c_j)^2-\frac{1}{4}\frac{b_j^2}{e_j}$ as discussed above. In that case, we are able to perform the double integral and obtain explicit solutions as we did in the four dimensional case. (Note that in the case $m=1$, equation (10.4) is equivalent to the Riemannian tensor being J-invariant in this way, and that in dimension four $m=1$ automatically.) In those solutions, the function h will be rational in the variable x.

If all of the A_i are of the form $A_i=b_ix+a_i$ then the metric will be Kahler. As this case was considered in the section on fiberwise Kahler toric manifolds, we restrict our attention to that case

in which at least one of the functions A_i is quadratic in x. That is, there is some $k > 1$ such that A_j is quadratic for $1 \leq j \leq k$ and A_j is linear if $k+1 \leq j \leq m$. When A_j is quadratic, it will sometimes be convenient to write A_j as

$$A_j = e_j(x+c_j)^2 - \frac{1}{4}\frac{b_j^2}{e_j} = e_j(x + c_j + \frac{1}{2}\frac{b_j}{e_j})(x + c_j - \frac{1}{2}\frac{b_j}{e_j}) = e_j x^2 + l_j x + t_j$$

where $l_j = 2e_j c_j$ and $t_j = e_j c_j^2 - \frac{1}{4}\frac{b_j^2}{e_j}$.

10.4.1 Non-Compact Metrics of Constant Scalar Curvature

In this section, we restrict our attention to the search for non-Kahler Hermitian metrics with constant scalar curvature and J-invariant Riemannian tensor. We construct metrics of this type on the interval $[0, \infty)$. We restrict our attention to the case in which $A_j(0) > 0$ for all j. In that case, the manifold will be a complex line bundle over the product manifold $M_1 \times ... \times M_m$. For the metric to be smooth and positive the following must hold

- $h(0) = 0$ and $h'(0) = 2$
- $A_j = e_j x^2 + l_j x + t_j > 0$ for $x \in [0, \infty)$ which implies that $e_j > 0$, $l_j > 0$, and $t_j > 0$ for all $1 \leq j \leq k$
- $A_j = b_j x + a_j > 0$ for $x \in [0, \infty)$ which implies that $b_j \geq 0$ and $a_j \geq 0$ for all $k+1 \leq j \leq m$ and
- $h(x) > 0$ for $x \in (0, \infty)$.

(To obtain the third condition, note that $e_j > 0$ for the metric to be positive for large values of x, $t_j > 0$ for the $A_j(0) > 0$. Also, because the A_j can have only negative roots, $c_j > |\frac{1}{2}\frac{b_j^2}{e_j}|$. From the definition of l_j we deduce that $l_j > 0$.)

In the constant scalar curvature case, h must satisfy (10.9) with $S = \beta$ where β is a constant. We set $h(x) = \frac{P(x)}{Q(x)}$ where

$$P(x) = \frac{1}{\prod_{j=1}^{k}(e_j x^2 + l_j x + t_j)^{\frac{d_j}{2}} \prod_{j=k+1}^{m}(b_j x + a_j)^{\frac{d_j}{2}}} \times$$

$$\int \left[\int \left(\sum_{j=1}^{k} \frac{d_j}{e_j x^2 + l_j x + t_j} + \sum_{j=k+1}^{m} \frac{d_j}{b_j x + a_j} - \beta \right) \prod_{j=1}^{k}(e_j x^2 + l_j x + t_j)^{\frac{d_j}{2}} \prod_{j=k+1}^{m}(b_j x + a_j)^{\frac{d_j}{2}} dx \right] dx$$

and

$$Q(x) = \prod_{j=1}^{k}(e_j x^2 + l_j x + t_j)^{\frac{d_j}{2}} \prod_{j=k+1}^{m}(b_j x + a_j)^{\frac{d_j}{2}}.$$

Note that because we have assumed that $A_j(x) > 0$ for all j when $x \geq 0$ we see that $Q(x) > 0$ for $x \in [0, \infty)$. Both P and Q are polynomials in x because each of the d_j is even.

After performing the double integration, the polynomial P can be written schematically as

$$P = f + wx + ... - \beta \frac{\prod_{j=1}^{k} e_j^{\frac{d_j}{2}} \prod_{j=k+1}^{m} b_j^{\frac{d_j}{2}}}{(r+2)(r+1)} x^{r+2}$$

where f and w are the constants of integration and $r = \sum_{j=1}^{k} d_j + \sum_{k+1}^{m} \frac{d_j}{2}$. The smoothness conditions at $x = 0$ determine the two constants of integration. To this, we first note that $h(0) = 0$ implies that $P(0) = 0$ as $Q(0) > 0$. Therefore,

$$h(0) = 0 \Rightarrow P(0) = 0 \Rightarrow f = 0.$$

Next, because $P(0) = 0$, we have that $h'(0) = \frac{P'(0)}{Q(0)}$. Therefore,

$$h'(0) = 2 \Rightarrow \frac{P'(0)}{Q(0)} = 2 \Rightarrow w = 2 \prod_{j=1}^{k} t_j^{\frac{d_j}{2}} \prod_{j=k+1}^{m} a_j^{\frac{d_j}{2}}.$$

This implies that

$$P = 2 \Rightarrow w = 2 \prod_{j=1}^{k} t_j^{\frac{d_j}{2}} \prod_{j=k+1}^{m} a_j^{\frac{d_j}{2}} x + ... - \beta \frac{\prod_{j=1}^{k} e_j^{\frac{d_j}{2}} \prod_{j=k+1}^{m} b_j^{\frac{d_j}{2}}}{(r+2)(r+1)} x^{r+2}.$$

It remains only to check that $h(x) > 0$ for $x \in (0, \infty)$. This is equivalent to demanding that $P(x) > 0$ for $x \in (0, \infty)$. Now, as $x \to \infty$

$$P(x) \to -\beta \frac{\prod_{j=1}^{k} e_j^{\frac{d_j}{2}} \prod_{j=k+1}^{m} b_j^{\frac{d_j}{2}}}{(r+2)(r+1)} x^{r+2}.$$

If $\beta > 0$, then $P(x) \to -\infty$ as $x \to \infty$ and the metric will fail to be positive for large values of x. Therefore, $\beta \leq 0$. If $\beta \leq 0$, then we see that in the polynomial $P(x)$, all of the coefficients of the x^k are positive as $b_j \geq 0$, $a_j > 0$. $e_j > 0$, $l_j > 0$, $t_j > 0$, and $d_j > 0$ for all j. Therefore, $P(x) > 0$ for $x \in (0, \infty)$. All of the smoothness conditions and the positivity condition are satisfied for all values of $a_j > 0$, $e_j > 0$, and $c_j > |\frac{1}{2} \frac{b_j^2}{e_j}|$. As discussed above, it is the b_i's which determine the principal orbit type. So given a collection of b_i, we that there exists a scalar-flat metric for all values of a_j, e_j, and c_j, as well as a metric with scalar curvature, say, $\beta = -1$. This gives the following theorem:

Theorem 10.5. *Let (M, g) be complex line-bundle over $M_1 \times ... \times M_m$ endowed with a non-Kahler Hermitian metric as described above with $A_j = e_j(x + c_j)^2 - \frac{1}{4} \frac{b_j^2}{e_j}$ for $1 \leq j \leq k$ and $A_j = b_j x + a_j$ for $k+1 \leq j \leq m$ with $k > 1$. If $b_j > 0$ for $k+1 \leq j \leq m$, every such manifold admits a scalar-flat metric (S=0) as well as a negative scalar curvature metric (S=-1) for every value of $a_j > 0$, $e_j > 0$, and $c_j > \frac{1}{2} \frac{b_j}{e_j}$. That is, each such manifold admits an $m+k$-parameter family of metrics of scalar-flat metrics and an $m + k$-parameter family of metrics with scalar curvature equal to -1.*

When $A_j(0) = 0$ for some j, additional smoothness conditions must be satisfied. As in the Kahler case discussed above, we let $A_1(0) = 0$ and $A_i(0) > 0$ for $2 \leq i \leq m$. As in the Kahler case,

we know that $\mathbb{C}P^{\frac{d_1}{2}}$. If A_1 is linear then $A_1 = b_1 x$. If, on the other hand A_1 is quadratic, then $A_1 = e_1(x + c_1)^2 - \frac{1}{4}\frac{b_1^2}{e}$. The condition $A_1 = 0$ determines the constant c_1 to give

$$A_1 = e_1 x^2 + b_1 x.$$

For the function A_1 to be positive on the interval $x \in (0, \infty)$, both b_1 and e_1 must be greater than zero.

In both the linear and the quadratic cases, the derivative of A_1 at 0 implies that $A'(0) = b_1$. Since the first derivatives of A_1 at $x = 0$ agree in both the linear and quadratic cases, the additional smoothness condition must be the same. As in the Kahler case, we have

$$b_1 = \frac{2}{d_1 + 2}.$$

In either case, the total manifold will be a $\mathbb{C}^{\frac{d_1+2}{2}}$-bundle over $M_2 \times ... \times M_m$.

Theorem 10.6. *Let (M, g) be a $\mathbb{C}^{\frac{d_1+2}{2}}$-bundle over $M_2 \times ... \times M_m$ with $A_j = e_j(x + c_j)^2 - \frac{1}{4}\frac{b_j^2}{e_j}$ for $2 \leq j \leq k$, $A_j = b_j x + a_j$ with $b_j > 0$ or all $k + 1 \leq j \leq k$, and A_1 satisfies either $A_1 = \frac{2}{d_1+2}x$ or $A_1 = ex^2 + \frac{2}{d_1+2}$. Every such manifold admits a scalar-flat metric ($S = 0$) as well as a constant negative scalar curvature metric (S=-1) for every value of $a_j > 0$, $e_j > 0$ and $c_j > \frac{1}{2}\frac{b_j}{e_j}$ for all $j \neq 1$.*

Proof. The proof of this theorem is almost identical to the proof of the previous theorem. We therefore omit the details. □

10.4.2 Compact Metrics of Constant Scalar Curvature on $\mathbb{C}P^1$-Bundles

Here, we consider the case in which the metric is defined on the interval $x \in [0, 1]$ (with h vanishing at the endpoints) and $A_i(x) > 0$ for $x \in [0, 1]$ for all $1 \leq i \leq m$. In this case, the manifold M is a $\mathbb{C}P^1$-bundle over $M_1 \times ... \times M_m$. Above, we proved the existence of constant scalar curvature Kahler metrics on such manifolds when at least one of the b_i is positive and at least one is negative. Therefore, we restrict our attention to the case in which $b_i \geq 0$ (we could have equivalently considered the case in which all of the b_i are nonpositive) and at least one of the b_i, say b_1, is positive. We also assume that $m \geq 2$. We make this last assumption because, as we saw in our analysis of the four-dimensional case, there exist manifolds of this type which do not admit constant scalar curvature Hermitian metrics with J-invariant Riemannian curvature when $m = 1$.

Theorem 10.7. *Let (M, g, J) be a $\mathbb{C}P^1$-bundle over $M_1 \times ... \times M_m$ with $m \geq 2$ as described above. If $b_i \geq 0$ for $1 \leq i \leq m$ and $b_1 > 0$, then M admits at least a one-parameter family of non-Kahler Hermitian metrics of constant scalar curvature with J-invariant Riemannian curvature tensor.*

Proof. To simplify the calculations, we set $A_i = b_i x + a_i$ for $1 \leq i \leq m - 1$ and $A_m = e_m x^2 + l_m x + t_m = ex^2 + lx + t$. Because one of the terms is quadratic, the metric will be non-Kahler.

Furthermore, we can assume without loss of generality that $b_j > 0$ for $2 \leq j \leq m-1$ since the metric would be a product if one of these b_j where to equal zero. For the metric to be smooth and positive definite the following must hold:

- $h(0) = 0$ and $h'(0) = 2$
- $h(1) = 0$ and $h'(1) = -2$
- $A_i = b_i x + a_i > 0$ for all $1 \leq i \leq m$ for $x \in [0,1]$ which implies that $a_i > 0$
- $A_m = ex^2 + lx + t > 0$ for $x \in [0,1]$ and
- $h(x) > 0$ for $x \in (0,1)$.

Let $c_i = \frac{a_i}{b_i} > 0$ for $1 \leq i \leq m-1$. Note that $c_i > 0$ because b_i is assumed to be greater than zero. We have

$$\prod_{j=1}^{m} A_i^{\frac{d_j}{2}} = \left(\prod_{j=1}^{m-1} b_j^{\frac{d_j}{2}} \right) \left(\prod_{j=1}^{m-1} (x + c_j)^{\frac{d_j}{2}} \right) (ex^2 + lx + t)^{\frac{d_m}{2}}$$

and

$$\sum_{j=1}^{m} \frac{d_j}{A_j} = \left(\sum_{j=1}^{m-1} \frac{\frac{d_j}{b_j}}{x + c_j} \right) + \frac{d_m}{ex^2 + lx + t}.$$

By changing the value of the a_i we can set

$$c_j = c > 0$$

for all $1 \leq j \leq m-1$. This implies that, in the constant scalar curvature case $S = \beta$,

$$h = \frac{1}{(x+c)^p (ex^2 + lx + t)^q} \int \left[\int \left(\frac{r}{x+c} + \frac{2q}{ex^2 + lx + t} - \beta \right) (x+c)^p (ex^2 + lx + t)^q dx \right] dx$$

where $p = \sum_{j=1}^{m-1} \frac{d_j}{2}$, $q = \frac{d_m}{2}$, and $r = \sum_{j=1}^{m-1} \frac{d_j}{b_j}$ which are all greater than zero.

Set $h(x) = \frac{T(x)}{L(x)}$ where

$$T(x) = \int \left[\int \left(\frac{r}{x+c} + \frac{2q}{ex^2 + lx + t} - \beta \right) (x+c)^p (ex^2 + lx + t)^q dx \right] dx$$

and

$$L(x) = (x+c)^p (ex^2 + lx + t)^q.$$

Schematically, we can write T as

$$T = F_{p+q+4} x^{p+q+4} + ... + F_w x^w + ... + Ex + F$$

where E and F are the constants of integration. It is straightforward to check that the smoothness conditions $h(0) = 0$ and $h'(0) = 2$ are satisfied when $F = 0$ and $E = 2c^p t^q$.

We will see that the two smoothness conditions at $x = 1$ will determine β and will place a condition on the constants c, l, and t. To obtain the constant scalar curvature metrics, it will be helpful to group the terms by the order of c and t. Doing this, we see that

$$T = [2c^p t^q x + c^{p-1}t^{q-1}(rt + 2qc)\frac{x^2}{2} + ...] - \beta[c^p t^q \frac{x^2}{2} + c^{p-1}t^{q-1}(pt + lqc)\frac{x^3}{6} + ...]$$

and

$$L = c^p t^q + c^{p-1}t^{q-1}(pt + lqc)x + ...$$

The condition $h(1) = 0$ implies that

$$\beta = \frac{2c^p t^q + \frac{1}{2}c^{p-1}t^{q-1}(rt + 2qc) + ...}{\frac{1}{2}c^p t^q + \frac{1}{6}c^{p-1}t^{q-1}(pt + lqc) + ...}.$$

As $c \to \infty$ and $t \to \infty$, we see that $\beta \to 4$.

The final smoothness condition states that $h'(1) = -2$ which is equivalent to $\frac{T'(1)}{L(1)} = -2$. This implies that

$$\beta = \frac{4c^p t^q + c^{p-1}t^{q-1}(rt + 2qc) + 2c^{p-1}t^{q-1}(pt + lqc) + ...}{c^p t^q + \frac{1}{2}c^{p-1}t^{q-1}(pt + lqc) + ...}.$$

Equating these two expressions for β implies that

$$\frac{2}{3}c^{2p-1}t^{2q-1}(pt + lqc) + K(c, t) = 0$$

where $K = K(c, t)$ is a polynomial in c and t in which each of the terms in K is of the form $K_{\eta\xi}c^\eta t^\xi$ such that $\eta \leq 2p$, $\xi \leq 2q$, $\eta + \xi < 2p + 2q - 1$, and the $K_{\eta\xi}$ are constants depending on the other terms. This implies that

$$B(c, t) \equiv pt + lqc + \frac{3}{2}\frac{K(c, t)}{c^{2p-1}t^{2q-1}} = 0. \tag{10.13}$$

Note that as c and t approach positive infinity, $\frac{3}{2}\frac{K(c,t)}{c^{2p-1}t^{2q-1}}$ approaches a constant. Recall that $A_m = e_m x^2 + 2e_m c_m x + e_m c_m^2 - \frac{1}{4}\frac{b_j^2}{e_m} = ex^2 + lx + t$. Letting $e_m = e$ be fixed positive number, equation (10.13) implies that

$$e[p(c_m^2 - \frac{1}{4}\frac{b_m^2}{e^2}) + 2qcc_m] + \frac{3}{2}\frac{K(c, t)}{c^{2p-1}t^{2q-1}} = 0.$$

As $c_m \to -\infty$ and $c \to \infty$, this equation, after dividing by ec_m, implies that

$$-pc_m - 2qc + C(c, c_m) = 0$$

where $C(c, c_m) \to 0$ as $c \to \infty$ and $c_m \to -\infty$. As p and q are positive, the above equation must (by an intermediate value argument similar to those used above) have a solution for $-c_m$ and c arbitrarily large. As $c_m \to \infty$ and $c \to \infty$, $A_m \to \infty$ and $A_i \to \infty$ for $1 \leq i \leq m - 1$. Furthermore, h will be positive in $(0, 1)$ for large values of c and c_m^2.

What we have demonstrated is that we can always solve the smoothness conditions from $-c_m$ and c arbitrarily large. Moreover, the resulting function h will be positive on $x \in (0, 1)$. This proves the theorem. □

10.4.3 Compact Metrics of Constant Scalar Curvature on $\mathbb{C}P^{\frac{d_1+2}{2}}$- Bundles

We now consider the case in which $A_1(0) = 0$. For the metric to be smooth, we know that $M_1 \cong \mathbb{C}P^{\frac{d_1}{2}}$, $b_1 = \frac{2}{d_1+2}$ and $A_1 = e_1x^2 + b_1x$ or $A_1 = b_1x$. In this case the manifold M is a $\mathbb{C}P^{\frac{d_1+2}{2}}$-bundle over $M_2 \times ... \times M_m$.

Above, we demonstrated that such manifolds admit constant scalar curvature Kahler metrics when $m = 1$ or when at least one of the b_i is positive and at least one is negative for $2 \leq i \leq m$. In this section we ask which of the remaining manifolds admits a constant scalar curvature Hermitian metric with J-invariant Riemannian curvature.

Theorem 10.8. *Let (M, g, J) be a $\mathbb{C}P^{\frac{d_1+2}{2}}$-bundle over $M_2 \times ... \times M_m$ with $m \geq 3$ as described above. If $b_i \geq 0$ for $3 \leq i \leq m$ and $b_2 > 0$, then M admits at least a one-parameter family of non-Kahler Hermitian constant scalar curvature metrics with J-invariant Ricci tensor. The following theorem provides a partial answer to this question.*

Proof. As in the proof of the theorem of the previous section we set $A_i = b_ix + a_i$ for $2 \leq j \leq m-1$ and set $A_m = e_m(x + c_m)^2 - \frac{1}{4}\frac{b_m^2}{e_m}$. The presence of a quadratic term prevents the metric from being Kahler. We take A_1 to be either quadratic or linear. The details of this proof follow almost identically to those of the previous theorem. For brevity, we omit the details. □

Chapter 11

Integrating the Scalar Curvature Equation on a Cohomogeneity-One Manifold

In this thesis, we were able to integrate the scalar curvature equation on a Kahler manifold of cohomogeneity-one under a semisimple Lie group in the monotypic case. We were also able to integrate the scalar curvature equation in a non-Kahler Hermitian case similar to the cohomogeneity-one Kahler case. In this section, we describe an integration of the scalar curvature equation which applies to a large class of cohomogeneity-one manifolds in the absence of a complex structure or special holonomy.

Let (M^n, g) be a manifold of cohomogeneity-one under the action of a compact, simply-connected Lie group G with connected isotropy subgroup K. This implies that G/K must be simply-connected. The Lie algebra $\mathfrak{g}$ decomposes as

$$\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}_0 \oplus ... \oplus \mathfrak{p}_m$$

where the $\mathfrak{p}_i$ are irreducible $\mathrm{Ad}(K)$-invariant subspaces. Let d_i denote the dimension of the summand $\mathfrak{p}_i$. We begin with two assumptions.

Assumption One:

The $\mathfrak{p}_i$ are distinct.

We restrict ourselves to the monotypic case because the metric, as it is automatically diagonalizable, is more easily described.

Assumption Two:

$$\dim\mathfrak{p}_0 = 1.$$

This is a more serious restriction; however, assumption two does hold in a variety of interesting cases. For example, if the cohomogeneity-one metric admits a compatible G-invariant almost-complex structure then there must be an invariant one-dimensional subspace in the isotropy representation. This can be seen by setting $\frac{\partial}{\partial t}$ to be the unit tangent vector to the horizontal distribution. The image of the vector field under the almost-complex structure, $J\frac{\partial}{\partial t}$, belongs to the tangent space to the vertical principal orbits. The vector field $J\frac{\partial}{\partial t}$ is invariant under the action of G because both $\frac{\partial}{\partial t}$ and J are invariant under the group action. Therefore, $J\frac{\partial}{\partial t}$ spans an invariant one-dimensional subspace. Of course, assumption two can hold even in the absence of an almost-complex structure.

Because K is connected and $\mathfrak{p}_0$ is one-dimensional we see that $\mathfrak{p}_0$ is in fact a trivial representation.

Lemma 11.1. *Let (M, g) be a cohomogeneity-one manifold under the action of a compact connected Lie group G with connected isotropy subgroup K satisfying assumptions one and two. On such a manifold,*

$$[\mathfrak{p}_0, \mathfrak{p}_i] \subseteq \mathfrak{p}_i \tag{11.1}$$

for all i.

Proof. Clearly, $[\mathfrak{p}_0, \mathfrak{p}_0] = 0$ as $\mathfrak{p}_0$ is one-dimensional. Furthermore, since K acts trivially on $\mathfrak{p}_0$, we see that $[\mathfrak{k}, \mathfrak{p}_0] = 0$. This implies that $\mathrm{ad}(\mathfrak{p}_0)$ acts on $\mathfrak{p}_1 \oplus ... \oplus \mathfrak{p}_m$ by K-morphisms. Because we have assumed that all of the $\mathfrak{p}_i$ are distinct, Schur's Lemma implies that $\mathrm{ad}(\mathfrak{p}_0) : \mathfrak{p}_i \to \mathfrak{p}_i$ for all $1 \leq i \leq m$. □

Under these assumptions we can write the metric, g, as

$$g = dt^2 + hQ|_{\mathfrak{p}_0} + \bigoplus_{i=1}^{m} A_i Q|_{\mathfrak{p}_i}$$

where Q is some $\mathrm{Ad}(K)$-invariant background metric on $\mathfrak{p} = \mathfrak{p}_0 \oplus ... \oplus \mathfrak{p}_m$. Let X_0 be an orthonormal basis for $Q|_{\mathfrak{p}_0}$ and let $\{Y_j^i\}_{j=1,...,d_i}$ be an orthonormal basis for $Q|_{\mathfrak{p}_i}$. Note also that h and the A_i are positive functions of t.

Let $x = x(t)$ be defined by the equation

$$\frac{1}{h}dx^2 = dt^2.$$

This implies that $\frac{dx}{dt} = \sqrt{h}$. We can now write the metric as

$$g = \begin{pmatrix} \frac{1}{h} & & & & \\ & h & & & \\ & & A_1 Id_{d_1} & & \\ & & & ... & \\ & & & & A_m Id_{d_m} \end{pmatrix}$$

relative to basis $(\frac{\partial}{\partial x}, X_0, Y_1^1, ..., Y_{d_m}^m)$.

The Einstein condition on this metric can be written schematically as

$$r\left(\frac{\partial}{\partial x}, \frac{\partial}{\partial x}\right) = \lambda g\left(\frac{\partial}{\partial x}, \frac{\partial}{\partial x}\right),$$

$$r(X_0, X_0) = \lambda g(X_0, X_0),$$

and

$$r(Y_k^i, Y_k^i) = \lambda g(Y_k^i, Y_k^i)$$

for all i where λ is the Einstein constant. Note that which $1 \leq k \leq d_i$ we choose is immaterial. In terms of the functions h and A_i it is straightforward to calculate that these equations become

$$-\frac{1}{2}h'' - h'\left(\sum_{j=1}^m \frac{d_j}{4}\frac{A_j'}{A_j}\right) + h\left[\sum_{j=1}^m \frac{d_j}{4}\left(\frac{A_j'}{A_j}\right)^2 - \sum_{j=1}^m \frac{d_j}{2}\frac{A_j''}{A_j}\right] = \lambda, \tag{11.2}$$

$$-\frac{1}{2}h'' - h'\left(\sum_{j=1}^m \frac{d_j}{4}\frac{A_j'}{A_j}\right) + \frac{1}{h}\hat{r}(X_0, X_0) = \lambda, \tag{11.3}$$

and

$$-\frac{1}{2}h'\frac{A_i'}{A_i} + h\left[-\frac{1}{2}\frac{A_i''}{A_i} + \frac{1}{2}\left(\frac{A_i'}{A_i}\right)^2 - \frac{A_i'}{A_i}\left(\sum_{j=1}^m \frac{d_j}{4}\frac{A_j'}{A_j}\right)\right] + \frac{1}{A_i}\hat{r}(Y_k^i, Y_k^i) = \lambda \tag{11.4}$$

for all $1 \leq i \leq m$. Recall that $\hat{r}$ is the Ricci curvature of the principal orbits as described in Chapter 2. More precisely, given a point x in the quotient space M/G, $\hat{r}|_x$ is the Ricci tensor of the principal orbit located at x.

We note that equations (11.2) and (11.3) together imply that

$$\frac{1}{h^2}\hat{r}(X_0, X_0) = \sum_{j=1}^m \frac{d_j}{4}\left(\frac{A_j'}{A_j}\right)^2 - \sum_{j=1}^m \frac{d_j}{2}\frac{A_j''}{A_j}. \tag{11.5}$$

The scalar curvature equation can be written as

$$S = -h'' - h'\left(\sum_{j=1}^m d_j \frac{A_j'}{A_j}\right) + h\left[-\sum_{j=1}^m d_j \frac{A_j''}{A_j} + \frac{3}{4}\sum_{j=1}^m d_j\left(\frac{A_j'}{A_j}\right)^2 - \left(\sum_{j=1}^m \frac{d_j}{2}\frac{A_j'}{A_j}\right)^2\right] + \hat{S} \tag{11.6}$$

where

$$\hat{S} = \frac{1}{h}\hat{r}(X_0, X_0) + \sum_{j=1}^m \frac{d_j}{A_j}\hat{r}(Y_k^j, Y_k^j).$$

is the scalar curvature of the homogeneous principal orbits and S is the scalar curvature of the total cohomogeneity-one manifold M. Specifically, for x a point in the quotient space, $\hat{S}|_x$ is the scalar curvature of the principal orbit located at the point x.

In [58], Wang and Ziller give a formula (1.3 on page 181) for the scalar curvature of a homogeneous manifold. As each of the principal orbits of M is a homogeneous manifold, G/K, we can use this formula to determine the form of $\hat{S}$. From that formula we see that $\hat{S}$ can be written in terms of the h and the A_i. More precisely, we can decompose $\hat{S}$ by setting

$$\hat{S} = S_h + S^*$$

where S_h includes all of the terms involving h and S^* does not include any terms with h in them but only terms involving the A_i. Because $\mathfrak{p}_0$ is trivial, we have the following proposition:

Proposition 11.2. *For (M, g) a cohomogeneity-one manifold satisfying assumptions one and two as above, we have that*

$$S_h = -h \sum_{j=1}^{m} \frac{d_j}{4} \frac{b_j^2}{A_j^2}$$

for some constants b_j.

Proof. Using lemma (11.1) and the fact that $\mathfrak{p}_0$ is trivial, we see from the formula in [58] that the only factors of $\hat{S}$ which involve the function h are of the form $-h\frac{d_j}{4}\frac{b_j^2}{A_j^2}$ for $1 \leq j \leq m$. □

The form of the function S^* will vary depending on the groups G and K. However, we stress that S^* does *not* involve the function h.

We next consider the term $\frac{1}{h}\hat{r}(X_0, X_0)$.

Proposition 11.3. *Let (M, g) be a cohomogeneity-one manifold satisfying assumptions one and two. We have that*

$$\frac{1}{h}\hat{r}(X_0, X_0) = h \left(\sum_{j=1}^{m} \frac{d_j}{4} \frac{b_j^2}{A_j^2} \right)$$

for some constants b_i.

Proof. Again, this result follows from lemma (11.1) and the fact the $\mathfrak{p}_0$ is trivial. Given these two facts, along with the discussion in [58] and equation 7.38 of [6], it is straightforward to calculate that this proposition holds. □

From the above we deduce immediately that

$$S_h = -\frac{1}{h}\hat{r}(X_0, X_0).$$

Furthermore, proposition (11.3) and equation (11.5) imply that

$$\sum_{j=1}^{m} d_j \mu_j = 0$$

where

$$\mu_j = 2\frac{A_j''}{A_j} - \left(\frac{A_j'}{A_j}\right)^2 + \frac{b_j^2}{A_j^2}.$$

We pause now to discuss the significance of equation (11.5). We define a map $\tilde{J}$ by the action $\tilde{J}\frac{\partial}{\partial x} = \frac{1}{h}X_0$ and $\tilde{J}X_0 = -h\frac{\partial}{\partial x}$. The map $\tilde{J}$ can be written as

$$\tilde{J} = \begin{pmatrix} & -h & & & \\ \frac{1}{h} & & & & \\ & & 0 & & \\ & & & ... & \\ & & & & 0 \end{pmatrix}$$

relative to basis $(\frac{\partial}{\partial x}, X_0, Y_1^1, ..., Y_{d_m}^m)$. Clearly, the metric g is invariant under the action of $\tilde{J}$, i.e. $g(\cdot,\cdot) = g\left(\tilde{J}\cdot, \tilde{J}\cdot\right)$. However, the Ricci tensor, r, is *not* automatically invariant under the action of $\tilde{J}$. In fact,

$$r(\cdot,\cdot)|_{\langle\frac{\partial}{\partial x}\rangle\oplus\mathfrak{p}_0} = r\left(\tilde{J}\cdot, \tilde{J}\cdot\right)|_{\langle\frac{\partial}{\partial x}\rangle\oplus\mathfrak{p}_0} \Leftrightarrow (11.5) \text{ holds} \Leftrightarrow \sum_{j=1}^{m} d_j\mu_j = 0.$$

We also note that the metric g and the map $\tilde{J}$ define a 2-form of rank two such that $\tilde{\omega} = g\tilde{J}$, i.e. $\tilde{\omega}(\cdot,\cdot) = g(\tilde{J}\cdot,\cdot)$. This can be written as

$$\tilde{\omega} = \begin{pmatrix} & -1 & & & \\ 1 & & & & \\ & & 0 & & \\ & & & ... & \\ & & & & 0 \end{pmatrix}$$

relative to basis $(\frac{\partial}{\partial x}, X_0, Y_1^1, ..., Y_{d_m}^m)$. Alternatively, we could write $\tilde{\omega}$ as

$$\tilde{\omega} = dx \wedge \theta_0$$

where θ_0 is the invariant dual 1-form to the vector field X_0 such that $\theta_0(X_0) = 1$.

We are now prepared to state and prove this section's main theorem.

Theorem 11.4. *Let (M, g) be a cohomogeneity-one manifold satisfying assumptions one and two. The function h can be written as*

$$h = \frac{1}{V^{\frac{1}{2}}}\int\left[\int (S^* - S)V^{\frac{1}{2}}dx\right]dx, \tag{11.7}$$

where $V^{\frac{1}{2}} = \prod_{j=1}^m A_j^{\frac{d_j}{2}}$, if and only if

$$\sum_{j=1}^{m} d_j\mu_j = 0.$$

Proof. Differentiating equation (11.7) implies that

$$S = -h'' - h'\left(\sum_{j=1}^{m} d_j\frac{A_j'}{A_j}\right) + h\left[-\sum_{j=1}^{m}\frac{d_j}{2}\frac{A_j''}{A_j} + \sum_{j=1}^{m}\frac{d_j}{2}\left(\frac{A_j'}{A_j}\right)^2 - \left(\sum_{j=1}^{m}\frac{d_j}{2}\frac{A_j'}{A_j}\right)^2\right] + S^*.$$

Comparing this to equation (11.6) implies that

$$S_h = h\left[-\sum_{j=1}^{m}\frac{d_j}{4}\left(\frac{A_j'}{A_j}\right)^2+\sum_{j=1}^{m}\frac{d_j}{2}\frac{A_j''}{A_j}\right].$$

Since $S_h = -\frac{1}{h}\hat{r}(X_0, X_0)$, this is equivalent to $\sum_{j=1}^{m} d_j\mu_j = 0$. □

Put another way, the scalar curvature equation can be integrated to give (11.7) if and only if the Ricci tensor on $\langle\frac{\partial}{\partial x}\rangle \oplus \mathfrak{p}_0$ is invariant under the map $\tilde{J}$. In the Einstein case, the Ricci tensor must be so invariant and, therefore, equation (11.7) *must* hold in the Einstein case.

Bibliography

[1] Abreu, Miguel, *Kahler Geometry of Toric Manifolds in Symplectic Coordinates*, (2003), Field Institute Communications, 35, 1-24.

[2] Abreu, M., *Kahler Geometry of Toric Varieties and Extremal Metrics*, (1998), Internat. J. Math., 9, 641-651.

[3] Apostolov, Vestislav; Armstrong, John; Draghici, Tedi, *Local Models and Integrability of Certain Almost Kahler 4-Manifolds*, (2002), Math. Ann., 323, no. 4, 633-666.

[4] Apostolov, V.; Gauduchon, P., *The Riemannian Goldberg-Sachs Theorem*, (1997), Int. J. Math., 8, 421-439.

[5] Bérard-Bergery, Lionel, *Sur des Nouvelles Variétés Riemanniennes d'Einstein*, (1981), Publication de l'Institut É. Cartan, 4, 1-60.

[6] Besse, A. L., *Einstein Manifolds*, (1987), Spring-Verlag, Berlin.

[7] Bredon, G. E., *Introduction to Compact Transformation Groups*, (1972), Academic Press, New York, USA.

[8] Bouguignon, J. P., *Une stratification de l'espace des structures riemanniennes*, (1975), Compositio Math., 30, 1-41.

[9] Calabi, E., it A Construction of Nonhomogeneous Einstein Metrics, (1975), Differential Geometry (Proc. Sympos. Pure Math., Vol. XXVII, Stanford Univ., Stanford, Calif., 1973), Part 2, Amer. Math. Soc., Providence, R.I., 17-24.

[10] Calabi, E., *Extremal Kahler Metrics*, (1982), Seminar on Differential Geometry, ed. S. T. Yau, Annals of Math. Studies, 102, Princeton University Press, 259-290.

[11] Candelas, P.; de la Ossa, X., *Comments on Conifolds*, (1990), Nucl. Phys. B, 342, 246-268.

[12] Dancer, A. S.; Strachan, I. A. B., *Einstein Metrics on Tangent Bundles of Spheres*, (2002), Class. Quantum Grav., 19, 4663-4670.

[13] Dancer, A. S.; Strachan, I. A. B., *Kahler-Einstein Metrics with $SU(2)$ Action*, (1994), Proc. Camb. Phil. Soc., 115, 513-525.

[14] Dancer, Andrew; Wang, McKenzie, *The Cohomogeneity One Einstein Equations from the Hamiltonian Viewpoint*, (2000), J. Reine AnGew. Math., 524, 97-128.

[15] Dancer, Andrew; Wang, McKenzie, *Kahler-Einstein Metrics of Cohomogeneity One*, (1998), Math. Ann., 312, 503-526.

[16] Delzant, T., *Hamiltoniens Périodiques et Image Convexe de l'Application Moment*, (1988), Bull. Soc. Math. France, 116, 315-339.

[17] Derdzinski, Andrzej, *Self-Dual Kahler Manifolds and Einstein Manifolds of Dimension Four*, (1983), Comp. Math., 49, 405-433.

[18] Donaldson, S., *Scalar Curvature and Stability of Toric Varieties*, (2002), J. Diff. Geom., 62, 289-349.

[19] Donaldson, S.; Kronheimer, P. B., The Geometry of Four-Manifolds, (1997), Oxford University Press.

[20] Falcitelli, M.; Farinola, A.; Salamon, S.; *Almost-Hermitian Geometry*, (1994), Differential Geom. Appl. 4, no. 3, 259-282.

[21] Freedman, M., *On the Topology of Four-Manifolds*. (1982), J. Diff. Geom., 17, 357-454.

[22] Fulton, William; Harris, Joe, *Representation Theory: A First Course*, (1991), Graduate Texts in Mathematics, Springer-Verlag, NY, USA.

[23] Fulton, William, *Introduction to Toric Varieties*, (1993), Annals of Mathematics Studies, 131, Princeton University Press.

[24] Futaki, A., *An Obstruction to the Existence of Einstein-Kahler Metrics*, (1983), Inv. Math., 73, 437-443.

[25] Futaki, A., *Kahler-Einstein Metrics and Integral Invariants*, Lecture Notes in Mathematics, 1314, Springer-Verlag.

[26] Gibbons, G. W.; Hawking, S. W., *Classification of Gravitational Instanton Symmetries*, (1979), Commun. Math. Phys., 66, 291-310.

[27] Gray, Alfred, *Curvature Identities for Hermitian and Almost Hermitian Manifolds*, (1976), Tohoku Math. J., (2) 38, no.4, 601-612.

[28] Guillemin, V., *Moment Maps and Combinatorial Invariants of Hamiltonian T^n-spaces*, (1994), Progress in Math. 122, Birkhauser.

[29] Hitchin, N. J., *The Moduli Space of Complex Lagrangian Submanifolds Sir Michael Atiyah: A Great Mathematician of the Twentieth Century*, (1999), Asian J. Math., 3, 77-91.

[30] Hitchin, N., *On Compact Four-Dimensional Einstein Manifolds*, (1974), J. Diff. Geom., 9, 435-442.

[31] Hwang, Andrew D.; Simanca, Santiago R., *Extremal Kahler Metrics on Hirzebruch Surfaces which are Locally Conformally Equivalent to Einstein Metrics*

[32] Jensen, G. R., *Homogeneous Einstein Spaces of Dimension Four*, (1969), J. Diff. Geom., 3, 309-349.

[33] Jost, Jürgen, *Riemannian Geometry and Geometrical Analysis*, (1998), Universitext, Springer-Verlag, NY, USA.

[34] Joyce, Dominic D., *Compact Manifolds with Special Holonomy*, (2000), Oxford University Press.

[35] Joyce, Dominic D., *Explicit Construction of Self-Dual Four-Manifolds*, (1995), Duke Math. J., 77, 519-552.

[36] Kazdan, J. L.; Warner, F. W., *Curvature Functions for Open 2-Manifolds*, (1974), Ann. of Math., 99, 203-219.

[37] Kerr, Megan, *New Examples of Homogeneous Einstein Metrics*, (1998), Michigan Math. J., 45, 115-134.

[38] Kobayashi, S.; Nomizu, K., *Foundations of Differential Geometry, Vol. II*, (1969), Interscience Wiley, New York, USA.

[39] Kundt, W.; Trumper, M., *Orthogonal Decomposition of Axi-Symmetric Stationary Space-Times*, (1966), Z. Phys., 192, 419-422.

[40] LeBrun, C., *Einstein Metrics on Complex Surfaces*, (1997), Lecture Notes in Pure and Applied Mathematics, Vol. 184, Eds. Anderson et al, Geometry and Physics, New York, 167-176.

[41] LeBrun, Claude; Wang, McKenzie, *Survey in Differential Geometry: Essays on Einstein Manifolds*, (1999), Supplement to the J. Diff. Geom., International Press, Boston, USA.

[42] Lee, John M., *Riemannian Manifolds: An Introduction to Curvature*, (1997), Graduate Texts in Mathematics, Springer-Verlag, NY, USA.

[43] Mabuchi, T., *Einstein-Kahler Forms, Futaki Invariants and Convex Geometry on Toric Fano Varieties*, (1987), Osaka J. Math., 24, 705-737.

[44] Madsen, A. B., *Compact Einstein-Weyl Manifolds with Large Symmetry Group*, (1995), Ph. D. thesis, Odense Univ.

[45] Mason, L.J.; Woodhouse, N. M., *Integrability, Self-Duality, and Twistor Theory*, (1996), Oxford University Press.

[46] Matsushima, Y., *Sur la Structure du Groupe d'Homéomorphismes Analytiques d'une Certaine Variété Kaehlérienne*, (1957), Nagoya Math. J., 11, 145-150.

[47] Obata, M., *The Conjectures on Conformal Transformations of Riemannian Manifolds*, (1961), Math. Soc., 12, 889-891.

[48] O'Neill, B., *The Fundamental Equations of a Submersion*, (1966), Mich. Math. J., 13, p. 459-469.

[49] Orlik, Peter; Raymond, Frank, *Actions of the Torus on Four Manifolds I*, (1970), Trans. Amer. Math. Soc., 152, 531-559.

[50] Page, Don, *A Compact Rotating Gravitational Instanton*, (1978), Physics Letters, 79B, 235-238.

[51] Pedersen, Henrik; Swann, Andrew; *Riemannian Submersions, Four-Manifolds, and Einstein-Weyl Geometry*, (1991), Proc. London. Math. Soc. (3), 66, 381-399.

[52] Petersen, Peter, *Riemannian Geometry*, (1999), Graduate Texts in Mathematics, Springer-Verlag, NY, USA.

[53] Poor, W. A., *Differential Geometric Structures*, (1981), McGraw-Hill, New York, USA.

[54] Sakane, Y., *Examples of Compact Einstein Kahler Manifolds with Positive Ricci Tensor*, (1986), Osaka Math. J., 23, 585-617.

[55] Siu, Y., *The Existence of Kahler-Einstein Metrics on Manifolds with Positive Anti-Canonical Line Bundle and Suitable Finite Symmetry Group*, (1988), Ann. Math., 127, 585-627.

[56] Stenzel, M. B., *Ricci-Flat Metrics on the Complexification of a Compact Rank One Symmetric Space*, (1993), Manuscr. Math., 80, 151-163.

[57] Wang, Jun; Wang, McKenzie, *Einstein Metrics on S^2-Bundles*, (1998), Math. Ann., 310, 497-526.

[58] Wang, M. Y.; Ziller, W., *Einstein Metrics in Principal Torus Bundles*, (1990), J. Diff. Geom., 31, 215-248.

[59] Ward, R. S., *Stationary Axisymmetric Space-Times: A New Approach*, (1983), Gen. Rel. Grav., 15, 105-9.

[60] Warner, Frank W., *Foundations of Differentiable Manifolds and Lie Groups*, (1983), Graduate Texts in Mathematics, Springer-Verlag, NY, USA.

[61] Tian, G.; Yau, S., *Kahler-Einstein Metrics on Complex Surfaces with* $c_1 > 0$, (1987), Comm. Math. Phys., 112, 175-203.